



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

3526

35



QL

561

N9

B86

CATALOGUE
OF
DIURNAL LEPIDOPTERA
OF THE FAMILY
SATYRIDÆ

IN THE
COLLECTION

OF THE
BRITISH MUSEUM. *(Nat. Hist) Dept
of Zool.*

BY
ARTHUR GARDINER BUTLER, F.L.S., F.Z.S., &c.

LONDON:
PRINTED BY ORDER OF THE TRUSTEES.
1868.



PRINTED BY TAYLOR AND FRANCIS,
RED LION COURT, FLEET STREET.

026 by 16 7-8.

PREFACE.

THE object of the present Catalogue is to give a complete list of all the genera and species of Satyridæ known to exist in the cabinets of European and American entomologists.

The localities given chiefly apply to the specimens in the National Collection; the letters B.M. after the species denote the specimens now contained in the British Museum; and the absence of these *letters* indicates the species which are desiderata, and therefore desirable to be procured for the Collection.

JOHN EDWARD GRAY.

British Museum,
Feb. 20, 1868.

Revised 7-20-34 K.M.P.



EXPLANATION OF THE PLATES.

I.

Fig.	Page
1. <i>Taygetis excavata</i> , <i>Butl.</i>	8
2. — <i>erubescens</i> , <i>Butl.</i>	10
3. — <i>armillata</i> , <i>Butl.</i>	8
4. <i>Euptychia rustica</i> , <i>Butl.</i>	32
5. <i>Taygetis Euptychidia</i> , <i>Butl.</i>	13
6. — <i>nymphosa</i> , <i>Butl.</i>	10
7. — <i>nympha</i> , <i>Butl.</i>	9

II.

1. <i>Melanitis Gnophodes</i> , <i>Butl.</i>	5
2. <i>Ceneis gigas</i> , <i>Butl.</i>	161
3. <i>Lethe arcuata</i> (= <i>arete</i> , <i>Cramer</i>)	114
4. <i>Erites elegans</i> , <i>Butl.</i>	147
5. <i>Paramecera Xicaque</i> , <i>Reakirt</i>	99
6. <i>Erebia mantoides</i> , <i>Butl.</i>	87
7. — <i>Hispania</i> , <i>Butl.</i>	86
8. — <i>fasciata</i> , <i>Butl.</i>	92
9. — <i>epipsodea</i> , <i>Butl.</i>	80
10. <i>Ceneis assimilis</i> , <i>Butl.</i>	163
11. <i>Tetraphlebia plumbeola</i> , <i>Butl.</i>	95
12. <i>Leptoneura cassina</i> , <i>Butl.</i>	72

III.

1. <i>Mycalesis saga</i> , <i>Butl.</i>	130
2. — <i>vulgaris</i> , <i>Butl.</i>	130
3. <i>Epinephele pallescens</i> , <i>Butl.</i>	65
4. <i>Mycalesis Junonia</i> , <i>Butl.</i>	146
5. — <i>Felderi</i> , <i>Butl.</i>	144
6. — <i>auricruda</i> , <i>Butl.</i>	131
7. — <i>Semperi</i> , <i>Butl.</i>	137
7a. — — (anal valves of ♂, magnified).	
8. — <i>angulosa</i> , <i>Butl.</i>	130
9. — <i>mandosa</i> , <i>Butl.</i>	139
10. — <i>Madjicosa</i> , <i>Butl.</i>	137
11. — <i>æthiops</i> , <i>Butl.</i>	141
12. — <i>justinella</i> , <i>Butl.</i>	135
13. — <i>fraterna</i> , <i>Butl.</i>	145
14. — <i>malsarida</i> , <i>Butl.</i>	134

IV.

Fig.		Page
1.	<i>Argyrophorus Williamsianus</i> , <i>Butl.</i>	159
2.	<i>Oxeoschistus simplex</i> , <i>Butl.</i>	180
3.	<i>Lymanopoda ferruginosa</i> , <i>Butl.</i>	169
4.	<i>Lasiophila Hewitsonia</i> , <i>Butl.</i>	182
5.	<i>Lymanopoda venosa</i> , <i>Butl.</i>	171
6.	— <i>Acræida</i> , <i>Butl.</i>	171
7.	<i>Pedaliodes angularis</i> , <i>Butl.</i>	176
8.	— <i>granulata</i> , <i>Butl.</i>	173
9.	<i>Hypocysta antirius</i> , <i>Butl.</i>	168
10.	<i>Pronophila obscura</i> , <i>Butl.</i>	184

V.

1.	<i>Hipio Cramer</i> , <i>Butl.</i>	5
2.	<i>Anchiphlebia Taygetina</i> , <i>Butl.</i>	107
3.	— (neurulation of archæa, <i>Hübner</i>)	106
4.	<i>Cænonympha marginalis</i> , <i>Butl.</i>	42
5.	<i>Euptychia perfuscata</i> , <i>Butl.</i>	18
6.	<i>Lasiophila orbifera</i> , <i>Butl.</i>	182
7.	<i>Steroma superba</i> , <i>Butl.</i>	172
8.	<i>Erites argentina</i> , <i>Butl.</i>	188

ERRATA.

- Page 14. For "Typical Species *Eutychia Herse*," read "Typical Species *Eutychia Herse*."
- „ 16. *E. Clarissa* must give way to *Penelope*, Fabr., described in the 'Systema Entomologiæ.'
- „ 17. For (*Eutychia divergens*) "p. 464. n. 14," read "p. 464. n. 15."
- „ 33. Omit note after *E. Bysses*, var. It was inserted in the MS. before the type of *E. Necys* had been examined.
- „ 107. *A. Morna*, Fabr. = *A. Philoctetes*, Linn. ♀.
- „ 114. *L. arcuata*, Butl. = *L. Arete*, Cram. ♂, hitherto considered identical with *L. Europa*.
- „ 138. *M. Hesione*, Cramer, must change places with *M. Medus*, Fabr., described in the 'Syst. Ent.'
- „ 166. For *Geitoneura*? *Cordace* read *Heteronympha Cordace* (Hab. Melbourne).

CATALOGUE

OF

DIURNAL LEPIDOPTERA.

FAMILY SATYRIDÆ, *Westwood* (1851).

Genus 1. CÆEROIS, *Hübner* (1816).

Typical Species *Cærois* (*Arcesilaë*) *Chorinæus*.

1. *Cærois Chorinæus*.

Papilio Chorinæus, *Fabricius*, *Syst. Ent.* p. 484 (1775).

Arpidea Chorinæus, *Duncan*, *Foreign Butterf.* p. 180, pl. 23 (1837).

Cærois Chorinæus, *Westwood* in *Gen. Diurn. Lepid.* p. 367. n. 1 (1851).

Papilio Arcesilaus, *Cramer*, *Pap. Exot.* iv. p. 11, pl. 294. figs. A-D (1782); *Stoll*, (transformations) pl. vi. (1787-91).

Cærois Arcesilaë, *Hübner*, *Verz. bek. Schmett.* p. 56. sp. 536 (1816).

Surinam (*Fabricius*), Ega, Tunantino.

B.M.

Genus 2. MELANITIS*, *Fabricius* (1808).

Typical Species *Melanitis Leda*.

1. *Melanitis Leda*.

Papilio Leda, *Linnaeus*, *Syst. Nat.* ii. p. 773. n. 150 (1766).

Oreas (*marmorata*) *Leda*, *Hübner*, *Samml. exot. Schmett.* Band i. pl. 91. figs. 1-4 (1806-27).

* *Melanitis* of the *Eurytelidae* is the *Elymnias* of *Hübner*.

Melanitis Leda, *Fabricius, Syst. Gloss. (Illiger's Mag. vi. p. 282, 1808)*; *F. Moore, Cat. Lep. Mus. East-Ind. Comp. p. 222. n. 461 (1857)*.

Hipio Leda, *Hübner, Verz. bek. Schmett. p. 56. n. 538 (1816)*.

Satyrus Leda, *Godart, Enc. Méth. ix. p. 478. n. 4 (1819)*.

Cyllo Leda, *Westwood in Gen. Diurn. Lepid. p. 361. n. 1 (1851)*.
Vars. 1, 2, 3, 3b, 4, 5, Butler, Ann. & Mag. Nat. Hist. (Jan. 1867).

Java, Ceylon, Moulmein, Northern India, Celebes, Ceram. B.M.

MELANITIS TRISTIS.

Cyllo tristis, *Felder, Reise der Fregatte Novara, iii. p. 464. n. 785 (1867)*.

Northern India (Felder).

Dr. Felder remarks with regard to this insect:—"The form of the wings differs proportionably from all the examples of *C. Leda*, from many localities, now before us. The front wings are longer in the inner margin; the hind wings likewise are broader, but shorter within than in *C. Leda*."

Notwithstanding, I cannot think it is more than a variety; for I am fully convinced that this species is capable of almost any amount of variation in form as well as in colour.

MELANITIS ISMENE.

Papilio Ismene, *Cramer, Pap. Exot. i. pl. 26. figs. A, B (1779)*.

Papilio Mycena, *Cram. ibid. iv. pl. 291. fig. F (1782)*.

Papilio Arcensia, *Cram. ibid. pl. 292. fig. C (1782)*.

Melanitis Banksia, *Moore, Cat. Lep. Mus. East-Ind. Comp. p. 222. n. 462 (1857)*.

Cyllo Leda, vars. *2a, 4a, 5a, 7-10, 11-18, 19, 20, 21, 23, Butler, Ann. & Mag. Nat. Hist. (January 1867)*.

Moreton Bay, Ashanti, Sierra Leone, Congo, North India.

B.M.

MELANITIS TAITENSIS.

Cyllo Leda, var. *Taitensis*, *Felder, Verhandl. zool.-bot. Gesellsch. in Wien, xii. (1862)*; var. *22, Butl. (1867)*.

Papilio Leda, *Cramer, Pap. Exot. iv. pl. 292. fig. A (1782)*.

Tahiti (Felder), Oceania.

B.M.

MELANITIS DURYODANA.

Cyllo Duryodana, *Felder, Reise der Fregatte Novara, vol. iii. p. 464. n. 786 (1867)*.

Bengal, Cachar, Assam (Felder).

"Larger than *C. Banksia* (F.); all the wings longer, and more strongly angulated."

MELANITIS OBSOLETA.

Cylo obsoleta, *Felder, Reise der Fregatte Novara*, iii. p. 464.
n. 787 (1867).

Celebes?, var. Java (Felder). B.M.

I think this must be my variety 8 of *M. Leda*; if so, it is only a form of *Ismene*.

MELANITIS SOLANDRA.

Papilio Solandra, *Fabricius, Syst. Ent.* p. 500 (1775).

Cylo Solandra, *Boisduval, Voy. dans l'Océanie, Ent.* pt. 1.
p. 142 (1832-35).

Cylo Helena, *Westwood in Gen. Diurn. Lepid.* p. 361. n. 2
(1851); *Bull. vars.* 1 a, 2 b, 5 b, 19 a, 20 a (1867).

? *Cylo fulvescens*, *Felder, Reise der Fregatte Novara*, iii. p. 465.
n. 788 (1867).

Moreton Bay, Mauritius, Cape of Good Hope. B.M.

MELANITIS BANKSIA.

Papilio Banksia, *Fabricius, Syst. Ent.* p. 499 (1775).

Satyrus Banksia, *Godart, Enc. Méth.* ix. p. 477. n. 3 (1819).

Cylo Banksia, *Westwood in Gen. Diurn. Lepid.* p. 361. n. 3
(1851).

Var. *Cylo Boisduvalia*, *Felder, Diagn. Lepid.* p. 48. n. 96
(1859); *Bull. vars.* 10 a, 18 a, 24-27, 28 a (1867).

Cape of Good Hope, Congo, Moreton Bay, Australia. B.M.

MELANITIS PHEDIMA.

Papilio Phedima, *Cramer, Pap. Exot.* iv. pl. 292. fig. B (1782);
Bull. var. 28 (1867).

Australia. B.M.

I have found it utterly impossible to separate the above forms specifically from one another; slight differences in form and in the colouring of the upperside are the only guides by which to distinguish the different named varieties; but even these are not constant. The true *Leda*, however, appears to occur only in India and the Indian Islands; but the slight variety *Ismene* (including *Mycena* and *Arcensia*) has a much wider range, being found in India, Java, Australia, and Africa. The ? *Taitensis* of Felder links this form to *Solandra*; the latter and *Banksia** appear to exist only in Australia and Africa, whilst *Phedima* seems to be strictly confined to Australia.

2. Melanitis Vamana.

Melanitis Vamana, *Moore, Cat. Lep. Mus. East-Ind. Comp.*
p. 223. n. 463 (1857).

* The *Banksia* of Moore's Catalogue is the *Ismene* of Cramer.

Darjeeling, North India (Moore), East Indies. B.M.
 Very possibly only another form of *Leda*.

3. **Melanitis Ambasara.**

Melanitis Ambasara, Moore, *Cat. Lep. Mus. East-Ind. Comp.*
 p. 223. n. 464 (1857).

Java (Moore).

This species appears to have been in the collection of Dr. Horsfield. It is possibly a variety of *Ismene*.

4. **Melanitis Bela.**

Melanitis Bela, Moore, *Cat. Lep. Mus. East-Ind. Comp.* p. 223.
 n. 465 (1857).

Northern India, Assam, Darjeeling (Moore), Sylhet. B.M.

5. **Melanitis Suyudana.**

Melanitis Suyudana, Moore, *Cat. Lep. Mus. East-Ind. Comp.*
 p. 224. n. 466 (1857).

Java.

Horsf. Collection.

6. **Melanitis Varaha.**

Melanitis Varaha, Moore, *Cat. Lep. Mus. East-Ind. Comp.*
 p. 224. n. 467 (1857).

Canara (Moore).

7. **Melanitis Erichsonia.**

Cylo Erichsonia, Felder, *Lep. Nov. Ins. Philipp. in Lep. Diagn.*
 p. 43. n. 95 (*Wien. Monats.* v. 1861).

Mindoro (Colls. Semper and Felder).

Like the preceding, but with more distinctly angulated wings. I have seen a specimen of this species in Mr. Bates's collection; it is smaller than *Suyudana*, and has the wings formed as in *Solandra*, Fabr.

8. **Melanitis Gokala.**

Melanitis Gokala, Moore, *Cat. Lep. Mus. East-Ind. Comp.*
 p. 224. n. 468 (1857).

Canara (Moore).

9. **Melanitis Gerdrudtus.**

Papilio Gerdrudtus, Fabricius, *Ent. Syst.* iii. pt. 1. p. 72. n. 224
 (1793).

Cyllo? Gerdruttus, *Westwood in Gen. Diurn. Lepid.* p. 361.
n. 4 (1851).

Hab. — ?

Appears to be a variety of *Leda*; but in some respects it appears to approach more nearly to *Vamana*.

10. *Melanitis Aswa*.

Cyllo Aswa, *F. Moore, Proc. Zool. Soc. London*, p. 769 (1865).

North-eastern Bengal (Moore).

11. *Melanitis Gnophodes*, sp. nov. Pl. II. fig. 1.

♂ ♀. Alæ supra velut in *Gnophode* (*Parmeno*) coloratæ, subtus variantes, coloribus *Phedimæ* vel *Banksiæ*.

Exp. alar. unc. $3\frac{3}{8}$ — $3\frac{5}{8}$.

India.

B.M.

The male of this species resembles *Gnophodes Parmeno* on the upper-side; the female has a brighter orange band, and resembles the Natal form of the same insect. It may possibly be a link between the two genera, though a true *Melanitis*.

Subgenus 1. *HIPIO*, *Hübner* (1816).

Typical Species *Hipio Constantia*.

1. *Hipio Constantia*.

Papilio Constantia, *Cramer, Pap. Exot.* ii. pl. 133. figs. A, B (1779).

Hipio Constantia, *Hübner, Verz. bek. Schmett.* p. 56. n. 537 (1816).

Satyrus Constantia, *Godart, Enc. Méth.* ix. p. 477 (1819).

Cyllo Constantia, *Boisduval, Voy. de l'Astrolabe, Ent.* pt. 1. p. 141. n. 2 (1832).

Amboyna (Boisduval), Northern India (Westwood), Darjeeling (Moore).

Not in the Collection of the British Museum, as stated by Westwood, but in the Collection of F. Moore.

2. *Hipio Crameri*.

Cyllo Crameri, *Builer, Ent. Month. Mag.* iii. p. 77. n. 4 (1866).

Locality not known.

B.M.

Allied to *Constantia*, but differing in form, size, and the number and position of the ocelli.

3. *Hipio amabilis*.

Cyllo amabilis, *Boisduval, Voy. de l'Astrolabe, Entom.* pt. 1. p. 140, pl. 2. figs. 1, 2 (1832).

New Ireland (Boisduval).

Allied to the preceding species, but of a slightly different form, a white band instead of a yellow one in the front wings, a white subapical spot on the underside, instead of an unequally geminate ocellus; the ocelli of the hind wings blue, and five instead of four in number.

4. *Hipio Atrax*.

Cylo Atrax, *Felder, Lep. Nov. Ins. Philipp. in Lep. Diagn.*
p. 42. n. 94 (*Wien. Monats.* v. 1861).

Luzon (Coll. Semper), Burias, Lochan (Coll. Felder). B.M.

5. *Hipio ? velutina*.

Cylo velutina, *Felder, Reise der Fregatte Novara*, iii. p. 463.
n. 784 (1867).

Celebes (Felder).

According to Felder, this belongs to the *Constantia* group; it does not, however, seem to be banded above.

The following species seems almost distinct enough to form a new genus; it appears to come near the genus *Elymnias* (*Eurytelidæ*) in several respects. The concave form of the discocellulars of the front wings, the more wedge-shaped hind-wing cell, the violet colouring of the upperside, and the large silky dark patch in the front wings are sufficient characters to warrant its separation from *Melanitis*. I have, however, separated it as a subgenus under the name of *Cyllogenes*.

Subgenus 2. CYLLOGENES, *Butler* (1868).1. *Cyllogenes Suradeva*.

Melanitis Suradeva, *Moore, Cat. Lep. Mus. East-Ind. Comp.* i.
p. 225. n. 469 (1857).

Darjeeling (Moore), Northern India. B.M.

Genus 3. GNOPHODES; *Westwood* (1851).

Typical Species *Gnophodes Parmeno*.

1. *Gnophodes Parmeno*.

Gnophodes Parmeno, *Hewitson in Gen. Diurn. Lepid.* p. 363.
n. 1, pl. 61. fig. 2 (1851).

Gnophodes Parmeus, *Chenu, Enc. d'Hist. Nat. Pap.* fig. 294
(1851-57).

Sierra Leone, Ashanti. B.M.

Local form. *Gnophodes Parmeno*, *Trimen, Rhop. Afric. Austral.*
p. 190. n. 107 (1866).

Port Natal. B.M.

2. *Gnophodes Morpena*, Westwood in litt.

Alæ supra piceæ, fascia subapicali ferruginea: posticæ maris macula magna ovali subcostali alba velut in speciebus *Euploæ*.

Alæ subtus violaceo-fuscæ, ochreo variegatæ, punctis ocellaribus, anticis quinque, posticis septem, ad plicas positis, ochreis, extus fusco cinctis, iride brunnea indistincta; fascia tenui lunulata ferruginea discali; maculis permagnis costalibus irregularibus ochreis anticis quatuor, posticis tribus; cella posticarum area externa albido rorata, in medio a fascia abbreviata nigrescente secta.

Exp. alar. unc. 3.

Congo, Ashanti.

♂ ♀, B.M.

The specimens of this species differ a little in the colouring of the underside, the ochreous colour giving place to pale olive.

3. *Gnophodes Betsimena*.

Cyllo Betsimena, *Boisduval, Fauna Ent. de Madag.* p. 58. n. 1 (1833); also of Westwood (1851).

Tamatave (Boisduval), Ashanti.

♀, B.M.

Seems to be allied to *Parmeno*, but with a pale whitish-yellow band.

Genus 4. *AMPHIDECTA*, Butler (1867).

Typical Species *Amphidecta pignerator*.

1. *Amphidecta pignerator*.

Amphidecta pignerator, Butler, *Ann. & Mag. Nat. Hist.* xx. p. 405, pl. 9. fig. 11 (1867).

Ega.

Coll. Bates.

This curiously formed species is evidently closely allied to the species of the genus *Taygetis*.

Genus 5. *TAYGETIS*, Hübner (1816).

Typical Species *Taygetis Virgilia*.

1. *Taygetis Mermeria*.

♀. *Papilio Mermeria*, Cramer, *Pap. Exot.* i. pl. 96. fig. B (1799); Stoll, *Suppl.* pl. 7. figs. 1, 1 a (transformations) (1787-91).

? *Satyrus Mermeria*, Godart, *Enc. Méth.* ix. p. 485. n. 23 (1819).

Taygetis Mermeria, Hübner, *Verz. bek. Schmett.* p. 55. n. 531 (1816).

Brazil, St. Paulo.

♀, B.M.

♂. Local form. *Pap. Mermeria*, Cramer, *Pap. Exot.* iii. pl. 289. figs. E, F (1782).

♀. *Taygetis Mermeria*, Hübner, *Samml. exot. Schmett.* ii. pl. 84. figs. 1, 2 (1806).

Rio Janeiro.

♂ ♀, B.M.

Var. Alæ subtus violaceo-fuscescentes.

Bolivia.

♂, B.M.

Local form. Faunus tenebrosus, *Blanchard in Voy. d'Orbigny*, p. 222, pl. 32. figs. 1, 2 (1847).

Ega.

♂, B.M.

Very like the preceding form, but with smaller ocellate spots on the underside of the front wings.

Local form. Alæ supra ochraceæ violaceo tinctæ, subtus multo pallidiores; ocellis fasciisque olivaceo-fuscis; fascia media posticarum purpurascens.

Nicaragua.

♀, B.M.

2. *Taygetis excavata*, sp. nov. Pl. I. fig. 1.

♂. Alæ supra olivaceo-fuscae, ad margines paulum obscuriores: anticæ apice acute producto: posticæ margine anali externo valde excavato; fere ut in *T. Celia*, Cramer.

Alæ subtus rufo-ferruginæ, velut in *Mermeria fasciatæ* et ocellatæ, fascia media posticarum virescente; area basali anticarum fuscescente, angulo anali albicante.

Exp. alar. unc. $3\frac{1}{2}$.

Honduras.

♂, B.M.

This may be only an extreme form of *Mermeria*. It most nearly approaches the variety figured by Blanchard under the name of *tenebrosus*, but it differs in the deep sinuations in the anal margin of the hind wings and in its much more gorgeous colouring below.

3. *Taygetis armillata*, sp. nov. Pl. I. fig. 3.

♂. Alæ supra olivaceo-fuscae obscuræ, margine externo obscuriore, ciliis pallidis: posticæ margine anali excavato.

Alæ subtus pallidiores: anticæ area apicali fuscescente, nitente; area interna olivacea; costa trifasciolata fasciis pallidis ochraceis, apud marginem externum ocellis quinque indistinctis regularibus roseo roratis; margine externo atomis albidis angulariter submarginato; posticæ cinereo-nigræ nitentes, roseo roratæ; fasciola subbasali ochreo-fusca convexa; fascia discali tenui, medio angulata, nigra, atomis albidis extus marginata, ocellis sex contiguis regularibus roseis nigro roratis, ochreo pupillatis, anali geminato; margine externo albido rorato et submarginato; apice aliquando roseo.

Exp. alar. unc. $3\frac{5}{8}$.

Bogota, var. Bolivia.

B.M.

Intermediate between *Mermeria* and *Chelys*.

4. *Taygetis Celia*.

Papilio Celia, *Cramer, Pap. Exot.* iii. pl. 242. fig. C (1782).

Taygetis Celia, *Hübner, Verz. bek. Schmett.* p. 56. n. 533 (1816).

Surinam (Cramer), Brazil.

B.M.

There is one very faded specimen of this species in the Collection.

5. *Taygetis Chelys*.

Papilio Chelys, *Fabricius, Ent. Syst.* iii. pt. 1. p. 80. n. 249 (1793).

Satyrus Chelys, *Godart, Enc. Méth.* ix. p. 484. n. 21 (1819).

Taygetis Chelys, *Westwood in Gen. Diurn. Lepid.* p. 356. n. 3 (1851).

Brazil.

♂, B.M.

Var. Alæ posticæ subtus fascia obscuriore castaneo-fusca subcostali.

Brazil.

♂ ♀, B.M.

6. *Taygetis Ypthima*.

Taygetis Ypthima, *Hübner, Samml. exot. Schmett.* ii. pl. 83. figs. 3, 4 (1806).

Brazil.

This species seems to be very closely allied to *Chelys*. There are no specimens of it in the Collection.

7. *Taygetis Larua*.

Taygetis Larua, *Felder, Reise der Fregatte Novara*, iii. p. 466. n. 790 (1867).

New Granada, Bogota (Felder).

This seems very close to *T. Ypthima*.

8. *Taygetis nympha*, sp. nov. Pl. I. fig. 7.

Alæ forma fere *Chelydis*; supra olivaceo-fuscæ, maris obscuriores linea submarginali nigrescente subintegra; marginibus roseo tinctis.

Alæ subtus pallidiores, lineis tribus, ocellisque fere velut in *Virgilia*, sed multo magis irregularibus; linea submarginali inter venas bidentata: *alæ* maris velut in *Virgilia* roseo suffusæ, ocellis brunneis: *alæ* feminae pallidæ area discali pallidiore; linea media intus fuscescente, extus albicante.

Exp. alar. ♂ unc. 3, ♀ unc. 3½.

Honduras.

♂ ♀, B.M.

Intermediate between *Chelys* and *Virgilia*.

9. *Taygetis Virgilia*.

♀. *Papilio Virgilia*, *Cramer, Pap. Exot.* i. pl. 96. fig. C (1779).

Taygetis Virgilia, *Hübner, Verz. bek. Schmett.* p. 55. n. 529 (1816).

♂. *Var. Papilio Rebecca*, *Fabricius, Ent. Syst.* iii. pt. 1. p. 75. n. 236 (1793).

Satyrus Rebecca, *Godart, Enc. Méth.* ix. p. 484. n. 22 (1819).
Taygetis Rebecca, *Westwood in Gen. Diurn. Lepid.* p. 357. n. 4 (1851).

Para, Brazil, Pernambuco, Venezuela.

B.M.

The female of what appears to be the typical *Rebecca* has the hind wings on the upperside distinctly margined with ferruginous; in the true *Virgilia* this colouring almost entirely disappears, and the colours of the underside become brighter.

10. *Taygetis nymphosa*, sp. nov. Pl. I. fig. 6.

♂. Alæ supra olivaceo-fuscae, forma fere *Virgiliae*, posticæ autem margine anali magis sinuato.

Alæ subtus pallidiores, lineis fere velut in *Virgilia*, discali autem ad marginem externum magis approximante, et linea submarginali, inter venas distincte bidentatæ; area tota apicali multo pallidiore, linea discali extus albido marginata; ocellis parvis, secundo, quinto et sexto posticarum nigris; aliter velut in *Virgilia*.

Exp. alar. unc. $2\frac{1}{8}$.

Bolivia.

B.M.

Closely allied to *nympha* and *Virgilia*, but quite distinct.

11. *Taygetis Chrysogone*.

Taygetis Chrysogone, *Hewitson in Gen. Diurn. Lepid.* p. 357. n. 11, pl. 60. fig. 4 (1851).

Venezuela, Bogota.

B.M.

12. *Taygetis erubescens*, sp. nov. Pl. I. fig. 2.

♂. Alæ supra velut in *Virgilia*; subtus roseo suffusæ et olivaceo hic illic roratæ; area interna anticarum olivaceo suffusa; area basali posticarum ferruginosa; fascia lata communi irregulari media pruno-ferruginea, nigro utrinque marginata, fusco varia, maculam magnam ferrugineam ovalem ad cellæ anticarum finem, alteramque in cella posticarum, includente; ocellis velut in *Virgilia*, magis autem ad fasciam mediam approximantibus; margine externo velut in *Virgilia*.

Exp. alar. unc. $3\frac{1}{8}$.

Bogota.

B.M.

Probably a local form of *Virgilia*, but in the form of its wings more like *Chrysogone*. The central band of the underside differs from that of *Virgilia*, the subbasal line being much more convex and the discal line irregularly undulated and less distinctly margined with pinky white.

13. *Taygetis Echo*.

Papilio Echo, *Cramer, Pap. Ecot.* i. pl. 57. figs. C, D (1779);
Fabricius, Spec. Ins. t. ii. p. 91. n. 399 (1781).

Taygetis Echo, *Westwood in Gen. Diurn. Lepid.* p. 357. n. 8 (1851).

Satyrus Betro, *Godart, Enc. Méth.* ix. p. 484. n. 20 (1819).

Hætera (Antirrhaea) (sic) Betro, *Westwood in Gen. Diurn. Lepid.* p. 366. n. 13 (1851).

Surinam (Cramer), Brazil, Ega.

B.M.

14. *Taygetis Cleopatra*.

Taygetis Cleopatra, *Felder, Wien. ent. Monatsch.* (1862).

Rio Negro and near Bahia (Felder), Brazil?

B.M.

Possibly this may prove to be nothing more than an extreme form of *Andromeda*, with which it is partially connected by the *Laches* of Fabricius.

T. CLEOPATRA, var. *Alæ subtus area basali lilacino suffusa.*

Brazil?

B.M.

15. *Taygetis Sylvia*.

Taygetis Sylvia, *H. W. Bates, Ent. Month. Mag.* iii. p. 153. n. 110 (1866).

Panama and Upper Amazons.

This seems, from the description, to be intermediate between *Cleopatra* and *Andromeda*. The ocelli of the hind wings seem to be as in *Para* specimens of the latter species. Mr. Bates says that it "resembles the smaller and more clearly marked individuals of *T. Andromeda*; but the fore-wing outer border of *T. Andromeda* is always angulated near the apex." It appears to differ also in the uniform brown colour of the base of the wings on the underside.

16. *Taygetis Andromeda*.

Papilio Andromeda, *Cramer, Pap. Exot.* i. pl. 96. fig. A (1779).

Satyrus Andromeda, *E. Doubleday, List Lep. Brit. Mus.* p. 119.

Fam. ix. (1844).

Taygetis Andromeda, *Westwood in Gen. Diurn. Lepid.* p. 357. n. 5 (1851).

Taygetis fatua, *Hubner, Verz. bek. Schmett.* p. 56. n. 532 (1816).

Santa Lucia, Bogota, Venezuela, Para.

B.M.

Var. *a.* *Alæ subtus area apicali ochreo variegata; ocellis apicalibus omnibus ochraceis.*

Brazil, Nicaragua.

B.M.

Var. *b.* *Alæ anticæ margine externo magis undato, apice distincte angulato; ocellis subtus obscurioribus fuscis; fasciis latioribus, externa posticarum medio magis angulata: alæ omnes lilacinæ.*

Venezuela.

B.M.

TAYGETIS LACHES.

Papilio Laches, *Fabricius, Ent. Syst.* iii. pt. 1. p. 229. n. 719 (1793).

Satyrus Laches, *Godart, Enc. Méth.* ix. p. 485. n. 24 (1819).

Oreas marmorea *Thamyra* (part.), *Hübner, Samml. exot. Schmett.* i. pl. 88. figs. 3, 4 (1806).

Para.

♀, B.M.

I am convinced, from a careful comparison of our specimens with Hübner's figures, that he has figured two distinct insects as sexes of *Thamyra*. The female figure appears to have been in a measure assimilated to that of the male, yet, notwithstanding, the difference in the ground-colour of the upperside and in the form of the central bands of the underside is at once apparent. The sexes of *Andromeda* do not thus differ, and I therefore conclude that the insects figured are distinct.

17. Taygetis Thamyra.

Papilio Thamyra, *Cramer, Pap. Exot.* iii. pl. 242. fig. B (1782).

Oreas marmorea *Thamyra* (part.), *Hübner, Samml. exot. Schmett.* i. pl. 88. figs. 1, 2 (not good) (1806).

Taygetis Thamyra, *Hübner, Verz. bek. Schmett.* p. 55. n. 530 (1816).

Brazil, Ega.

♂, B.M.

This insect, though it comes near *Laches*, must, I think, be considered a distinct species; the deep pitchy colouring of the upperside, the shorter and more rounded form of the hind wings, the deeper colouring of the underside, the outer band of the hind wings angulated and placed much nearer to the middle of the wing, and the more regular ocelli with ochre-tinted pupils readily separate it from all the varieties of *Andromeda*.

18. Taygetis calliomma.

Taygetis calliomma, *Felder, Wien. entom. Monatschr.* (1862); *Diagn. Lepid.* p. 48. n. 140; *Nov. Voy.* iii. tab. 66. fig. 7 (1867).

Bogota.

19. Taygetis albinotata.

Taygetis albinotata, *Butler, Ann. & Mag. Nat. Hist.* xx. 3rd ser. p. 216, pl. 4. figs. 6, 7 (1867).

Bolivia.

B.M.

This species is one of the most distinct and beautiful forms in the genus; we only have one specimen, in bad condition.

20. *Taygetis satyrina*.

Taygetis satyrina, *H. W. Bates, Ent. Month. Mag.* i. p. 179. n. 72 (1865).

Guatemala (central valleys).

"Not closely allied to any other known species of the genus" (Bates in *Ent. Mag.*).

21. *Taygetis inornata*.

Taygetis inornata, *Felder, Reise der Fregatte Novara*, iii. p. 466. n. 792 (1867).

New Granada, Bogota (Felder).

"Belongs to the *Valentina* group; is the size of *Marpessa*."

22. *Taygetis Valentina*.

Papilio Valentina, *Cramer, Pap. Exot.* iii. pl. 242. fig. A (1782).

Satyrus Valentina, *Godart, Enc. Méth.* ix. p. 486. n. 26 (1851).

Surinam (Cramer), Cayenne.

B.M.

Appears to be intermediate between *Andromeda* and *Penelea*.

23. *Taygetis Penelea*.

♀. *Papilio Penelea*, *Cramer, Pap. Exot.* ii. pl. 101. fig. G (1779); also of *Fabricius* (1788).

Satyrus Penelea, *Godart, Enc. Méth.* ix. p. 486. n. 28 (1819).

Bogota, Santa Lucia, Venezuela.

B.M.

24. *Taygetis Marpessa*.

Debis *Marpessa*, *Hewitson, Exot. Butterf.* iii. p. 73, pl. 37. fig. 2 (1862); *Taygetis Marpessa* (corrections).

Amazon (Hewitson).

25. *Taygetis neonympha*.

Taygetis neonympha, *Felder, Reise der Fregatte Novara*, iii. p. 467. n. 793 (1867).

New Granada, Bogota (Felder).

"Somewhat resembles *Euptychia* (*neonympha*) *Antonina*," *Felder*.

26. *Taygetis Euptychidia*, sp. nov. Pl. I. fig. 5.

♂. Alæ forma *Euptychia* (*Libye*); supra olivaceæ marginibus paulum roseo tinctæ, linea submarginali valde indistincta undata, altera marginali et margine ipso fuscis.

Alæ subtus olivaceæ multo pallidiores, violaceo tinctæ, lineis tribus communibus nigris valde irregularibus sinuatis et dentatis, media subobliqua; linea marginali et margine ipso fuscis; puncto ad cellæ anticarum finem, altero lineam subbasalem extus attingente, ochreis;

linea discali posticarum extus griseo nebuloſa: anticæ ocellis quinque, ſecundo et quinto nigris, tertio et quarto ochraceis ochreo pupillatis, aliis albo pupillatis: poſticæ ocellis quinque ſimilibus, ſed quinto maximo.

Exp. alar. unc $2\frac{5}{16}$.

Brazil, Para.

B.M.

A local form of *T. Valentina*.

27. *Taygetis anophthalma*.

Taygetis anophthalma, *Felder, Reise der Fregatte Novara*, iii. p. 467. n. 794 (1867).

New Granada, Bogota (Felder).

"The ſmalleſt known ſpecies of *Taygetis*, in the form of its wings very like *T. neonympha*, but differing in the convex outer margin of the front wing and the ſhorter hind wing, in appearance reſembling the *Erebia*. The palpi have longer cirri, and the wings are more deeply fringed."

Genus 6. EUPTYCHIA, *Hübner* (1816).

Typical ſpecies *Euptychia Herse*.

Section *Megisto*, *Hübner* (1816).

Typical ſpecies *Megisto (Cymela) Eurytus*.

1. *Euptychia Ocirrhoë*.

Papilio Ocirrhoë, *Fabricius, Gen. Ins.* p. 260 (1776); *Ent. Syst.* iii. pt. 1. p. 96. n. 297 (1793).

Euptychia Ocirrhoë, *Westwood in Gen. Diurn. Lepid.* p. 373. n. 1 (1851).

Oreas (strigata) Ocyrrhoë, *Hübner, Samml. exot. Schmett.* i. pl. 85. figs. 1-4 (1806).

Euptychia Ocyrrhoë, *Hübner, Verz. bek. Schmett.* p. 54. n. 505 (1816).

Satyrus Ocyrrhoë, *Godart, Enc. Méth.* ix. pp. 464 & 489. n. 41 (1819).

Papilio Hesione, *Sulzer, Gesch. der Insekt.* p. 144, t. 17. figs. 3, 4 (1776).

Papilio Cissia, *Cramer, Pap. Exot.* iii. p. 3, pl. 194. figs. D, E (1782).

Euptychia Cissia, *E. Doubleday, List Lep. Brit. Mus.* i. p. 122 (1844).

Panama, Brazil, Honduras, Caraccas.

B.M.

Var. 1 a. *Butler, Monog. Proc. Zool. Soc.* p. 459 (1866).

Polochie Valley, Guatemala.

Coll. Salvin.

Var. 2 b. *Butl. ibid.* *Calpurnia, Feld.* ?

Obydos (Coll. Bates), Panama.

B.M.

2. Euptychia Lydia.

Papilio Lydia, *Cramer, Pap. Exot.* ii. pl. 148. figs. C, D (1779), but not of *Fabricius, Ent. Syst.*

Euptychia Lydia, *Hübner, Verz. bek. Schmett.* p. 54. n. 510 (1816); also of *Westwood* (*Ocirrhoë*, ♀ ?), *Butler* (1866).

Satyrus Lydius, *Godart, Enc. Méth.* ix. pp. 464 & 489. n. 40 (1819).

Surinam (*Cramer*).

3. Euptychia Calpurnia.

Neonympha Calpurnia, *Felder, Reise der Fregatte Novara*, iii. p. 484. n. 836 (1867).

Cayenne (*Felder*).

I think this must be identical with my *Ocirrhoë*, var. 1 b, received lately from Peru, but without the subanal spot on the upperside of the hind wings. It may be a distinct species.

4. Euptychia Ocypete.

Papilio Ocypete, *Fabricius, Gen. Ins. Mant.* p. 260 (1776); *Ent. Syst.* iii. pt. 1. p. 96. n. 296 (1793).

Euptychia Ocypete, *Hübner, Verz. bek. Schmett.* p. 54, n. 506 (1816).

Satyrus Ocypete, *Godart, Enc. Méth.* ix. pp. 463 & 488. n. 34 (1819).

Surinam (*Fabricius*). Var. Para.

B.M.

EUPTYCHIA TERENTIA.

Neonympha Terentia, *Felder, Reise der Novara*, iii. p. 483. n. 833 (1867).

E. Ocypete, var., *Butler, Monog. Proc. Zool. Soc.* p. 460 (1866).

Cayenne (*Felder*), Venezuela, Para, Tapajos (*Coll. Bates*). B.M.

5. Euptychia Helle.

Papilio Helle, *Cramer, Pap. Exot.* iii. pl. 194. figs. F, G (1782).

Papilio Ocypete, (*synonym*) *Fabricius*; *Eupt. Ocyp.* (*syn.*) *Westwood*.

Surinam (*Cramer*).

Var. *Butl. Monog. Proc. Zool. Soc.* p. 460 (1866).

Brazil, Para.

B.M.

6. Euptychia Myncea.

Papilio Myncea, *Cramer, Pap. Exot.* iv. pl. 193. fig. C (1782).

Euptychia Myncea, *Hübner, Verz. bek. Schmett.* p. 54. n. 516 (1816).

Satyrus Myncea, *Godart, Enc. Méth.* ix. pp. 463 & 488. n. 36 (1819).

- Neonympha clerica*, *Herrich-Schäffer, Lep. Ind. Syst.* (1864).
 Brazil. B.M.
- Var. *a.* *Butler, Monog. Proc. Zool. Soc.* p. 461 (1866).
 Para. B.M.
- Var. *b.* *Butl. ibid.*
 Para. Coll. Bates.
- Var. *c.* *Butl. ibid.*
 Honduras. B.M.
7. ***Euptychia Palladia.***
Euptychia Palladia, *Butler, Proc. Zool. Soc.* p. 461, pl. 39. fig. 21 (1866).
 Tapajos. Coll. Bates.
8. ***Euptychia terrestris.***
Euptychia terrestris, *Butler, Proc. Zool. Soc.* p. 462, pl. 39. fig. 1 (1866).
 Brazil, Para. B.M.
- Var. *Butl. ibid.*
 Lower Amazons. Coll. Bates.
9. ***Euptychia Clarissa.***
Papilio Clarissa, *Cramer, Pap. Exot.* iv. pl. 293. figs. D, E (1782).
Euptychia Clarissa, *Hübner, Verz. bek. Schmett.* p. 54. n. 503 (1816).
Papilio Penelope, *Fabricius, Ent. Syst.* iii. pt. 1. p. 96. n. 298 (1793).
Satyrus Penelope, *Godart, Enc. Méth.* ix. pp. 463 & 496. n. 39 (1819).
 Brazil, Para. B.M.
10. ***Euptychia Camerta.***
Papilio Camerta, *Cramer, Pap. Exot.* iv. pl. 293. fig. F (1782).
Neonympha Camerta, *Westwood in Gen. Diurn. Lepid.* p. 375. n. 3 (1851).
 Surinam (Cramer).
11. ***Euptychia Themis.***
Euptychia Themis, *Butler, Suppl. Monog. Proc. Zool. Soc.* p. 104. n. 1, pl. 12. fig. 13 (1867).
 Hab. — ? Coll. Hewitson.

12. *Euptychia usitata*.

Euptychia usitata, *Butler, Monog. Proc. Zool. Soc.* p. 463. n. 11,
pl. 39. fig. 2 (1866).

Venezuela. B.M.

Var. *a. Butl. ibid.*

Venezuela. B.M.

Var. *b. Butl. ibid.*

Polochie Valley, Guatemala. Coll. Salvin.

13. *Euptychia similis*.

Euptychia similis, *Butler, Monog. Proc. Zool. Soc.* p. 463. n. 12
(1866).

Guatemala. Coll. Salvin.

Var. *Butl. ibid.*, and pl. 12. fig. 10 (1867).

Nicaragua. B.M.

14. *Euptychia Pieria*.

Euptychia Pieria, *Butler, Monog. Proc. Zool. Soc.* p. 463. n. 13,
pl. 39. fig. 3 (1866).

Honduras. B.M.

15. *Euptychia austera*.

Euptychia austera, *Butler, Monog. Proc. Zool. Soc.* p. 464. n. 14,
pl. 39. fig. 4 (1866).

Bogota. B.M.

16. *Euptychia divergens*.

Euptychia divergens, *Butler, Monog. Proc. Zool. Soc.* p. 464.
n. 14, pl. 40. fig. 3 (1866).

Rio Negro. Colls. Bates & Salvin.

17. *Euptychia Eurytus*.

Papilio Eurytus, *Fabricius, Syst. Ent.* p. 487. n. 194 (1775-
78).

Euptychia Eurytus, *Butler, Proc. Zool. Soc.* p. 465. n. 16 (1866).

Papilio Eurytris, *Fabricius, Ent. Syst.* iii. pt. 1. p. 157. n. 485
(1793).

Neonympha Eurytris, *Westwood in Gen. Diurn. Lepid.* p. 357.
n. 1 (1851).

Satyrus Eurythris, *Godart, Enc. Méth.* ix. pp. 465 & 495. n. 57
(1819); also of *Boisduval & Leconte*.

Papilio Cymela, *Cramer, Pap. Exot.* ii. pl. 132. figs. C, D (1779).

Megisto Cymelia, *Hübner, Verz. bek. Schmett.* p. 54. n. 517
(1816).

United States. B.M.

18. **Euptychia Periphas.**

Satyrus Periphas, *Godart, Enc. Méth.* ix. pp. 465 & 495. n. 62 (1819).

Neonympha Periphas, *Westwood in Gen. Diurn. Lepid.* p. 375. n. 5 (1851).

Euptychia Periphas, *Butler, Proc. Zool. Soc.* p. 465. n. 17 (1866), pl. 12. fig. 16 (1867).

Brazil.

Coll. Hewitson.

19. **Euptychia Lethe.**

Euptychia Lethe, *Butler, Monog. Proc. Zool. Soc.* p. 465. n. 18 (1866), pl. 12. fig. 1 (1867).

Venezuela.

B.M.

20. **Euptychia perfuscata**, sp. nov.

♂. Alæ supra olivaceo-fuscae, stria marginali subintegra, altera angulis alternis undata submarginali fuscis; margine ipso nigro.

Alæ subtus pallidiores violaceo leviter tinctæ atomis ochreis omnino roratae; striis duabus mediis subintegris obliquis leviter undulatis fuscis; interna anticarum introrsum directa, lineis marginalibus velut supra nigris; fascia nebulosa discali ocellos includente abbreviata: alæ anticae ocellis quatuor regularibus, tertio paulum majore, primo et quarto nigris minoribus argenteo unipupillatis, aliis bipupillatis, omnibus fulvo cinctis: alæ posticae ocellis quinque, primo minimo, secundo unipupillato; tertio et quarto fuscis, aliis nigris argenteo bipupillatis, omnibus distincte fulvo cinctis; ocello minimo interno-anali, aliquando geminato, nigro.

Exp. alar. unc. $1\frac{3}{4}$.

Brazil, Para.

B.M.

21. **Euptychia Erigone.**

Euptychia Erigone (E. Mæpius, var. ?), *Butler, Proc. Zool. Soc.* p. 466. n. 20 (1866).

St. Paulo.

B.M.

Var. *Butl. ibid.*

Brazil, Ega.

Coll. Bates.

22. **Euptychia argyrospila.**

Euptychia argyrospila, *Butler, Proc. Zool. Soc.* p. 467. n. 21 (1866), pl. 11. fig. 12 (1867).

Brazil, Ega.

Coll. Bates.

23. **Euptychia Crantor.**

Papilio Crantor, *Fabricius, Ent. Syst.* iii. pt. 1. p. 158. n. 489 (1793); *Donovan, Ins. India*, pl. 37. fig. 4 (1800).

Satyrus Crantor, *Godart, Enc. Méth.* ix. p. 488. n. 37 (1819).

Euptychia Crantor, *Westwood in Gen. Diurn. Lepid.* p. 373. n. 7 (1851).

Hab. — ?

24. ***Euptychia* *Ocnus*.**

Euptychia *Ocnus*, *Butler, Proc. Zool. Soc.* p. 467. n. 23 (1866), pl. 11. fig. 8 (1867).

Tapajos (Coll. Bates), Villa Nova.

B.M.

25. ***Euptychia* *Eriphule*.**

Euptychia *Eriphule*, *Butler, Proc. Zool. Soc.* p. 468. n. 24. pl. 39. fig. 6 (1866).

Pernambuco.

B.M.

26. ***Euptychia* *Celmis*.**

Satyrus *Celmis*, *Godart, Ent. Méth.* ix. pp. 463 & 489. n. 38 (1819).

Euptychia *Celmis*, *Westwood in Gen. Diurn. Lepid.* p. 374. n. 26 (1851); *Butler, descr.* (1866).

Brazil (Godart).

27. ***Euptychia* *Electra*.**

Euptychia *Electra* (*E. Celmis*, var. ??), *Butler, Proc. Zool. Soc.* p. 468. n. 26, pl. 39. fig. 7 (1866).

Bahia.

B.M.

28. ***Euptychia* *variabilis*.**

Euptychia *variabilis*, *Butler, Proc. Zool. Soc.* p. 469. n. 27, pl. 39. fig. 8 (1866).

Pernambuco, Rio Janeiro.

B.M.

Var. *Butler, ibid.*

Rio Janeiro.

B.M.

29. ***Euptychia* *angularis*.**

Euptychia *angularis*, *Butler, Proc. Zool. Soc.* p. 106. n. 4, pl. 12. fig. 8 (1867).

Minas Geraes, Brazil.

Coll. Hewitson.

30. ***Euptychia* *straminea*.**

Euptychia *straminea*, *Butler, Proc. Zool. Soc.* p. 106. n. 3, pl. 12. fig. 9 (1867).

Minas Geraes, Brazil.

Coll. Hewitson.

31. *Euptychia affinis*.

Euptychia affinis, *Butler, Proc. Zool. Soc.* p. 469. n. 30, pl. 39.
fig. 9 (1866).

Pernambuco, Rio Janeiro.

B.M.

32. *Euptychia vestigiata*.

Euptychia vestigiata, *Butler, Proc. Zool. Soc.* p. 105. n. 2,
pl. 12. fig. 17 (1867).

Minas Geraes, Brazil.

Coll. Hewitson.

33. *Euptychia Pimpla*.

Neonympha Pimpla, *Felder, Wien. ent. Monatschr.* vi. p. 177.
n. 156 (1862).

Euptychia Pimpla, *Butler, Monog. Proc. Zool. Soc.* p. 470. n. 32
(1866).

Rio Negro, New Granada (Felder).

34. *Euptychia renata*.

Papilio renata, *Cramer, Pap. Exot.* iv. pl. 326. fig. A (1782).

Mycalesis ? renata, *Westwood in Gen. Diurn. Lepid.* p. 394.
n. 13 (1851).

Neonympha renata, *Herrich-Schäffer, Lep. Ind. Syst.* (1864).

Euptychia renata, *Butler, Proc. Zool. Soc.* p. 470. n. 33 (1866).

Surinam (Cramer), Rio Janeiro.

B.M.

Var. *a.* *E. Pimpla*, var., *Felder, Wien. ent. Monatschr.* vi. p. 177.
n. 156 (1862). *E. renata*, var., *Butl.* (1866).

Var. *b.* *Butler, Proc. Zool. Soc.* p. 470 (1866).

Tocantins River, Brazil.

Coll. Salvin.

35. *Euptychia Phronius*.

Satyrus Phronius, *Godart, Enc. Méth.* ix. pp. 466 & 496. n. 65
(1819).

Neonympha Phronius, *Westwood in Gen. Diurn. Lepid.* p. 376.
n. 21 (1851).

Euptychia Phronius, *Butler, Monog. Proc. Zool. Soc.* p. 470.
n. 34 (1866).

Brazil (Godart).

36. *Euptychia Pæon*.

Satyrus Pæon, *Godart, Enc. Méth.* ix. pp. 464 & 498. n. 43
(1819).

Euptychia Pæon, *Westwood in Gen. Diurn. Lepid.* p. 374. n. 23
(1851).

Brazil (Godart).

37. **Euptychia ochracea.**

Euptychia ochracea, *Butler, Proc. Zool. Soc.* p. 107. n. 5,
pl. 11. fig. 5 (1867).

Brazil.

Coll. Hewitson.

38. **Euptychia marmorata.**

Euptychia marmorata, *Butler, Proc. Zool. Soc.* p. 471. n. 37
pl. 40. fig. 1 (1866).

Rio Janeiro.

B.M.

39. **Euptychia Grimon.**

Satyrus Grimon, *Godart, Enc. Méth.* ix. pp. 464 & 490. n. 44
(1819).

Neonympha Grimon, *E. Doubleday, List Lep. Brit. Mus.* App.
p. 33 (1847); also of *Westwood* (1851).

Euptychia Grimon, *Butler, Proc. Zool. Soc.* p. 472. n. 39
(1866).

Brazil.

B.M.

40. **Euptychia Argante.**

Papilio Argante, *Cramer, Pap. Exot.* iii. pl. 204. figs. C, D
(1782).

Megisto Argante, *Hübner, Verz. bek. Schmett.* p. 54. n. 518
(1816).

Euptychia Argante, *Westwood in Gen. Diurn. Lepid.* p. 374.
n. 27 (1851).

Satyrus Argulus, *Godart, Enc. Méth.* ix., pp. 463 & 488. n. 35
(1819).

Surinam (*Cramer*).

41. **Euptychia ambigua.**

Euptychia ambigua, *Butler, Proc. Zool. Soc.* p. 472. n. 41,
pl. 39. fig. 10 (1866).

Rio Janeiro.

B.M.

42. **Euptychia modesta.**

Euptychia modesta, *Butler, Proc. Zool. Soc.* p. 473. n. 42
(1866), pl. 12. fig. 12 (1867).

Para.

Coll. Bates.

Var. *Butl. ibid.*

Cameta.

Coll. Bates.

43. *Euptychia Huebneri*.

Euptychia Huebneri, *Butler, Proc. Zool. Soc.* p. 473. n. 43,
pl. 39. fig. 11 (1866).

Para.

B.M.

44. *Euptychia Galesus*.

Satyrus Galesus, *Godart, Enc. Méth.* ix. pp. 465 & 496. n. 64
(1819).

Neonympha Galesus, *Westwood in Gen. Diurn. Lepid.* p. 376.
n. 20 (1819).

Euptychia Galesus, *Butler, descr. Proc. Zool. Soc.* p. 473. n. 44
(1866).

Euptychia Canthe, var., *Herrich-Schäffer, Ind. Syst.* (1864).

Brazil (Godart).

45. *Euptychia Vesper*.

Euptychia Vesper, *Butler, Proc. Zool. Soc.* p. 108. n. 8, pl. 12.
fig. 19 (1867).

Hab. — ?

Coll. Hewitson.

46. *Euptychia armilla*.

Euptychia armilla, *Butler, Proc. Zool. Soc.* p. 108. n. 9, pl. 12.
fig. 21 (1867).

Minas Geraes, Brazil.

Coll. Hewitson.

47. *Euptychia liturata*.

Euptychia liturata, *Butler, Proc. Zool. Soc.* p. 107. n. 7, pl. 12.
fig. 18 (1867).

Hab. — ?

Coll. Hewitson.

48. *Euptychia Sosybius*.

Euptychia Sosybius, *Fabricius, Ent. Syst.* iii. pt. 1. p. 219.
n. 684 (1793).

Satyrus Sosybius, *Godart, Enc. Méth.* ix. pp. 465 & 495. n. 63
(1819); *Boisduval and Leconte, Icon. Lep. Am. Sept.* 1.
pl. 63. figs. 1-4 (metamorph.) (1829).

Neonympha Sosybius, *Westwood in Gen. Diurn. Lepid.* p. 375.
n. 2 (1851).

Euptychia Sosybius, *Butler, Proc. Zool. Soc.* p. 474. n. 48
(1866).

East Florida.

♂ ♀, B.M.

49. Euptychia fallax.

Neonympha fallax, *Felder, Wien. ent. Monatschr.* vi. p. 177. n. 157 (1862).

Euptychia fallax, *Butler, Proc. Zool. Soc.* p. 474. n. 49, descr. (1866).

Rio Negro (Felder), Venezuela. B.M.

Var. *a.* *Butl. ibid.* (1866).

Venezuela. B.M.

Var. *b.* *Butl. ibid.* (1866).

Ega. Coll. Bates.

50. Euptychia Atalanta.

Euptychia Atalanta, *Butler, Proc. Zool. Soc.* p. 474. n. 50 pl. 39. fig. 12 (1866).

Venezuela. B.M.

Var. *Butl. ibid.* (1866).

Para. B.M.

51. Euptychia Hermes.

Papilio Hermes, *Fabricius, Ent. Syst.* iii. pt. 1. p. 158. n. 486 (1793), but not of *Herbst*.

Satyrus Hermes, var., *Godart, Enc. Méth.* ix. p. 487. n. 33 (part.) (1819).

Neonympha Hermes, *E. Doubleday, List Lep. Brit. Mus.* p. 138 (1844).

Euptychia Hermes, *Butler, Proc. Zool. Soc.* p. 475. n. 51 (1866).

Oreas (strigata) Canthe, *Hübner, Samml. exot. Schmett.* Bd. i. pl. 87. figs. 1-4 (1806).

Euptychia Canthe, *Westwood in Gen. Diurn. Lepid.* p. 373. n. 22 (1851).

Pernambuco, Rio Janeiro, Para, Nicaragua, Honduras. B.M.

52. Euptychia undulata.

Euptychia undulata, *Butler, Proc. Zool. Soc.* p. 475. n. 52, pl. 39. fig. 13 (1866).

Para. B.M.

53. Euptychia binalinea.

Euptychia binalinea, *Butler, Proc. Zool. Soc.* p. 475. n. 53, pl. 39. fig. 14 (1866).

Venezuela, Pernambuco. B.M.

54. Euptychia Poltys.

Neonympha Poltys, *Prittwitz, Entom. Zeit. herausg. von dem ent. Vereine, Stettin*, p. 311 (1865).

Euptychia Poltys, *Butler, Proc. Zool. Soc.* p. 476. n. 54, descr. (1866).

Corcovado (Prittwitz).

55. Euptychia Acmenis.

(*Oreas strigata*) Megisto Acmenis, *Hübner, Samml. exot. Schmett. Zutr.* figs. 233, 234 (1806).

Megisto Acmenis, *Hübner, Verz. bek. Schmett.* p. 54. n. 510 (1816).

Neonympha Acmenis, *Westwood in Gen. Diurn. Lepid.* p. 375. n. 14 (1851).

Euptychia Acmenis, *Butler, Proc. Zool. Soc.* p. 476. n. 55 (1866).

Baltimore (Hübner), Southern States? (Edwards in litt.).

56. Euptychia Eoüs.

Euptychia Eoüs, *Butler, Proc. Zool. Soc.* p. 477. n. 56, pl. 39. fig. 15 (1866).

Brazil, Para.

B.M.

57. Euptychia Phares.

Satyrus Phares, *Godart, Enc. Méth.* ix. pp. 464 & 491. n. 47 (1819).

Neonympha Phares, *Westwood in Gen. Diurn. Lepid.* p. 375. n. 18, pl. 66. fig. 4 (1851).

Euptychia Phares, *Butler, Proc. Zool. Soc.* p. 477. n. 57 (1866).

Brazil, Venezuela.

B.M.

58. Euptychia Pharella.

Euptychia Pharella, *Butler, Proc. Zool. Soc.* p. 478. n. 58, pl. 39. fig. 16 (1866).

Rio Janeiro.

B.M.

59. Euptychia Harmonia.

Euptychia Harmonia, *Butler, Proc. Zool. Soc.* p. 478. n. 59, pl. 39. fig. 17 (1866).

Quito.

B.M.

60. Euptychia Epinephele.

Neonympha Epinephele, *Felder, Reise der Fregatte Novara*, iii. p. 476. n. 817 (1867).

Mexico (Felder).

61. Euptychia Jovita.

Neonympha Jovita, *Felder, Reise der Fregatte Novara*, iii.
p. 477. n. 819 (1867).

Bogota (Felder).

This species seems from the description to be nearly allied to *E. Harmonia*.

62. Euptychia Alcinoë.

Neonympha Alcinoë, *Felder, Reise der Fregatte Novara*, iii.
p. 477. n. 820 (1867).

Bogota, Venezuela (Felder).

Like the preceding, but constantly larger.

63. Euptychia Sylvina.

Neonympha Sylvina, *Felder, Reise der Fregatte Novara*, iii.
p. 478. n. 821 (1867).

Bahia (Felder).

Closely allied to *E. Alcinoë*, *Felder*. The male from which the description was taken had the discoidal cell longer than in *Alcinoë*. If this was the only distinction, it may have been a malformation.

64. Euptychia Numeria.

Neonympha Numeria, *Felder, Reise der Fregatte Novara*, iii.
p. 478. n. 822 (1867).

Bahia (Felder).

Seems chiefly to differ from the preceding in form and size.

65. Euptychia Numilia.

Neonympha Numilia, *Felder, Reise der Fregatte Novara*, iii.
p. 478. n. 823 (1867).

Bogota (Felder).

"Not larger than *Phares*. The wings, especially the hind wing, considerably shorter than in the preceding species."

66. Euptychia Pompilia.

Neonympha Pompilia, *Felder, Reise der Fregatte Novara*, iii.
p. 479. n. 824 (1867).

Bogota, Mexico (Felder).

According to *Felder*, larger than the preceding species, to which it is allied.

67. Euptychia peloria.

Neonympha peloria, *Felder, Reise der Fregatte Novara*, iii.
p. 479. n. 825 (1867).

Venezuela (Felder).

. "Size and form of wings as in *Sylvina*."

68. Euptychia Manasses.

Neonympha Manasses, *Felder, Reise der Fregatte Novara*, iii.
p. 479. n. 826 (1867).

Bahia (Felder).

Larger and broader than the preceding, of which it is probably a variety. If Dr. Felder had given a figure of one of the foregoing species, it would have been much more easy to understand his complex comparative descriptions; but without such aid it is next to impossible to make anything of them.

69. Euptychia yphthima.

Neonympha yphthima (*sic*), *Felder, Reise der Fregatte Novara*,
iii. p. 481. n. 830 (1867).

Bahia (Felder).

This insect, according to Dr. Felder, reminds one of the larger species of *Yphthima* in the form of its wings. The description of the species will very nearly fit *E. nebulosa*.

70. Euptychia Phineus.

Euptychia Phineus, *Butler, Proc. Zool. Soc.* p. 478. n. 60,
pl. 39. fig. 18 (1866).

Neonympha Eugenia, *Felder, Reise der Fregatte Novara*, iii.
p. 476. n. 818 (1867).

Venezuela.

B.M.

71. Euptychia nebulosa.

Euptychia nebulosa, *Butler, Proc. Zool. Soc.* p. 479. n. 61
(1866), pl. 12. fig. 2 (1867).

Venezuela.

B.M.

72. Euptychia Nossis.

Euptychia Nossis, *Hewitson, Exot. Butterf.* iii. pl. 49. fig. 1
(1862).

Quito.

Colls. W. W. Saunders and W. C. Hewitson.

73. Euptychia Pronophila.

Euptychia Pronophila, *Butler, Proc. Zool. Soc.* p. 107. n. 6,
pl. 12. fig. 20 (1867).

Rio Janeiro.

Coll. Hewitson.

74. Euptychia hilara.

Neonympha hilara, *Felder, Reise der Fregatte Novara*, iii. p. 485. n. 837 (1867).

Bogota (Felder).

The description of this insect seems to come very near indeed to *Mollina*. According to Felder, it differs in being larger and the hind wing not rounded.

75. Euptychia Saturnus.

Euptychia Saturnus, *Butler, Proc. Zool. Soc.* p. 479. n. 64, pl. 39. fig. 19 (1866).

Neonympha Laccine, *Felder, Reise der Novara*, iii. p. 480. n. 828 (1867).

Venezuela, Brazil.

B.M.

76. Euptychia Vesta.

Euptychia Vesta, *Butler, Proc. Zool. Soc.* p. 479. n. 65, pl. 39. fig. 20 (1866).

Neonympha Griphe, *Felder, Reise der Novara*, iii. p. 480. n. 827 (1867).

Bogota (Felder), Venezuela.

B.M.

77. Euptychia Enyo.

Euptychia Enyo, *Butler, Proc. Zool. Soc.* p. 480. n. 66, pl. 39. fig. 22 (1866).

Cuenca.

B.M.

78. Euptychia Mollina.

(*Oreas strigata*) *Euptychia Mollina*, *Hübner, Samml. exot. Schmett. Zuträge*, figs. 105, 106 (1806).

Euptychia Mollina, *E. Doubleday, List. Lep. Brit. Mus.* App. p. 31 (1847).

Euptychia Molina (sic), *Hübner, Verz. bek. Schmett.* p. 54. n. 509 (1816).

Para, Ega.

B.M.

79. Euptychia Westwoodii.

Euptychia Westwoodii, *Butler, Proc. Zool. Soc.* p. 481. n. 68 (1866), pl. 12. fig. 3 (1867).

Neonympha Mollina, *Westwood, Gen. Diurn. Lepid.* p. 375. n. 17 (1851).

Honduras.

B.M.

Var. *Butl. ibid.*

Venezuela, Peru.

B.M.

80. *Euptychia picea*.

Euptychia picea, *Bulter, Proc. Zool. Soc.* p. 481. n. 69 (1866),
pl. 12. fig. 6 (1867).

Ega. B.M.

Var. *a.* *Bulter, ibid.*

Brazil. B.M.

Var. *b.* *Bulter, ibid.*

Tapajos. B.M.

Var. *c.* *Bulter, ibid.*

Para. Colls. Salvin and B.M.

81. *Euptychia Mæpius*.

Satyrus Mæpius, *Godart, Enc. Méth.* ix. pp. 464 & 490. n. 45
(1819).

Euptychia Mæpius, *Westwood in Gen. Diurn. Lepid.* p. 373.
n. 21 (1851); *Bulter, descr.* (1866).

Guiana (Godart). Coll. Boisduval.

This species is closely allied to *E. picea*. In my monograph of the genus it is considered to be closely allied to *E. Lethe*; but the type in Dr. Boisduval's Collection shows its true position to be as above.

82. *Euptychia Lysidice*.

♂. *Papilio Lysidice*, *Cramer, Pap. Exot.* ii. pl. 169. figs. C, D
(1779); also of *Fabricius*.

(*Oreas strigata*) *Lysidice*, *Hübner, Samml. exot. Schmett.* Bd. i.
pl. 86. figs. 1-4 (1806).

Erycina Lysidice, *Godart, Enc. Méth.* ix. p. 583. n. 92 (1819).

Euptychia Lysidice, *Westwood in Gen. Diurn. Lepid.* p. 373.
n. 12 (1851).

Euptychia Lysidice (sic), *Hübner, Verz. bek. Schmett.* p. 54.
n. 507 (1816).

♀. *Papilio Doris*, *Cramer, Pap. Exot.* i. pl. 8. figs. B, C (1779);
also of *Fabricius* (1793).

Euptychia Doris, *Westwood in Gen. Diurn. Lepid.* p. 373. n. 13
(1851).

Surinam (Cramer), Para, Tapajos, Ega, Nauta, Brazil. B.M.

83. *Euptychia glaucina*.

Euptychia glaucina, *H. W. Bates, Ent. Month. Mag.* i. p. 202.
n. 76 (1865).

Guatemala. Coll. Salvin.

84. **Euptychia agrota.**

Euptychia agrota, *Butler, Proc. Zool. Soc.* p. 482. n. 72 (1866),
pl. 11. fig. 2 (1867).

Para (♂ ♀, Coll. Hewitson). ♀, B.M.

85. **Euptychia pilata.**

Euptychia pilata, *Butler, Proc. Zool. Soc.* p. 483. n. 73, pl. 40.
fig. 3 (1866).

Ega. B.M.

86. **Euptychia Brixiola.**

Euptychia Brixiola, *Butler, Proc. Zool. Soc.* p. 483. n. 74, pl. 40.
fig. 9 (1866).

Para. Col. Bates.

87. **Euptychia Brixius.**

Satyrus Brixius, *Godart, Enc. Méth.* ix. pp. 464 & 490. n. 42
(1819).

Euptychia Brixius, *Westwood in Gen. Diurn. Lepid.* p. 374.
n. 24 (1851); also *Butl. descr.* (1866).

Brazil (Godart).

88. **Euptychia celestis.**

Euptychia celestis, *Butler, Proc. Zool. Soc.* p. 484. n. 76, pl. 40
fig. 5 (1866).

Ega. ♂ ♀, B.M.

Var. *Butl. ibid.* (1866). ? *Urania* ♀.

Para. ♀, Coll. Salvin.

89. **Euptychia Urania.**

Euptychia Urania, *Butler, Proc. Zool. Soc.* p. 484. n. 77, pl. 40.
fig. 6 (1866).

Cameta. ♂, Coll. Bates.

90. **Euptychia Lea.**

♀. *Papilio* Lea, *Cramer, Pap. Exot.* ii. pl. 151. figs. C, D
(1779).

Satyrus Lea, *Godart, Enc. Méth.* ix. pp. 464 & 492. n. 50 (1819).

Euptychia Lea, *Westwood in Gen. Diurn. Lepid.* p. 373. n. 17
(1851).

Euptychia Ebusa (part.), *Hübner, Verz. bek. Schmett.* p. 54.
n. 515 (1816).

Surinam (Cramer).

91. **Euptychia Junia.**

Papilio Junia, *Cramer, Pap. Exot.* iv. pl. 292. figs. D, E (1782).

Euptychia Junia, *Hübner, Samml. exot. Schmett. Beiträge*, figs. 627, 628 (1806).

Satyrus Lea (part.), *Godart, Enc. Méth.* ix. p. 492. n. 50 (1819).

Surinam (*Cramer*), Brazil. ♂, B.M.

Var. *Butler, Proc. Zool. Soc.* p. 485. n. 79 (1866).

Para. ♀, B.M.

92. **Euptychia Philippa.**

Euptychia Philippa, *Butler, Proc. Zool. Soc.* p. 485. n. 80 (1866), pl. 11. fig. 3 (1867).

Ega. ♂ ♀, B.M.

93. **Euptychia Antonoë.**

Papilio Antonoë, *Cramer, Pap. Exot.* i. pl. 59. figs. E, F (1779).

Euptychia Antonoë, *Butler, Proc. Zool. Soc.* p. 486. n. 81 (1866).

Euptychia Hermessa, *Hübner, Verz. bek. Schmett.* p. 54. n. 508 (1816).

Satyrus Hermes (part.), *Godart, Enc. Méth.* ix. p. 487 (1819).

♂ (Coll. Salvin). ♀, Para, Bolivia. B.M.

94. **Euptychia Sabina.**

Neonympha Sabina, *Felder, Reise der Fregatte Novara*, iii. p. 483. n. 834 (1867).

Cayenne (*Felder*).

Dr. Felder considers that this species, from the shape of its wings, will form a link between his *Terentia* (which, if correctly made out, is only a variety of *ocypete*) and the *Junia* group; but the latter insects have the ocelli all black beneath, whilst in the description of *Sabina* we read "*ocellis bipupillatis, tertio et quarto completis sed fuscis.*"

95. **Euptychia gigas.**

Euptychia gigas, *Butler, Proc. Zool. Soc.* p. 486. n. 82, pl. 40. fig. 7 (1866).

Mexico. B.M.

96. **Euptychia Libye.**

Papilio Libye, *Linnaeus, Syst. Nat.* ii. p. 772. n. 146 (1766); *Sulzer, Gesch.* pl. 17. fig. 7. *P. Lybie* (sic), *ejusd.* p. 145 (1776).

Var. ?. *Papilio Hermes*, *Herbst, Natursyst. Schmett.* viii. p. 68. n. 32, tab. 192. fig. 4 (1796).

Neonympha? *Antonoë*, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 138 (1844); *Neonymph. Anton. Westwood* (1851).

Euptychia Antonoë, *Westwood*, *Gen. Diurn. Lepid.* p. 374. n. 30 (1851).

Neonympha Harpyia, *Felder*, *Reise der Novara*, iii. p. 482. n. 832 (1867).

Pernambuco, Bahia, Rio, Honduras, Para, Quito. B.M.

Var. *Butler*, *Proc. Zool. Soc.* p. 486. n. 83 (1851).

Quito. B.M.

97. ***Euptychia libyoidea*.**

Euptychia libyoidea, *Butler*, *Proc. Zool. Soc.* p. 487. n. 84 (1866), pl. 11. fig. 13 (1867).

Nicaragua. B.M.

98. ***Euptychia obscura*.**

Euptychia obscura, *Butler*, *Proc. Zool. Soc.* p. 487. n. 85 (1866), pl. 11. fig. 9 (1867).

Bolivia. B.M.

99. ***Euptychia Necys*.**

Satyrus Necys, *Godart*, *Enc. Méth.* ix. pp. 466 & 511. n. 100 (1819).

Euptychia Necys, *Westwood in Gen. Diurn. Lepid.* p. 373. n. 14 (1851); *Bull. descr.* (1866).

Euptychia vastata, *Butler*, *Proc. Zool. Soc.* p. 487. n. 86 (1866), pl. 12. fig. 11 (1867).

Brazil (Godart), Venezuela (Westwood), Rio Grande (Coll. Bates).

100. ***Euptychia Quantius*.**

Satyrus Quantius, *Godart*, *Enc. Méth.* ix. pp. 463 & 487. n. 31 (1819).

Neonympha Quantius, *Westwood in Gen. Diurn. Lepid.* p. 376. n. 23 (1851).

Euptychia Quantius, *Butler*, *descr. Proc. Zool. Soc.* p. 488. n. 87 (1866).

Brazil (Godart), Venezuela. B.M.

? Var. *a. Bull. ibid.*

Rio Grande. Coll. Bates.

Var. *b. Bull. ibid.*

Rio Janeiro. B.M.

101. *Euptychia rustica*, sp. nov. Pl. I. fig. 4.

♂. Alæ supra rufo-fuscæ, posticæ margine externo minime undulato.

Alæ subtus velut supra fuscæ, area apicali paulum dilutior; linea discali arcuata undulata, altera submarginali sinuata, tertia marginali indistinctissima, fuscis obscurioribus; margine ipso nigro, ciliis cinereo-fuscis brevibus: anticæ macula squamisque nonnullis subapicalibus fulvo-lacteis albisve: posticæ punctis quinque submarginalibus albis.

Exp. alar. unc. 2.

Bolivia.

B.M.

Allied to *Quantius*; differs in the form of the central lines and large subapical spot on the underside.

102. *Euptychia Polyphemus*.

Euptychia Polyphemus, Butler, *Proc. Zool. Soc.* p. 488. n. 88 (1866), pl. 12. fig. 5 (1867).

Bogota.

B.M.

Section EUPTYCHIA, true, Hübner.

Typical species *E. Herse*.

103. *Euptychia Cluena*.

Papilio (Nymph.) *Cluena*, Drury, *Ill.* iii. p. 9, pl. 7. figs. 5, 6 (1782).

Euptychia Cluena, Westwood in *Gen. Diurn. Lepid.* p. 373. n. 3 (1851); see also *P. Z. S.* p. 489 (1866).

Papilio Clueria, Fabricius, *Ent. Syst.* iii. pt. 1. p. 229. n. 716 (1793).

Satyrus Clueria, Godart, *Enc. Méth.* ix. p. 492. n. 51 (1819).

Euptychia Clueria, E. Doubleday, *List Lep. Brit. Mus.* i. p. 122 (1844).

Brazil.

? B.M.

Note.—It is doubtful whether the species in the National Collection is the true *Cluena* of Drury, as stated in the "Monograph of *Euptychia*," *P. Z. S.* (1866).

104. *Euptychia sericeella*.

Euptychia sericeella, H. W. Bates, *Ent. Month. Mag.* i. p. 202. n. 75 (1865).

Guatemala.

(Coll. Salvin).

105. *Euptychia Ebusa*.

Papilio Ebusa, Cramer, *Pap. Exot.* iv. pl. 292. figs. F, G (1782).

Euptychia Ebusa, Hübner, *Verz. bek. Schmett.* p. 54. n. 515 (1816).

Papilio aranea, *Fabricius, Ent. Syst.* iii. pt. 1. p. 97. n. 299 (1793).

Satyrus aranea, *Godart, Enc. Méth.* ix. pp. 464 & 492. n. 49 (1819).

Pernambuco, Venezuela, Bahia, Para.

B.M.

106. ***Euptychia Byses*.**

Satyrus Byses, *Godart, Enc. Méth.* ix. pp. 466 & 496. n. 67 (1819).

Euptychia Byses, *Westwood, Gen. Diurn. Lepid.* p. 373. n. 9 (1851); *Butler, Proc. Zool. Soc.* pl. 12. fig. 15 (1867).

Rio Janeiro.

B.M.

Var. *Butler, ibid.* (1866) (possibly the *E. Necys* of Godart: it does not, however, exactly suit the description of that species).

Brazil.

B.M.

107. ***Euptychia Chloris*.**

Papilio Chloris, *Cramer, Pap. Exot.* iv. pl. 293. figs. A, B (1782).

Euptychia Chloris, *Westwood in Gen. Diurn. Lepid.* p. 373. n. 20 (1851).

Euptychia Chlorimene, *Hübner, Verz. bek. Schmett.* p. 54. n. 514 (1816).

Satyrus Tolumnia ♀, *Godart, Enc. Méth.* ix. p. 491. n. 48 (1819).

Brazil, Bahia, Para.

B.M.

108. ***Euptychia Herse*.**

Papilio Herse, *Cramer, Pap. Exot.* i. pl. 10. figs. C, D (1779).

Euptychia Herse, *Hübner, Verz. bek. Schmett.* p. 54. n. 502 (1816).

Satyrus Herseis, *Godart, Enc. Méth.* ix. pp. 465 & 495. n. 60 (1819).

Euptychia Herseis, *Westwood in Gen. Diurn. Lepid.* p. 373. n. 11 (1851).

Brazil, Venezuela, Para.

B.M.

Var. *Butler, Proc. Zool. Soc.* p. 490. n. 94 (1866).

Para, Tapajos.

Coll. Bates and B.M.

109. ***Euptychia Callichloris*.**

Euptychia Callichloris, *Butler, Proc. Zool. Soc.* p. 490. n. 95, pl. 40. fig. 10 (1866).

(Brazil), Ega.

Coll. Bates.

110. *Euptychia Hewitsonii*.

Euptychia Hewitsonii, *Butler*, *Proc. Zool. Soc.* p. 491. n. 96,
pl. 40. fig. 4 (1866).

Para. B.M.

Var. *Butler*, *ibid.* (1866).

Ega. B.M.

111. *Euptychia Agatha*.

Euptychia Agatha, *Butler*, *Proc. Zool. Soc.* p. 492. n. 97, pl. 40.
fig. 8 (1866).

Para. B.M.

112. *Euptychia Tolumnia*.

Papilio Tolumnia, *Cramer*, *Pap. Exot.* ii. pl. 130. figs. F, G
(1779); also of *Fabricius* (1793).

Euptychia Tolumnia, *Hübner*, *Verz. bek. Schmett.* p. 54. n. 513
(1816).

Satyrus Tolumnia, *Gedart*, *Enc. Méth.* ix. pp. 464 & 491. n. 48
(1819).

Guiana, Bahia (Westwood). Coll. Hewitson.

113. *Euptychia Ayaya*.

Euptychia Ayaya, *Butler*, *Proc. Zool. Soc.* p. 492. n. 99, pl. 40.
fig. 11 (1866).

Tapajos. Coll. Bates.

114. *Euptychia tricolor*.

Euptychia tricolor, *Hewitson*, *Ann. & Mag. Nat. Hist.* vi. p. 440
(1850); *Gen. Diurn. Lepid.* pl. 65. fig. 3 (1851).

Tapajos, Fonteboa. ♂, B.M.

115. *Euptychia Batesii*.

Euptychia Batesii, *Butler*, *Proc. Zool. Soc.* p. 493. n. 101. pl. 40.
fig. 16 (1866).

Tapajos. ♂, B.M.

116. *Euptychia Iris*.

Neonympha Iris, *Felder*, *Reise der Fregatte Novara*, iii. p. 483.
n. 835 (1867).

Bogota (Felder).

Seems to be closely allied to my *E. Ayaya*.

117. **Euptychia Nortia.**

Euptychia Nortia, *Hewitson, Exot. Butterf.* iii. pl. 44. n. 2. fig. 2 (1862).

Tapajos.

B.M.

118. **Euptychia Gera.**

Euptychia Gera, *Hewitson, Ann. & Mag. Nat. Hist.* vi. p. 439 (1850); *Gen. Diurn. Lepid.* pl. 63. fig. 4 (1851).

Tapajos.

Colls. Bates and Hewitson.

119. **Euptychia Metagera.**

Euptychia Metagera, *Butler, Proc. Zool. Soc.* p. 494. n. 104 (1866), pl. 11. fig. 4 (1867).

Upper Amazons.

Coll. Bates.

120. **Euptychia hiemalis.**

Euptychia hiemalis, *Butler, Proc. Zool. Soc.* p. 494. n. 105 (1866), pl. 12. fig. 4 (1867).

Amazons.

B.M.

Section NEONYMPHA, true, *Hübner*.

Typical species *Neonympha (Helicta) Phocion*.

121. **Euptychia Furina.**

Euptychia Furina, *Hewitson, Exot. Butterf.* iii. pl. 44. fig. 4 (1862).

Tapajos.

♂, B.M.

122. **Euptychia Junonia.**

Euptychia Junonia, *Butler, Proc. Zool. Soc.* p. 495. n. 107 (1866), pl. 11. fig. 11 (1867).

Tapajos.

B.M.

123. **Euptychia gemmula.**

Euptychia gemmula, *Butler, Proc. Zool. Soc.* p. 495. n. 108 (1866), pl. 11. fig. 7 (1867).

♂, Rio Janeiro (Coll. Bates), Brazil.

♀, B.M.

124. **Euptychia Doxes.**

Satyrus Doxes, *Godart, Enc. Méth.* ix. pp. 465 & 493. n. 54 (1819).

Euptychia Doxes, *Westwood in Gen. Diurn. Lepid.* p. 374.
n. 28 (1851); descr. *Proc. Zool. Soc.* p. 496. n. 109 (1866).
Brazil (Godart).

125. ***Euptychia Erycina*.**

Euptychia Erycina, *Butler, Proc. Zool. Soc.* p. 496. n. 110
(1866), pl. 11. fig. 6 (1867).
Brazil. ♂, B.M.

126. ***Euptychia Latia*.**

Euptychia Latia, *Butler, Proc. Zool. Soc.* p. 496. n. 111, pl. 40.
fig. 14 (1866).
Bahia. ♀, B.M.

127. ***Euptychia cosmophila*.**

♂ (*Oreas fimbriata*) *Neonympha cosmophila*, *Hübner, Sammlung
exot. Schmett. Beiträge*, figs. 255, 256 (1806).
Neonympha cosmophila, *Westwood in Gen. Diurn. Lepid.*
p. 375. n. 15 (1851).
Euptychia cosmophila, *Butler, Proc. Zool. Soc.* p. 497. n. 112
(1866).
♀, *Satyrus argenteus*, *Swainson, Zool. Ill.* i.-iii. pl. 159
(1820-23).
Euptychia Pagyris (*syn.*), *Westwood in Gen. Diurn. Lepid.*
p. 373. n. 8 (1851).
Brazil, Bahia. B.M.

128. ***Euptychia Pagyris*.**

Satyrus Pagyris, *Godart, Enc. Méth.* ix. pp. 464 & 491. n. 46
(1819).
Euptychia Pagyris, *Westwood in Gen. Diurn. Lepid.* p. 373.
n. 8 (1851); *Butler, Proc. Zool. Soc.* p. 497. n. 113 (1866),
pl. 11. fig. 1 (1867).
Brazil. Coll. Hewitson.

129. ***Euptychia Itonis*.**

Euptychia Itonis, *Hewitson, Exot. Butterf.* iii. pl. 44. fig. 3
(1862).
Para. B.M.

130. ***Euptychia Clorimena*.**

Papilio Clorimena, *Stoll, Pap. Exot.* pl. 13. figs. 2, 2b (1791).
Euptychia Clorimena, *Butler, Proc. Zool. Soc.* p. 498. n. 115
(1866).

Neonympha Clorimene (*sic*), *Herrich-Schäffer, Ind. Syst. Lep. Exot.* p. 59. n. 12 (1864).

Neonympha Clerimene (*sic*), *Hübner, Verz. bek. Schmett.* p. 65. n. 624 (1816); also of *Westwood* (1851).

Surinam (Stoll), Cayenne.

Colls. Salvin & Hewitson.

131. *Euptychia Salvini*.

Euptychia Salvini, *Butler, Proc. Zool. Soc.* p. 498. n. 116, pl. 40. fig. 13 (1866).

Panama.

Coll. Salvin.

132. *Euptychia Phocion*.

Papilio Phocion, *Fabricius, Ent. Syst.* iii. pt. 1. p. 218. n. 683 (1793).

Satyrus Phocion, *Godart, Enc. Méth.* ix. pp. 465 & 494. n. 59 (1819).

Neonympha Phocion, *Westwood in Gen. Diurn. Lepid.* p. 375. n. 6 (1851).

United States.

B.M.

Var. AREOLATA.

Papilio areolatus, *Smith & Abbot's Lep. Georgia*, p. 25, tab. 13 (1797).

Satyrus areolatus, *Godart, Enc. Méth.* ix. pp. 465 & 494. n. 58 (1819); also of *Boisduval & Leconte* (1833).

Neonympha areolatus, *Westwood in Gen. Diurn. Lepid.* p. 375. n. 6 (1851).

Euptychia areolata, *Butler, Proc. Zool. Soc.* p. 498. n. 117 (1866).

Oreas (fimbriata) Helicta, *Hübner, Samml. exot. Schmett.* i. pl. 95 (1806).

Neonympha Helicta, *Hübner, Verz. bek. Schmett.* p. 65. n. 622 (1816).

Georgia, United States.

Coll. Edwards.

133. *Euptychia Pyracmon*.

Euptychia Pyracmon, *Buller, Proc. Zool. Soc.* p. 499. n. 118 (1866), pl. 11. fig. 10 (1867).

Mexico.

B.M.

134. *Euptychia gemma*.

Neonympha gemma, *Hübner, Sammlung exot. Schmett. Zuträge*, figs. 7, 8 (1806); also of *Westwood* (1851).

Satyrus gemma, *Boisduval & Leconte, Hist. des Léop.* pl. 62. figs. 1-3, 4, 5 (metamorph.) (1833).

Polochie Valley (Coll. Salvin), United States.

B.M.

135. Euptychia Saundersii.

Euptychia Saundersii, *Butler, Proc. Zool. Soc.* p. 500. n. 120,
pl. 40. fig. 17 (1866).

Ega (Coll. Bates), Pernambuco.

B.M.

136. Euptychia mima.

Euptychia mima, *Butler, Proc. Zool. Soc.* p. 500. n. 121 (1866),
pl. 12. fig. 7 (1867).

? *Neonympha Antonina*, *Felder, Reise der Novara*, iii. p. 482.
no. 831 (1867).

Tapajos.

Coll. Bates.

137. Euptychia fumata.

Euptychia fumata, *Butler, Proc. Zool. Soc.* p. 109. n. 10, pl. 12.
fig. 14 (1867).

Hab. — ?

Coll. Hewitson.

138. Euptychia Pacarus.

Satyrus Pacarus, *Godart, Enc. Méth.* ix. pp. 465 & 495. n. 61
(1819).

Neonympha Pacarus, *Westwood in Gen. Diurn. Lepid.* p. 375.
n. 4 (1851).

Euptychia Pacarus, *Butler, Proc. Zool. Soc.* p. 501. n. 123 (descr.)
(1866).

Brazil (Godart).

139. Euptychia insignis.

Euptychia insignis, *Butler, Proc. Zool. Soc.* p. 501. n. 124,
pl. 40. fig. 12 (1866).

Hab. — ?

Coll. Bates.

140. Euptychia Erichtho.

Euptychia Erichtho, *Butler, Proc. Zool. Soc.* p. 501. n. 125,
pl. 40. fig. 15 (1866).

Pernambuco, Rio, Ega, Para.

B.M.

Undetermined Species.

141. Euptychia ?? Peribæa.

Papilio Peribæa, *Fabricius, Ent. Syst.* iii. 1. p. 234. n. 730
(1793).

Satyrus Peribæa, *Godart, Enc. Méth.* ix. pp. 463 & 486. n. 29
(1819).

Neonympha Peribæa, *Westwood in Gen. Diurn. Lepid.* p. 376.
n. 24 (1851).

Surinam (Fabricius).

The part of the 'Proceedings Philad.' in which the following are described has not yet been obtained:—

142. *Euptychia Lupita*.

Neonympha Lupita, *Reakirt, Proc. Acad. Nat. Sci. Philad.*
p. 331 (1866).

Mexico (Reakirt).

143. *Euptychia Xicaque*.

Neonympha Xicaque, *Reakirt, Proc. Acad. Nat. Sci. Philad.*
p. 336 (1866).

Mexico (Reakirt).

Genus 7. CÆNONYMPHA, *Hübner* (1816).

Typical Species *Cænonympha (Ædipe) Geticus*.

1. *Cænonympha Geticus*.

♀. *Papilio* (pleb. rural.) *Geticus*, *Esper, Schmett.* i. pt. 2.
tab. 102, cont. 57. fig. 2; ♂, tab. 107, cont. 62. fig. 5 (1777).

Papilio Iphigenus, *Herbst, Natursyst. Schmett.* viii. p. 108. n. 60,
tab. 198. figs. 5–8 (1796).

Pyrenees, Sweden.

♂ ♀, B.M.

Var. ŒDIPPUS.

♂. *Papilio Œdippus*, *Fabricius, Mant. Ins.* ii. p. 31. n. 335
(1787); *Ent. Syst.* iii. pt. 1. p. 218. n. 681 (1793); also of
Herbst.

Papilio Œdippe, *Borkhausen, Naturgesch. eur. Schmett.* ii. p. 205.
n. 23–24 (1788).

Papilio Œdipus (*sic*), *Ochsenheimer, Schmett.* v. *Eur.* i. p. 315.
n. 71 (1807).

Cænonympha Œdipus (*sic*), *E. Doubleday, List Lep. Brit. Mus.*
i. p. 143 (1844); also of *Westwood* (1851).

Hipparchia Œdipus, *Eversmann, Faun. Lepid. Volgo-Ural.*
p. 36. n. 29 (1844).

Cænonympha Œdipe (*sic*), *Hübner, Verz. bek. Schmett.* p. 65.
n. 626 (1816).

Var. Satyrus Œdipus (*sic*), *Godart, Enc. Méth.* ix. p. 544. n. 169
(1819); *Lép. de France*, ii. p. 141. n. 53, pl. 19. figs. 5, 6
(1822).

Var. Papilio Miris, Fabricius, Ent. Syst. Suppl. v. p. 429. n. 693, 694 (1798).

Papilio Pylarge, Hübner, Eur. Schmett. i. pl. 52. figs. 245, 246 (1805).

Russia.

♂ ♀, B.M.

C. Edippus appears to be the Russian form of *Geticus*; it differs from it chiefly in having less conspicuous ocelli and bands on the underside. The females differ much more than the males; the number of ocelli in the front wings of these two forms varies in our specimens from five to none.

2. *Cænonympha Hero.*

Papilio Hero, Linnæus, Fauna Suecica, ed. 2. p. 274. n. 1047 (1761); Syst. Nat. ii. p. 793. n. 255 (1766); also of Fabricius, Syst. Ent., but not Ent. Syst., and of Borkhausen, Herbst, and Hübner.

Papilio (pleb. rural.) Hero, Esper, Schmett. i. pt. 1. tab. 22. fig. 4 (1777).

Maniola Hero, Schrank, Fauna Boica, i. pt. 2. p. 179. n. 1310 (1801).

Satyrus Hero, Godart, Enc. Méth. ix. p. 544. n. 170 (1819).

Hipparchia Hero, Stephens, Ill. Haust. i. p. 68. sp. 18 (1828); also of Eversmann (1844).

Cænonympha Hero, E. Doubleday, List. Lep. Brit. Mus. i. p. 143 (1844); also of Westwood (1851).

Papilio Sabacus, Fabricius, Syst. Ent. p. 530. n. 371 (1775).

France, Sicily ?

B.M.

3. *Cænonympha Amyntas.*

♀. *Papilio Amyntas, Poda, Ins. Mus. Græc. p. 79. n. 52 (1761).*

Papilio Hero (part.), Linnæus, Fauna Suecica, ed. 2. p. 274. n. 1047 (1761); also of Fabricius, Ent. Syst., but not Syst. Ent.

Papilio Iphis, Denis, Wiener Verz. p. 321. n. 25 (1775); Hübner, Eur. Schmett. i. tab. 53. figs. 249-251 (var.) (1805).

Satyrus Iphis, Godart, Enc. Méth. ix. p. 545. n. 171 (1819); Léop. de France, ii. 145. n. 55, pl. 20. figs. 3, 4 (1822).

Cænonympha Iphis, E. Doubleday, List Lep. Brit. Mus. i. p. 143 (1844).

Hipparchia Iphis, Eversmann, Fauna Lepid. Volgo-Ural. p. 35. n. 27 (1844).

Maniola Manto, Schrank, Fauna Boica, i. pt. 2. p. 180. n. 1313 (1801).

Papilio (pleb. rural.) Tiphon, Esper, Schmett. i. pt. 1. tab. 35. Suppl. xi. figs. 3, 4 (1777).

Papilio Tiphon, Schneider, Syst. Besch. p. 126. n. 63 (1787).

Papilio Glycerion, Borkhausen, Naturgesch. eur. Schmett. i. p. 90. n. 27 (1788).

♀. *Cænonympha Mandane*, *W. F. Kirby, Proc. Ent. Soc. Lond.* p. 85 (1862).

Var. Papilio (Hipparchia) Iphis, Freyer, Neuere Beiträge, iv. p. 118. n. 601, tab. 355. figs. 3, 4 (1842).

Var. Cænonympha Anaxagorus, Assmuss, Bresl. Zeitschr. f. d. Entom. x. p. 9 (1856).

Germany, Polish Ukraine.

♂ ♀, B.M.

♂. *Var. Posticæ subtus grisescentes, ocellis obsoletis.*

Germany.

B.M.

4. *Cænonympha arcanioides*.

Satyrus arcanioides, Pierret, Ann. Soc. Ent. France, t. vi. p. 306, *arcanoides*, pl. 12. fig. 5 (1837).

Cænonympha ascanioides (sic), Westwood in Gen. Diurn. Lepid. p. 397. n. 5 (1851).

Barbary (Pierret).

Local form. ?Papilio (Hipparchia) arcanoides (sic), Freyer, Neuere Beiträge, v. p. 125. n. 816, tab. 457. n. 1 (1845).

Switzerland (Freyer).

B.M.

It is quite likely that the Swiss specimens may be a race of *arcanioides*; they do not, however, agree with the figure of that species in all respects. We have an insect from Algeria, which Mr. Adam White had labelled as this species; but it more nearly resembles *Arcania*, and is probably distinct from both.

5. *Cænonympha Arcania*.

Papilio Arcania, Linnæus, Fauna Suecica, ed 2. p. 273. n. 1045 (1761); also of *Borkhausen* (1788).

Papilio (Nymph. Gemm.) Arcania, Hufnagel, Tab. Berl. Mag. pt. 2. p. 72. n. 31 (1766).

Hipparchia Arcania, Eversmann, Faun. Lepid. Volgo-Ural. p. 37. n. 30 (1844).

Papilio Arcanius, Linnæus, Syst. Nat. ii. p. 791. n. 242 (1766).

Hipparchia Arcanius, Stephens, Ill. Haust. i. p. 69. sp. 19 (1828).

Cænonympha Arcanius, E. Doubleday, List Lepid. Brit. Mus. i. p. 142 (1844).

Cænonympha Ascanius (sic), Westwood in Gen. Diurn. Lepid. p. 397. n. 4 (1851).

Papilio Amyntas, Scopoli, Ent. Carn. p. 174. n. 457 (1763), but not of *Poda*.

Var. Papilio (pleb. rural.) arcanius, Esper, Schmett. i. pt. 1. tab. 21. fig. 4 (1777).

Satyrus Arcanius, Godart, Enc. Méth. ix. p. 546. n. 172 (1819); *Lép. de France*, i. p. 174, pl. 8. fig. 3 (1821).

Papilio Arcania, Hübner, Eur. Schmett. i. tab. 51. figs. 240-242 (1805).

Germany.

B.M.

Var. Papilio Naidion (not Naidon), *Borkhausen, Naturgesch. eur. Schmett.* i. p. 91. n. 28, pl. 1. figs. 1, 2 (1788).

Alæ posticæ subtus fascia albida et linea argentea obsoletis.

Near Darmstadt (Borkhausen).

This is a very variable species. In some specimens the subapical ocellus is wanting; in some there are three distinct ocelli on the upperside of the hind wings: the yellowish band on the underside also varies in outline.

6. *Cænonympha marginalis*, sp. nov.

Alæ supra velut in Arcania.

Anticæ subtus velut in Arcania, sed apice haud pallido, et oculo distincto; posticæ velut in Satyrione, fascia autem alba angustiore, et ocellis multo majoribus.

Exp. alar. unc. 1½.

Algeria.

B.M.

This species was registered as *C. arcanoides*, an insect with which it has but little affinity.

7. *Cænonympha Satyrion*.

Papilio [Dan. fest. (pleb. rural.)] *Satyrion*, *Esper, Schmett.* i. pt. 2. tab. 122. cont. 77. fig. 2 (1777).

Papilio Satyrion, *Ochsenheimer, Schmett. v. Eur. i.* p. 322. n. 74 (1807).

Papilio (*Hipparchia*) *Satyrion*, *Freyer, Neuere Beiträge*, iv. p. 137. n. 626, tab. 367. figs. 1, 2 (1842).

♂. *Satyrus Satyrion*, *Herrich-Schäffer, Eur. Schmett.* i. figs. 289, 290 (1845).

Satyrus Phileus, *Godart, Enc. Méth.* ix. p. 547. n. 174 (1819); *Lép. de France*, ii. p. 143. n. 54, pl. 15. figs. 1, 2 (1822).

Papilio (*Hipparchia*) *Philea*, *Freyer, Neuere Beiträge*, iv. p. 137. n. 627, tab. 367. figs. 3, 4 (1842).

Cænonympha Philea, *E. Doubleday, List Lep. Brit. Mus.* i. p. 142 (1844); also of *Westwood* (1851).

Var. Papilio Philea, *Hübner, Eur. Schmett.* i. tab. 53. figs. 254, 255 (1805).

Germany.

♂ ♀, B.M.

8. *Cænonympha Leander*.

Papilio (pleb. rural.) *Leander*, *Esper, Schmett.* i. pt. 2. tab. 89. cont. 39. fig. 5 (1777).

Papilio Leander, *Fabricius, Ent. Syst.* iii. pt. 1. p. 222. n. 693 (1793); also (part.) *Ochsenheimer*.

Papilio (*Hipparchia*) *Leander*, *Freyer, Neuere Beiträge*, ii. tab. 110. fig. 1 (1836).

Satyrus Leander, *Godart, Enc. Méth.* ix. p. 548. n. 176 (1819); also of *Duponchel*, *Boisduval*, and *Herrich-Schäffer*.

Cænonympha Leander, *E. Doubleday, List Lepid. Brit. Mus.* i. p. 141 (1844).

Hipparchia Leander, *Eversmann, Fauna Lepid. Volgo-Ural.* p. 37. n. 31 (1844).

Papilio Philaidilis, *Borkhausen, Naturgesch. eur. Schmett.* i. p. 93. n. 32 (1788).

Papilio Clite, *Hübner, Eur. Schmett.* i. tab. 103. figs. 526, 527 (1805).

Hungary, Russia.

B.M.

Ochsenheimer has quoted *Amaryllis* as a synonym of this species; it is, however, quite distinct. Westwood, in the 'Genera of Diurnal Lepidoptera,' quotes *Amaryllis* of Cramer, but makes *Amarillis* (sic) of Herbst a distinct species; yet Herbst's figure is copied from that of Cramer, and must therefore be identical with it.

9. *Cænonympha* *Amaryllis*.

Papilio *Amaryllis*, *Cramer, Pap. Exot.* iv. pl. 391. figs. A, B (1782).

Hipparchia *Amaryllis*, *Eversmann in Nouv. Mém. Soc. Imp. Nat. Mosc.* ii. p. 350, tab. 20. figs. 5, 6 (1832).

Papilio (*Hipparchia*) *Amaryllis*, *Freyer, Neuere Beiträge*, iii. pl. 283. figs. 3, 4 (1839).

Satyrus *Amaryllis*, *Herrich-Schäffer, Eur. Schmett.* i. figs. 188, 189, 287, 288 (1845).

Cænonympha *Amaryllis*, *Westwood in Gen. Diurn. Lepid.* p. 397. n. 7 (1851).

Papilio *Amarillis* (sic), *Herbst, Natursyst. Schmett.* viii. p. 35, tab. 186. figs. 1, 2 (1796).

Russia.

B.M.

10. *Cænonympha* *Rinda*.

Cænonympha *Rinda*, *Ménétriés, Bull. Phys. Acad. Pétersb.* xvii. p. 217. n. 18 (1859); *Schrenck's Amurlande*, p. 42. n. 83, tab. 4. fig. 1 (1859).

River Amoor.

Possibly an extreme variety of *Amaryllis*.

11. *Cænonympha* *Corinna*.

Papilio *Corinna*, *Hübner, Eur. Schmett.* i. tab. 105. figs. 536, 537 (1805); also of *Ochsenheimer*.

Papilio (*Hipparchia*) *Corinna*, *Freyer, Neuere Beiträge*, i. tab. 68. fig. 2 (1833).

Satyrus *Corinna*, *Boisduval, Sp. Gén. Léop. Ind.* p. 33. n. 273 (1840).

Cænonympha *Corinna*, *E. Doubleday, List Lep. Brit. Mus.* p. 141 (1844); also of *Westwood*.

Satyrus *Corinnus*, *Godart, Enc. Méth.* ix. p. 540. n. 173 (1819); *Lép. de France*, ii. p. 150. n. 57, pl. 20. figs. 7, 8 (1822).

Papilio Corinna, *Ochsenheimer* (synonym).

Papilio Corinus, *Westwood* (synonym).

Papilio Norax, *Bonelli* in *Mem. Accad. Torino*, xxx. t. 2. fig. 2 (1826).

Corsica.

B.M.

12. *Cænonympha Thyrsis*.

Papilio (*Hipparchia*) *Thyrsis*, *Freyer*, *Neuere Beiträge*, v. p. 157. n. 860, tab. 475. fig. 1 (1845).

Satyrus Thyrsis, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 297–300 (1845).

Cænonympha Thyrsis, *Westwood* in *Gen. Diurn. Lepid.* p. 397. n. 10 (1851).

Turkey in Europe?

B.M.

13. *Cænonympha Dorus*.

Papilio (pleb. rural.) *Dorus*, *Esper*, *Schmett.* i. pt. 2. tab. 78, cont. 28. fig. 1 (1777).

Papilio Dorus, *Herbst*, *Natursyst. Schmett.* viii. tab. 188. figs. 5–8 (1796); also of *Schneider & Ochsenheimer*.

Satyrus Dorus, *Godart*, *Enc. Méth.* ix. p. 547. n. 175 (1819); *Lép. de France*, ii. p. 148. n. 56, pl. 20. figs. 5, 6 (1822); also of *Boisduval*.

Cænonympha Dorus, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 141 (1844); also of *Westwood*.

Papilio Dorion, *Hübner*, *Eur. Schmett.* i. tab. 52. figs. 247, 248 (1805), *Dorilis*, text, p. 42. n. 77.

Papilio Dorilis, *Borkhausen*, *Naturgesch. eur. Schmett.* i. p. 93. n. 31 (1788).

Papilio Palemon, *Ernst*, *Pap. Eur.* i. pl. 68, *Suppl.* xiv. fig. 57 (1779).

Papilio Lizetta, *Cramer*, *Pap. Exot.* iv. pl. 373. figs. F, G (1782).

Papilio Hanibal, *Herbst*, *Natursyst. Schmett.* viii. p. 48. n. 22 (1796).

South Europe, Pyrenees.

♂ ♀, B.M.

Var. Alæ supra minime fulvescentes, ocellis minus distinctis.

Germany.

♂ ♀, B.M.

Var. Alæ anticæ supra maris rufescentes, feminæ ad margines fuscæ: alæ subtus multo pallidiores ochreo-cinerascentes.

Spain.

♂ ♀, B.M.

14. *Cænonympha pamphiloides*.

Cænonympha pamphiloides, *Reakirt*, *Proc. Ent. Soc. Philad.* vi. p. 146 (footnote) (1866).

Colorado Territory.

Like *Pamphilus* on the upperside and on the underside of the front wings. Seems to differ chiefly in the longer fringe to the wings, the irregular banding of the hind wings, and more distinct ocellation of the underside. If these are the only differences, it may be considered a local variety of *Pamphilus*; for the form of the central bands and the constancy of the ocelli cannot be at all relied upon in this genus. We have *Davus* varying by slow degrees from the European *Philoxenus* to the *inornata* of the New World; and our specimens of *Geticus* vary from a silver-banded insect, with ten large distinct ocelli on the underside, to an insect having only a fine silver line and five small ocelli in the hind wings.

15. *Cænonympha Pamphilus*.

Papilio Pamphilus, *Linnaeus, Fauna Suecica*, ed. 2. p. 273. n. 1044 (1761); also of *Fabricius, Lewin, &c.*

Papilio (pleb. rural.) *Pamphilus*, *Esper, Schmett.* i. pt. 1. tab. 21. fig. 3 (1777).

Maniola Pamphilus, *Schrank, Fauna Boica*, ii. pt. 1. p. 180. n. 1512 (1801).

Satyrus Pamphilus, *Godart, Enc. Méth.* ix. p. 549. n. 179 (1819).

Hipparchia Pamphilus, *Stephens, Ill. Haust.* i. p. 69. sp. 20 (1828).

Cænonympha Pamphilus, *E. Doubleday, List Lep. Brit. Mus.* i. p. 141 (1844).

Cænonympha Pamphile, *Hübner, Verz. bek. Schmett.* p. 65. n. 633 (1816).

Satyrus Pamphile, *Godart, Léop. de France*, i. p. 176. n. 60, pl. 8 second. (1821).

Papilio Menalcas, *Poda, Ins. Mus. Græc.* p. 78. n. 50 (1761); also of *Scopoli* (1763).

Papilio Nephelē, *Hufnagel, Tab. in Berl. Mag.* ii. p. 78. n. 43 (1766); *Hübner, Eur. Schmett.* i. pl. 51. figs. 237-239 (1805).

Var. Papilio (pleb. rural.) *Pamphilus*, *Esper, Schmett.* i. pt. 2. tab. 78. cont. 28. fig. 4 (1777).

South Europe.

B.M.

Var. LYLUS.

Papilio (pleb. rural.) *Lyllus*, *Esper, Schmett.* i. pt. 2. tab. 122. cont. 77. fig. 1 (1777); also of *Oechsenheimer*.

Satyrus Lyllus, *Godart, Enc. Méth.* ix. p. 548. n. 178 (1819); *Lép. de France*, ii. p. 152. n. 58, pl. 20. figs. 9, 10 (1822).

Papilio (*Hipparchia*) *Lyllus*, *Freyer, Neuere Beiträge*, vi. p. 29. n. 898, tab. 499. fig. 1 (1852).

Cænonympha Lyllus, *Westwood in Gen. Diurn. Lepid.* p. 398. n. 17 (1851).

Satyrus Pamphilus, var., *Boisduval*, *Sp. Gén. Lép. Ind.* n. 33, sp. 277 (1840).

Cænonympha Pamphilus, var., *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 142 (1844).

Var. Papilio Pamphila, *Hübner*, *Eur. Schmett.* i. pl. 109. figs. 557, 558 (1805).

Var. Satyrus Lyllus, var., *Herrich-Schäffer*, *Eur. Schmett.* i. tab. 90. figs. 430, 431 (1843-45).

South Europe.

B.M.

16. *Cænonympha Davus* (? *Tullia*, *Müller*).

Papilio Davus, *Fabricius*, *Gen. Ins.* p. 259 (1776); *Ent. Syst.* iii. pt. 1. p. 221. n. 690 (1793); also of *Herbst*.

Satyrus Davus, *Godart*, *Lép. de France*, ii. p. 155. n. 59, pl. 21. figs. 1, 2 (1822).

Hipparchia Davus, *Stephens*, *Ill. Haust.* i. p. 67. sp. 17 (1828).

Cænonympha Davus, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 141 (1844).

Papilio Hero, *De Geer*, *Ueberf. v. Goeze*, ii. pt. 1. p. 147, tab. 2. fig. 4 (1776); but not of *Linnaeus*.

Maniola Tiphon, *Schrank*, *Fauna Boica*, ii. pt. 1. p. 179. n. 1311 (1801).

Papilio Typhon, *Haworth*, *Lep. Brit.* p. 16 (1803).

Papilio Tullia, *Hübner*, *Eur. Schmett.* i. tab. 52. figs. 243, 244 (1805).

? *Var. Papilio* (Nymph.) *Tullia*, *Müller*, *Faun. Fridr.* p. 36. n. 332 (1764).

Germany, England.

B.M.

Var. PHILOXENUS.

Papilio (pleb. rural.) *Philoxenus*, *Esper*, *Schmett.* i. pt. 2. tab. 54. cont. 4. fig. 3, tab. 78, cont. 28. fig. 3 (1777).

Cænonympha Philoxena, *Hübner*, *Verz. bek. Schmett.* p. 65. sp. 634 (1816).

Papilio Musarion, *Borkhausen*, *Naturgesch. eur. Schmett.* i. p. 92. n. 30 (1788).

Germany.

B.M.

Var. POLYMEDA.

Papilio Polymeda, *Jermyns*, *Engl. Butterf.* p. 47 (1824).

Papilio Polydama, *Haworth*, *Lep. Brit.* p. 16 (1803); *Stephens*, *Ill. Haust.* i. p. 66. sp. 16, pl. 7. fig. 3 (1828).

Merioneth.

Brit. Coll. B.M.

Var. IPHIS.

Papilio Iphis, *Borkhausen, Rhein. Magaz.* i. p. 241. n. 23 (1793).

Hipparchia Iphis, *Stephens, Ill. Haust.* i. p. 64, pl. 7. figs. 1, 2 (1828).

England, Germany.

B.M.

Var. LAIDION.

Papilio Laidion, *Borkhausen, Naturgesch. eur. Schmett.* i. p. 91. n. 29, pl. 1. figs. 5, 6 (1788).

England.

B.M.

Var. ISIS.

Papilio Isis, *Thunberg, Dissert. Acad.* iii. p. 45 (1801).

Satyrus Isis, *Zetterstedt, Ins. Lappon.* p. 905. n. 16 (1828);
Herrich-Schäffer, Eur. Schmett. i. figs. 293-296 (1843-45).

Cænonympha Isis, *Westwood in Gen. Diurn. Lepid.* p. 398. n. 14 (1851).

Papilio Davus, var., *Ochsenheimer, Schmett. v. Eur.* p. 303 (1807).

Erebia Davus, var., *Dalman, Pap. Suec.* p. 83. n. 14 (1824).

Papilio (*Hipparchia*) *Demophile*, *Freyer, Neuere Beiträge*, v. p. 97. n. 780, tab. 439. figs. 3, 4 (1845).

Lapland (Zetterstedt & Freyer), Newfoundland.

B.M.

17. *Cænonympha inornata.*

Cænonympha inornata, *Edwards, Proc. Acad. Nat. Sci. Philad.* p. 163 (1861).

Lake Winnipeg (Edwards), Vancouver's Island, West Coast.

B.M.

18. *Cænonympha California.*

Cænonympha California, *Westwood & Hewitson, Gen. Diurn. Lepid.* p. 398. n. 15, pl. 67. fig. 2 (1851).

Satyrus Californius, *Boisduval, Ann. Ent. Soc. France*, p. 309. n. 60 (1852).

California.

♂ ♀, B.M.

Var. CERES.

Cænonympha Ceres, *Butler, Ent. Month. Mag.* iii. p. 78 (1866).

California.

♀, B.M.

Scarcely sufficiently distinct to rank as a species.

19. *Cænonympha ochracea*.

Cænonympha ochracea, *Edwards, Proc. Acad. Nat. Sci. Philad.*
p. 163 (1861).

Lake Winnipeg, California, Kansas (Edwards).

The position of this species cannot be determined; it may come near *Davus*; but nothing is said in the description concerning its affinities.

20. *Cænonympha galactinus*.

Satyrus galactinus, *Boisduval, Ann. Ent. Soc. France*, p. 309.
n. 61 (1852).

California.

B.M.

21. *Cænonympha Saadi*.

Satyrus Saadi, *Kollar, Denkschrift K. Akad. Wissenschaft. Wien*,
p. 52. sp. 9 (1851).

Hipparchia (*Cænonympha*) *Iphias*, *Eversmann, Bull. Soc. Imp. Nat. Mosc.* pt. i. p. 618. n. 7 (1851).

Russian Armenia (Eversmann), Persia.

B.M.

22. *Cænonympha Sunbecca*.

Hipparchia Sunbecca, *Eversmann in Bull. Soc. Imp. Nat. Mosc.*
p. 538, tab. 7. figs. 4a, 4b (1843).

Satyrus Sunbecca, *Herrich-Schäffer, Eur. Schmett.* i. tab. 120.
figs. 611, 612 (1845).

Cænonympha Sunbecca, *Westwood in Gen. Diurn. Lepid.* p. 398.
n. 18 (1851).

Dry plains near Lake Noor Saisan (Eversmann).

Genus 8. *TRIPHYSA* *, *Zeller* (1850).

Typical Species *Triphysa* (*Tircis*) *Phryne*.

1. *Triphysa Phryne*.

Papilio (*Nymph. Gemm.*) *Phryne*, *Pallas, Reise*, pt. i. n. 60
(1776).

Papilio (*pleb. rural.*) *Phryne*, *Esper, Schmett.* i. pt. 2. tab. 89.
cont. 39. figs. 3, 4; var. ?, tab. 113. cont. 68. fig. 6 (1777).

Papilio Phryne, *Borkhausen, Naturgesch. eur. Schmett.* i. pp. 108
& 246. n. 52 (1788); *Hübner, Eur. Schmett.* i. pl. 44. figs.
200, 201, pl. 140. figs. 708, 709 (1805).

Cænonympha Phryne, *E. Doubleday, List Lep. Brit. Mus.* i.
p. 142 (1844); also of *Westwood* (1851).

* The genus *Phryne* of Herrich-Schäffer was rejected by Zeller as having been previously used for a genus of Reptilia.

Satyrus Phryne, *Herrich-Schäffer*, *Eur. Schmett.* i. tab. 22. figs. 106, 107 (1845).

Papilio Phryneus, *Fabricius*, *Mant. Ins.* ii. p. 33. n. 354 (1787).

Satyrus Phryneus, *Godart*, *Enc. Méth.* ix. p. 529. n. 119 (1819); also of *Boisduval & Duponchel* (1832).

Papilio Tircis, *Cramer*, *Pap. Exot.* iv. pl. 373. figs. C, D (1782); also of *Herbst* (1796).

Phryne Tircis, *Herrich-Schäffer*, *Syst. Bearb.* p. 90 (1843).

Triphysa Tircis, *Zeller*, *Ent. Zeit.* p. 311 (1850).

Eastern Russia.

♂ ♀, B.M.

2. *Triphysa Dohrnii*.

Triphysa Dohrnii, *Zeller*, *Ent. Zeit.* p. 308 (1850); *Herrich-Schäffer*, *Eur. Schmett.* i. tab. 133. figs. 641, 642 (1843-56).

Southern Russia.

Genus 9. AULOCERA, *Butler* (1867).

Typical Species *Aulocera Brahminus*.

1. *Aulocera Brahminus*.

Satyrus Brahminus, *Blanchard* in *Jacquemont's Voyage in India*, pl. 2. figs. 4-6 (1844); *Westwood* (1851).

Aulocera Brahminus, *Butler*, *Ent. Month. Mag.* iv. p. 121. n. 1 (1867), (neuraton) fig. 1.

Satyrus Swaha, *Kollar* in *Hügel's Kaschmir*, p. 444, tab. 14. figs. 1, 2 (1844).

Satyrus Saraswati (*synonym*), *Moore*, *Cat. Lep. Mus. East-Ind. Comp.* i. p. 229. n. 483 (1857).

North India, Nepaul, Himalayas, Cashmere.

B.M.

2. *Aulocera Saraswati*.

Satyrus Saraswati, *Kollar* in *Hügel's Kaschmir*, p. 445, tab. 14. figs. 3, 4 (1844); *Westwood* (1851).

Aulocera Saraswati, *Butler*, *Ent. Month. Mag.* iv. p. 121. n. 2 (1867).

North India.

B.M.

3. *Aulocera Padma*.

Satyrus Padma, *Kollar* in *Hügel's Kaschmir*, p. 445, tab. 15. figs. 1, 2 (1844); *Westwood* (1851).

Aulocera Padma, *Butler*, *Ent. Month. Mag.* iv. p. 122. n. 3 (1867).

North India.

B.M.

4. *Aulocera Avatara*.

Satyrus Avatara, *F. Moore, Cat. Lep. Mus. East-Ind. Comp. i.*
p. 229. n. 481 (1857).

Aulocera Avatara, *Butler, Ent. Month. Mag. iv. p. 122. n. 4*
(1867).

North India (Moore).

5. *Aulocera Scylla*.

Aulocera Scylla, *Butler, Ent. Month. Mag. iv. p. 122. n. 5* (1867).

Near Sylhet.

B.M.

Genus 10. *HIPPARCHIA*, *Fabricius* (1808).

Typical Species *Hipparchia Fagi* (*Hermione*).

1. *Hipparchia Proserpina*.

Papilio Proserpina, *Denis, Wien. Verz. p. 169. fam. F. n. 23*
(1775); *Hübner, Eur. Schmett. i. pl. 26. figs. 119-121* (1805).

Papilio (Nymph. Gemm.) *Proserpina*, *Esper, Schmett. v. Eur. i.*
pt. 1. tab. 39. *Suppl. xv. fig. 2, tab. 42. Suppl. xviii. fig. 3*
(1777).

Satyrus Proserpina, *Westwood in Gen. Diurn. Lepid. p. 390. n. 12*
(1851).

Papilio Circe, *Fabricius, Ent. Syst. iii. pt. 2. p. 233. n. 728* (1793).

Satyrus Circe, *Godart, Enc. Méth. ix. p. 513. n. 108* (1819);

Lép. de France, i. p. 131. n. 41, pl. 7 secund. fig. 1 (1821).

Hipparchia Circe, *E. Doubleday, List Lep. Brit. Mus. pt. 1. p. 131*
(1844).

Germany, Corfu.

B.M.

2. *Hipparchia Fagi*.

Papilio Fagi, *Scopoli, Ent. Carn. p. 152. n. 428, fig. 428* (1763).

Papilio Hermione, *Linnaeus, Syst. Nat. ii. p. 773. n. 149* (1766);

Hübner, Eur. Schmett. i. pl. 27. figs. 122-124 (1805).

Satyrus Hermione, *Godart, Enc. Méth. ix. p. 515. n. 110* (1819);

Lép. de France, i. p. 137. n. 43, pl. 7 secund. fig. 2 (1821).

Papilio (Nymph. Gemm.) *Hermione major*, *Esper, Schmett. i.*
pt. 1. tab. 8. fig. 3, pt. 2. tab. 70. cont. 20. fig. 4 (1777).

Germany, South Europe.

B.M.

Local form. Athens.

B.M.

3. *Hipparchia Alcyone*.

Papilio Alcyone, *Denis, Wien. Verz. p. 169. fam. F. n. 21* (1775);

Hübner, Eur. Schmett. i. pl. 27. figs. 125, 126 (1805).

Satyrus Alcyone, *Boisduval, Icon. Hist. Lép. p. 202. n. 2, pl. 40.*
figs. 5, 6 (var.) (1832).

Papilio (Nymph. Gemm.) *Hermione minor*, *Esper*, *Schmett.* i. pt. 1. tab. 8. fig. 2 (1777).

Papilio Hermione ♂, *Herbst*, *Natursyst. Schmett.* viii. tab. 215. figs. 1, 2 (1796).

Satyrus Hermione, var., *Boisduval*, *Sp. Gén. Ind. Meth.* p. 30. n. 241 (1840).

Germany.

B.M.

It is not to be credited that this should be only a variety of *Fagi* (*Hermione*); the form of the central band is utterly unlike that species.

4. *Hipparchia Marmoræ*.

Papilio Marmoræ, *Hübner*, *Eur. Schmett.* i. pl. 164. figs. 814–817 (1805).

Satyrus Neomiris, *Godart*, *Enc. Méth. ix. Suppl.* p. 827. n. 111 (1819); *Lép. de France*, ii. p. 88. n. 28, pl. 40. figs. 1, 2 (1822).

Satyrus Neomyris (sic), *Boisduval*, *Icon. Hist. Lép.* pl. 42. figs. 6–8 (1832); and also of *Westwood* (1851).

Hipparchia Neomyris, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 132 (1844).

Papilio (*Satyrus*) *Iolaus*, *Bonelli in Mem. Accad. Torino*, xxx. p. 179, sp. 4a, tav. 3. figs. 1 ♂, 1 ♀ (1826).

Corsica.

B.M.

5. *Hipparchia Anthelea*.

♂. *Papilio Anthelea*, *Hübner*, *Eur. Schmett.* i. pl. 174. figs. 861, 862 (1805).

Papilio (*Hipparchia*) *Anthelea*, *Freyer*, *Neuere Beiträge*, iii. p. 101. n. 436, tab. 265. fig. 2. (1839).

♀. *Satyrus Anthelea*, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 303, 304 (1843–45); ♂. *Boisduval & Duponchel* (1832).

♂ (part.). *Hipparchia Anthelea*, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 131 (1844).

Crete.

♂, B.M.

Var. PONTICA.

Papilio (*Hipparchia*) *Pontica*, *Freyer*, *Neuere Beiträge*, v. p. 158. n. 861, pl. 475. figs. 2, 3 (1845).

Satyrus Pontica, *Westwood in Gen. Diurn. Lepid.* p. 390. n. 22 (1851).

Turkey, Crete, Smyrna.

♂ ♀, B.M.

6. *Hipparchia Thelephassa*.

♂ ♀. *Eumenis Thelephassa*, *Hübner*, *Exot. Schmett.* ii. pl. 85. figs. 1–4 (1806).

♂. *Satyrus Thelephassa*, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 305, 306, (♀) 178, 179 (1843-45); ♀, *Duponchel*, *Lép. de France*, i. p. 184. n. 42, pl. 28. figs. 1, 2 (1832).

♀. *Papilio* (*Hipparchia*) *Anthelea*, *Freyer*, *Neuere Beiträge*, iii. tab. 265. fig. 3 (1839).

♀. *Satyrus Anthelea*, *Boisduval*, *Icon. Hist. Lép.* pl. 41. figs. 3, 4 (1832); also of *Hewitson* (1851).

Local form, ♂ ♀. *Hipparchia Thelephassa*, *Klug & Ehrenberg*, *Symb. Phys. Ins.* tab. 29. figs. 1-4 (1829-45).

Syria (Klug), Persia, Russia.

♂ ♀, B.M.

Though allied to *Anthelea*, this species must surely be distinct; *Pontica*, however, is only a slight variety, although Westwood has retained it as a species.

7. *Hipparchia Pelopea*.

Hipparchia Pelopea, *Klug in Ehrenb. & Hempr. Symb. Phys.* tab. 29. figs. 5-8 (1829-45).

Satyrus Pelopea, *Westwood in Gen. Diurn. Lepid.* p. 391. n. 29 (1851).

Syria, Persia.

B.M.

8. *Hipparchia Hippolyte*.

Papilio (Nymph. Gemm.) *Hippolyte*, *Esper*, *Schmett.* i. pt. 2. p. 164 (1777).

Papilio Hippolyte, *Zetterstedt, Ins. Lappon.* p. 901. n. 3 (1840).

Satyrus Hippolyte, *Boisduval, Icon. Hist. Lép.* p. 210. n. 5, pl. 42. figs. 1, 2 (1832).

Hipparchia Hippolyte, *E. Doubleday, List Lep. Brit. Mus.* i. p. 132 (1844).

Papilio Hippolyte (*sic*), *Herbst, Natursyst. Schmett.* viii. p. 200. n. 118, tab. 211. figs. 3, 4 (1796).

Papilio (*Hipparchia*) *Hippolyte*, *Freyer, Neuere Beiträge*, iii. tab. 278. figs. 1, 2 (*Hippolyte*), p. 119. n. 455 (1839).

Papilio (Nymph. Gemm.) *Agave*, *Esper, Schmett.* i. pt. 2. tab. 84. cont. 34. fig. 4, ♂ (1777).

Papilio Agave, *Borkhausen, Naturgesch. Schmett.* i. p. 102. n. 43 (1788); *Hübner, Eur. Schmett.* i. pl. 30. figs. 139, 140 (1805).

Papilio Alcyone, *Fabricius, Ent. Syst.* iii. pt. 1. p. 231. n. 724 (1793).

Satyrus Alcyone, *Godart, Enc. Méth.* ix. p. 517. n. 112 (1819).
Russia.

B.M.

9. *Hipparchia Hübneri*.

Satyrus Hübneri, *Felder, Reise der Fregatte Novara*, iii. tab. 69. figs. 8, 9 (published 1867).

Eastern Himalayas (Felder).

Closely allied to *H. Hippolyte*, Esper; according to Felder most nearly allied to *H. Agave*, Esper.

10. **Hipparchia Beroë.**

Satyrus Beroë, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 108–111 (1843).

Papilio (*Hipparchia*) *Beroë*, *Freyer*, *Neuere Beiträge*, v. p. 53. n. 724, tab. 415. figs. 1, 2 (1845).

Var. ♀. *Satyrus Mamurra*, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 314, 315 (1843–45).

Turkey, Crete.

B.M.

11. **Hipparchia Geyeri.**

Satyrus Geyeri, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 301, 302 (1843–45).

Ararat.

12. **Hipparchia Autonoe.**

Papilio (*Nymph. Gemm.*) *Autonoe*, *Esper*, *Schmett.* i. pt. 2. tab. 86. cont. 36. figs. 1–3 (1777).

Papilio Autonoe, *Borkhausen*, *Naturgesch. Schmett.* i. p. 103. n. 45 (1788).

Satyrus Autonoe, *Godart*, *Enc. Méth.* ix. p. 518. n. 114 (1819).

Papilio (*Hipparchia*) *Autonoe*, *Freyer*, *Neuere Beiträge*, iii. p. 127. n. 465, tab. 283. figs. 1, 2 (1839).

Hipparchia Autonoe, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 132 (1844).

Papilio Autonoe (*sic*), *Herbst*, *Natursyst. Schmett.* viii. p. 205. n. 120 (1796).

Var. ♀? *Papilio Autonoe*, *Hübner*, *Eur. Schmett.* i. pl. 30. figs. 137, 138 (1805).

Russia.

B.M.

13. **Hipparchia Arethusa.**

Papilio Arethusa, *Denis*, *Wien. Verz.* p. 169. n. 16 (1775); *Hübner*, *Eur. Schmett.* i. pl. 190. figs. 937, 938 (1805).

Papilio (*Nymph. Gemm.*) *Arethusa*, *Esper*, *Schmett.* i. pt. 2. tab. 69. cont. 19. figs. 3, 4 (1777).

Satyrus Arethusa, *Godart*, *Enc. Méth.* ix. p. 517. n. 113 (1819); *Lép. de France*, i. p. 141. n. 45, pl. 7 tert. fig. 2 (1821).

Hipparchia Arethusa, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 132 (1844).

Papilio Arethusus, *Herbst*, *Natursyst. Schmett.* viii. p. 166, tab. 206. figs. 3–6 (1796).

Russia, Germany, Odessa.

B.M.

Var. ERYTHIA.

♂. *Papilio Erythia*, *Hübner, Eur. Schmett. i. pl. 115. figs. 591, 592 (1805).*

♀ *Var. ? Papilio Arethusa, ibid. pl. 34. figs. 154, 155.*

Satyrus Arethusa, var. Boabdil, Herrich-Schäffer, Eur. Schmett. i. figs. 474-476 (1843-45).

Spain.

B.M.

Var. BOABDIL.

Satyrus Boabdil, Rambur, Faun. Ent. de l'Andal. pl. 12. figs. 1, 2 (1842).

Papilio (Hipparchia) Arethusa, Freyer, Neuere Beiträge, vi. p. 134. n. 1032, tab. 567. fig. 3 (1852).

Spain (Rambur), Bordeaux.

B.M.

14. *Hipparchia Heidenreichii.*

Satyrus Heidenreichii, Lederer, Verhandl. zool.-botan. Gesellsch. Wien, iii. p. 359, pl. 1. fig. 2 (1853).

Altai Mountains.

B.M.

15. *Hipparchia Briseis.*

Papilio Briseis, Linnæus, Syst. Nat. ii. p. 770. n. 139 (1766); Hübner, Eur. Schmett. i. pl. 28. figs. 130, 131 (1805).

Papilio (Hipparchia) Briseis, Freyer, Neuere Beiträge, vi. p. 3. n. 871, tab. 481 (1852).

Satyrus Briseis, Godart, Enc. Méth. ix. p. 514. n. 109 (1819); Léop. de France, i. p. 134. n. 42, pl. 7. fig. 1 (1821); also of Herrich-Schäffer.

Hipparchia Briseis, Westwood & Humphreys, Brit. Butterf. p. 68, pl. 19. figs. 1, 2 (1840).

Papilio Ianthé major, Esper, Schmett. i. pt. 1. tab. 26. Suppl. ii. fig. 1; minor, ibid. fig. 2 (1777).

Maniola Ianthé, Schrank, Faun. Boic. ii. pt. 1. p. 182. n. 1316 (1801).

Papilio Dædale, Bergsträsser, Nomenclat. Inst. pt. 4. tab. 94. figs. 1, 2 (1780); also of Borkhausen (1788).

Germany, South Europe.

B.M.

Var. ♀. Papilio (Nymph. Gemm.) pirata, Esper, Schmett. i. pt. 2. tab. C. cont. 55. fig. 3 (1777).

Papilio pirata, Hübner, Eur. Schmett. i. pl. 118. figs. 604, 605 (1805).

South Europe.

B.M.

Local form, ♂. Fascia anticarum latiore et minus interrupta, margine postico posticarum albicante, maculis subtus minoribus.

Persia.

B.M.

Var. PRIEURI.

Satyrus Prieuri, *Pierret in Ann. Soc. Ent. France*, vi. p. 304. pl. 12. fig. 6 (1837); also of *Westwood* (1851).

Barbary.

Only a local form of *Briseis*.

16. *Hipparchia Persephone.*

Papilio Persephone, *Hübner, Eur. Schmett.* i. pl. 115. figs. 589, 590, and pl. 140. figs. 710, 711 (1805).

Papilio anthe, *Ochsenheimer, Schmett. v. Eur.* i. p. 169. n. 2 (1807).

Satyrus anthe, *Godart, Enc. Méth.* ix. p. 826. nos. 109, 110 (1819); also of *Duponchel* and *Boisduval* (1832).

Russia.

B.M.

♀. *Local form.* *Anticæ* subtus ferruginosæ, maculis nigris minoribus; *posticæ* fascia media albicante, area basali magis fuscescente.

Syria (Mount Hermon).

B.M.

Satyrus anthe, *var. Caucasica*, *Hanifa, Nordmann, in Bull. Soc. Imp. Nat. Mosc.* p. 406, tab. 9. figs. 1-3 (1851).

Local form. *Fasciis* supra ferrugineis, maculis nigris minoribus; *anticarum* subtus magis ferrugineis; *fascia* media *posticarum* minus albicante; aliter velut præcedens.

Persia.

♀, B.M.

This form is very like *Pelopea* on the upperside, but without the white intermacular spots.

Ochsenheimer has substituted the name *anthe* for *Persephone* on the ground of this name having been previously used by *Fabricius*; Such a name, however, does not seem to exist amongst the *Satyridae*. *Persiphone* (sic), *Ent. Syst.* iii. pt. 1. p. 542, seems to represent an *Acræa*, not a *Satyrus*.

17. *Hipparchia Bischoffii.*

Satyrus Bischoffii, *Herrich-Schäffer, Eur. Schmett.* i. figs. 307-310 (1843-45); also of *Hewitson* (1851).

Satyrus Bischoffii (sic), *Schaum, Bericht.* p. 231 (1851).

Russia.

B.M.

18. *Hipparchia Semele.*

Papilio Semele, *Linnaeus, Syst. Nat.* i. p. 474 (1758), ii. p. 772. n. 148 (1766); *Hübner, Eur. Schmett.* i. pl. 31. figs. 143, 144, pl. 167. figs. 826, 827 (1805); also of *Herbst* (1796).

Papilio (Nymph. Gemm.) *Semele*, *Esper, Schmett.* i. pt. 1. tab. 8. fig. 1 (var. ♀?) (1777).

Satyrus Semele, *Godart, Enc. Méth.* ix. p. 516. n. 111 (1819).

Hipparchia Semele, *Stephens, Ill. Haust.* i. p. 56. sp. 4 (1828).

Papilio Danaë, *Hufnagel, Berl. Mag.* ii. p. 82. n. 50 (1766).

Var. Satyrus Semele, *Herrich-Schäffer, Eur. Schmett.* i. figs. 182 (1843-1845).

England, Germany, South Europe. B.M.

Var. ♂. *Alis supra nigrescentibus, fasciis omnibus obsoletis.*

Europe. B.M.

Var. ♂. *Coloribus consuetis, forma autem Aristæi; alis anticis brevioribus.*

Jersey. B.M.

Var. ARISTÆUS.

Papilio (Satyrus) Aristæus, *Bonelli in Mem. R. Accad. Torino*, xxx. p. 177. sp. 3, tav. 2. figs. 1♂, 1♀ (1826); also of *Herrich-Schäffer*.

Satyrus Aristæus, *Westwood in Gen. Diurn. Lepid.* p. 390. n. 25 (1851).

Papilio Semele, *Hübner, Eur. Schmett.* i. pl. 168. figs. 832-835 (1805).

Satyrus Semele, *var. ?*, *Boisduval, Sp. Gén. Ind.* p. 31. n. 247 (1836).

Var. Papilio (Hipparchia) Aristæus, *Freyer, Neuere Beiträge*, v. p. 25. n. 690, tab. 397. fig. 1 (1845).

Corsica. B.M.

19. *Hipparchia Statilinus*.

Papilio Statilinus, *Hufnagel, Tab. in Berl. Mag.* ii. p. 84. n. 52 (1766); also of *Herbst and Ochsenheimer*.

Papilio (Hipparchia) Statilinus, *Freyer, Neuere Beiträge*, vi. p. 30. n. 899, tab. 499. figs. 2, 3 (1852).

Satyrus Statilinus, *Herrich-Schäffer, Eur. Schmett.* i. figs. 177; *Martianii* ♀, 190, 191 (1843-45).

Papilio Arachne, *Denis, Wien. Verz.* p. 169. fam. F. n. 18 (1775).

Papilio (Nymph. Gemm.) Arachne, *Esper, Schmett.* i. pt. 2. tab. 95. cont. 50. figs. 3, 4.

Papilio (Nymph. Gemm.) Fauna, *ibid.* pt. 1. tab. 29. *Suppl.* v. fig. 1, and pt. 2. tab. 63. cont. 13. fig. 7 (1777).

Papilio Fauna, *Sulzer, Gesch. Ins.* pt. 2. p. 145, tab. 17. figs. 8, 9 (1776); also of *Hübner* (1805).

Satyrus Fauna, *Godart, Enc. Méth.* ix. p. 520. n. 121 (1819); *Lép. de France*, i. p. 143. n. 46, pl. 7 tert. fig. 3 (1821); also of *Westwood* (1851).

Hipparchia Fauna, *E. Doubleday, List Lepid. Brit. Mus.* i. p. 130 (1844).

Papilio (Nymph. Gemm.) *Fidia*, *Esper*, *Schmett.* i. pt. 2. tab. 52. cont. 2. fig. 4 (1777).

Papilio Fidia, *Herbst*, *Natursyst. Schmett.* viii. tab. 200. figs. 1, 2 (1796).

France.

♂ ♀, B.M.

Var. ALLIONIA.

Papilio (Nymph. Gemm.) *Allionia*, *Esper*, *Schmett.* i. pt. 2. tab. 105. cont. 60. fig. 4 (1777).

Papilio Allionia, *Cyrrillus*, *Ent. Neap.* sp. 1. tab. 2. fig. 13 (1787-92); also of *Fabricius* and *Ochsenheimer*.

? *Papilio Allionius*, *Herbst*, *Natursyst. Schmett.* viii. p. 130. n. 71, tab. 201. figs. 5, 6 (1796).

? *Papilio Actæa*, *De Villers*, *Ent. Linn.* tab. iv. fig. 9 (1789).

South Europe, Naples.

♂, B.M.

This is only a variety of the true *Statilinus*; it is slightly larger, and with the central band below more distinctly margined with white.

Var. Subtus fascia albida magis albicante, ocellis anticarum supra obsoletis.

Athens.

♂, B.M.

20. *Hipparchia fatua*.

♂ ♀. *Papilio* (*Hipparchia*) *fatua*, *Freyer*, *Neuere Beiträge*, v. p. 54. n. 725, tab. 415. figs. 3, 4 (1845).

♀. *Satyrus fatua*, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 192, 193 (1845).

Papilio Allionii, *Hübner*, *Eur. Schmett.* i. pl. 165. figs. 818, 819 (1805); but not *Allionia* of *Esper*.

Athens.

♂ ♀, B.M.

21. *Hipparchia Pisidice*.

Hipparchia Pisidice, *Klug in Ehrenb. & Hempr. Symb. Phys.* t. 29. figs. 9, 10 (1829-45).

Satyrus Pisidice, *Westwood in Gen. Diurn. Lepid.* p. 391. n. 30 (1851).

Mount Sinai.

Seems to be closely allied to *Statilinus* (var. *Allionia*), but wants the second ocellus on the underside of the front wings, and differs in other particulars.

22. *Hipparchia Fidia*.

Papilio Fidia, *Linnaeus*, *Syst. Nat.* 12th ed. ii. pt. 1. p. 770. n. 138 (1766); *Hübner*, *Eur. Schmett.* i. pl. 32. figs. 147, 148 (1805);

also of *Fabricius*, *Ochsenheimer*, *Borkhausen*, &c.

Papilio (Nymph. Gemm.) *Fidia*, *Esper*, *Schmett.* i. pt. 1. tab. 49. *Suppl.* xxv. fig. 3 (1777).

Satyrus Fidia, *Godart*, *Enc. Méth.* ix. p. 520. n. 120 (1819); *Lép. de France*, ii. p. 90. n. 29, pl. 11. figs. 3, 4 (1822); also of *Westwood*.

Hipparchia Fidia, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 130 (1844).

Papilio Lusitanica, *Petiver*, *Gazoph.* i. tab. 7. fig. 5. n. 147 (1709).

Germany, France.

B.M.

Var. *Alæ multo majores, maculis supra distinctioribus.*

Gibraltar.

B.M.

23. *Hipparchia Parisatis*.

Satyrus Parisatis, *Kollar*, *Denkschrift K. Akad. Wissenschaft. Wien*, p. 52. sp. 7 (1850).

Satyrus macrophthalmus, *Eversmann* in *Bull. Soc. Imp. Nat. Mosc.* p. 615 (1851).

Russian Armenia (Eversm.), Persia.

B.M.

24. *Hipparchia Chiliensis*.

Satyrus Chiliensis, *Guérin*, *Voyage Coquille, Zool. Ins.* p. 280; *Atlas, Ins.* n. 16. figs. 4, 5 (1832).

Erebia Chiliensis, *Westwood* in *Gen. Diurn. Lepid.* p. 380. n. 53 (1851).

Erebia Chilensis (*sic*), *E. Doubleday*, *List Lep. Brit. Mus.* pt. 1. p. 127 (1844); also of *Gay* (1852).

Chili.

B.M.

25. *Hipparchia* ? *Flora*.

Satyrus Flora, *Philippi*, *Linnaea Entomologica*, xiv. p. 267. n. 4 (1860).

Chili (near Corral Valdivia).

Herr Philippi does not state the affinities of this species; but, judging from his description of the coloration of the underside, it seems to come near *Chiliensis*. Felder, however, seems to look upon it as an *Epinephele*.

26. *Hipparchia* ? *Boisduvalii*.

Erebia Boisduvalii, *Blanchard* in *Gay's Fauna Chilena*, vii. p. 32. n. 2 (1852).

Chili.

Seems to be allied to *Chiliensis*, but has the subapical ocellus of the front wings bipupillate on the underside.

27. *Hipparchia* † *Ridingsii*.

Satyrus Ridingsii, *Edwards*, *Proc. Ent. Soc. Philad.* iv. p. 201 (1866); *Reakirt*, vi. p. 145 (1866).

Colorado Territory.

Seems, from the description, to be allied to *Chiliensis*; but as no notice is taken of its affinities, it is impossible to determine its true position. It may be a species of *Eneis*.

Genus HIPPARCHIA, *Fabricius*.

Subgenus 1. COSMOSATYRUS *, *Felder* (1867).

Typical Species *Cosmosatyrus leptoneuroides*.

1. *Cosmosatyrus leptoneuroides*.

Cosmosatyrus leptoneuroides, *Felder*, *Reise der Fregatte Novara*, iii. p. 495. n. 857 (1867).

Chili (*Felder*).

Subgenus 2. SATYRUS, *Westwood* (nec *Godart*)†.

Typical Species *Satyrus Actæa*.

1. *Satyrus Actæa*.

Papilio (Nymph. Gemm.) *Actæa*, *Esper*, *Eur. Schmett.* i. pt. 2. tab. 57. cont. 7. figs. 1a, 1b (1777).

Papilio Actæa, *Borkhausen*, *Naturgesch. eur. Schmett.* i. p. 98. n. 38 (1788); *Hübner*, *Eur. Schmett.* i. pl. 33. figs. 151, 152, pl. 119. figs. 610, 611 (1805).

Satyrus Actæa, *Godart*, *Enc. Méth.* ix. p. 522. n. 124 (1819); *Lép. de France*, i. p. 145. n. 47, pl. 7 quart. fig. 1 (1821).

♀. *Papilio Actæus*, *Herbst*, *Natursyst. Schmett.* viii. tab. 201. figs. 1, 2 (1796).

Germany.

♂ ♀, B.M.

Var.? *Anticæ supra ocello subanali magno nigro*.

Germany?

♀, B.M.

Var.? *Papilio Actæus*, *Herbst*, *Natursyst. Schmett.* viii. p. 123, tab. 200. figs. 7, 8 (♂), but not tab. 201. figs. 1, 2 (1796).

Spain.

♂ ♀, B.M.

* Nearly allied to the *Chiliensis* group, but differing in its longer palpi and shorter upper discocellular in the front wing.

† The type of *Godart's Satyrus* was *Hiptio Constantia* of *Cramer*.

Satyrus. Alis apice anticarum minus acuto, antennis subintegris, lineis mediis subtus undulatis.

Var. PODARCE.

Satyrus Podarce, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 49-51 (1843-45).

Spain.

♂, B.M.

We have only a single specimen of this insect, unless the preceding variety is another form of it. It seems scarcely to be a distinct species.

2. *Satyrus Pimpla.*

Satyrus Pimpla, *Felder*, *Reise der Fregatte Novara*, iii. tab. 69. figs. 10, 11 (1867).

Eastern Himalayas (Felder).

Most nearly allied to *Actæa*, *Esper*.

3. *Satyrus Cordula* (? *S. Ferula*).

Papilio Cordula, *Fabricius*, *Ent. Syst.* iii. pt. 1. p. 226. n. 708 (1793); var. ♀, *Hübner*, *Eur. Schmett.* i. pl. 121. figs. 619, 620 (1805).

♀. *Satyrus Cordula*, *Godart*, *Enc. Méth.* ix. p. 522. n. 123 (1819); *Lép. de France*, ii. p. 94. n. 31, pl. 12. figs. 3, 4 (1822) as ♂.

Hipparchia Cordula, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 129 (1844).

Italy.

♂ ♀, B.M.

? *Var. Papilio Ferula*, *Fabricius*, *op. cit.* p. 225. n. 707 (1793).

Italy.

Var. BRYCE.

Papilio Bryce, *Hübner*, *Eur. Schmett.* i. pl. 144. figs. 724-727 (1805); also of *Ochsenheimer* (1807).

Satyrus Bryce, *Godart*, *Enc. Méth.* ix. p. 522. n. 123 (1819); *Lép. de France*, ii. p. 92. n. 30, pl. 12. figs. 1, 2 (1822); also of *Westwood* (1851).

Papilio Hippolytus, *Herbst*, *Natursyst. Schmett.* viii. p. 126. n. 67, tab. 201. figs. 3, 4 (1796).

♀. *Var. ? Papilio Hippodice*, *Hübner*, *Eur. Schmett.* i. pl. 142. figs. 718, 719 (1805).

Russia.

♂ ♀, B.M.

A local race of *S. Cordula* of *Fabricius*.

Hippolytus of *Herbst* will remain as a synonym of this race, as the name *Hippolyte* has already been used in *Hipparchia* proper.

Var. Pæas.

Papilio (Nymph. Gemm.) *Pæas*, *Esper*, *Schmett.* i. pt. 2. tab. 112. cont. 67. fig. 1 (1777).

Papilio *Pæas*, *Hübner*, *Eur. Schmett.* i. pl. 29. figs. 132, 133 (1805).

Papilio *Actæa*, *Fabricius*, *Ent. Syst.* iii. pt. 1. p. 225. n. 705 (1793).

Var.? *Papilio* *Proserpina*, *Cyrill.* *Ent. Neap.* i. tab. 2. fig. 11 (1787).

Papilio *Cyrillus*, *Herbst*, *Natursyst. Schmett.* viii. p. 165. n. 99, tab. 206. figs. 1, 2 (1796).

Russia.

♂ ♀, B.M.

4. *Satyrus Glaucopæ.*

Epinephelæ *Glaucopæ*, *Felder*, *Reise der Fregatte Novara*, iii. tab. 67. figs. 5, 6 (1867).

Brazil (Felder).

Allied to *Bryce*. It does not seem to be allied to *Coctei*, as *Felder* states in his description.

5. *Satyrus Abd-el-Kader.*

♂. *Satyrus* *Abd-el-Kader*, *Pierret* in *Ann. Soc. Ent. de France*, vi. p. 19, pl. 1. figs. 5, 6 (1837).

♀. *Lucas*, *Expl. Algér. Léop.* pl. 2. figs. 3, 3a (1849).

Algeria.

Allied to *S. Bryce*.

6. *Satyrus Virbius.*

Papilio (*Hipparchia*) *Virbius*, *Freyer*, *Neuere Beiträge*, v. p. 135. n. 828, tab. 463. figs. 1, 2 (1845).

Satyrus *Virbius*, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 45–48 (1843–45).

South Russia.

Intermediate between *Bryce* and *Phædra*.

Section *MINOIS*, *Hübner*.

Typical Species *Minois* (*Phædra*) *Dryas*.

7. *Satyrus Dryas.*

Papilio *Dryas*, *Scopoli*, *Ent. Carn.* p. 153. n. 429, fig. 429 (1763).

Papilio (Nymph. Gemm.) *Dryas*, *Esper*, *Schmett.* i. pt. 1. tab. 40.

Suppl. lxvi. figs. 1, 2 (1777).

Papilio *Phædra*, *Linnaeus*, *Syst. Nat.* ii. p. 773. n. 150 (1766);

Hübner, Eur. Schmett. i. pl. 28. figs. 127-129 (1805); also of *Fabricius, Ochsenheimer, Borkhausen, and Herbst.*

Maniola Phædra, Schrank, Faun. Boic. p. 181 (1801).

Hipparchia Phædra, Stephens, Ill. Haust. p. 56. sp. 5 (1828); also of *Wood.*

Var. ♀. Satyrus Phædra, Godart, Enc. Méth. ix. p. 523. n. 126 (1819); *Lép. de France*, i. p. 147. n. 48, pl. 7 quart. fig. 2 (1821); also of *Westwood* (1851).

Papilio (Nymph. Gemm.) Briseis, Esper, Schmett. i. pt. 1. tab. 6. fig. 1 (1777).

Papilio Athene, Borkhausen, Naturgesch. eur. Schmett. i. p. 71. n. 8 (1788).

France, Switzerland, Germany. B.M.

Var. Alis multo majoribus, ocellis permagnis.

Japan. B.M.

Var. Magnitudine consueta, sed alis magis fuscescentibus.

Hakodadi and Northern India. B.M.

Var. Major et paulum pallidior, ocellis magnis.

North China. B.M.

8. *Satyrus Alope.*

Papilio Alope, Fabricius, Ent. Syst. iii. pt. 1. p. 229. n. 715 (1793).

Satyrus Alope, Godart, Enc. Méth. ix. p. 524. n. 127 (1819); *Boisduval & Leconte, Icon. Lép. Am. Sept.* p. 228, pl. 59 (1833).

Enodia? Alope, E. Doubleday, List Lep. Brit. Mus. i. p. 136 (1844).

Satyrus (Enodia) Alope, Westwood in Gen. Diurn. Lepid. p. 392. n. 45 (1851).

Enodia Alope, Reakirt, Proc. Ent. Soc. Philad. vi. p. 145 (1866).

United States. B.M.

9. *Satyrus Pegala.*

Papilio Pegala, Fabricius, Ent. Syst. iii. pt. 1. p. 230. n. 720 (1793).

Satyrus Pegala, Godart, Enc. Méth. ix. p. 524. n. 128 (1819).

Satyrus (Enodia) Pegala, Westwood in Gen. Diurn. Lepid. p. 392. n. 46 (1851).

North America.

This species appears from the description to be closely allied to *Alope*; it, however, has only one ocellus in the front wings.

10. *Satyrus Ariane*.

Satyrus Ariane, *Boisduval*, *Ann. Soc. Ent. France*, p. 307. n. 58 (1852).

Enodia Ariane, *Reakirt*, *Proc. Ent. Soc. Philad.* vi. p. 145 (1866).

California.

♂ ♀, B.M.

11. *Satyrus Nephela*.

Hipparchia Nephela, *Kirby in Fauna Boreali-Amer.* p. 297. n. 415 (1837).

Erebia Nephela, *Westwood in Gen. Diurn. Lepid.* p. 380. n. 42 (1851).

Canada (Kirby), Canada West.

♂ ♀, B.M.

Kirby's description of this species is not good; it applies best to the female. The preceding species is probably a local form of this insect.

12. *Satyrus Cænonympha*.

Epinephele Cænonympha, *Felder, Reise der Fregatte Novara*, iii. tab. 69. figs. 14, 15 (1867).

Eastern Himalayas, Ladak, Suroo (Felder).

Allied to *Sthenele*, *Boisd.*

13. *Satyrus Sthenele*.

Satyrus Sthenele, *Boisduval*, *Ann. Soc. Ent. France*, p. 308. n. 60 (1852).

Satyrus silvestris, *Edwards*, *Proc. Acad. Nat. Sci. Philad.* p. 162 (1861).

California.

B.M.

14. *Satyrus Hyperantus*.

Papilio Hyperantus, *Linnaeus*, *Fauna Suec.* p. 273. n. 1041 (1761); *Syst. Nat.* i. p. 471 (1758), ii. p. 768. n. 127 (1766).

Papilio Hyperanthus (sic), *Füssli*, *Schweiz. Ins.* p. 29. n. 556 (1775).

Papilio (Nymph. Gemm.) *Hyperanthus* (sic), *Esper*, *Schmett.* i. pt. 1. tab. 5. fig. 1 (1777).

Satyrus Hyperanthus, *Godart*, *Enc. Méth.* ix. p. 538. n. 162 (1819).

Hipparchia Hyperanthus, *Stephens*, *Ill. Haust.* i. p. 60. sp. 10 (1828).

Enodia Hyperanthe, *Hübner*, *Verz. bek. Schmett.* p. 61. n. 589 (1816).

Papilio Polymeda, *Scopoli*, *Ent. Carn.* p. 157. n. 434. fig. 434 (1763).

Var. Papilio Arete, Müller, *Faun. Fridr.* p. 36. n. 330.

Var. Papilio vidua, *ibid.* n. 331 (1764).

Prussia, Germany, France, South Europe.

B.M.

Genus 11. *ELINA*, Blanchard (1852).

Typical Species *Elina vanessoides*.

1. *Elina vanessoides*.

Elina vanessoides, Blanchard in *Gay's Fauna Chilena*, vii. p. 28, lam. 5. figs. 5, 6 (1852).

Chili.

2. *Elina Lefebvrei*.

Satyrus Lefebvrei, Guérin, *Voy. de la Coquille*, p. 281 (1829).

Satyrus Montrolii, Feisthamel, *Magas. de Zool.* ix. pl. 20 (1839).

Lasiommata Montrolii, Westwood in *Gen. Diurn. Lep.* p. 387. n. 15 (1851).

Elina Montrolii, Gay, *Fauna Chilena*, vii. p. 28, lam. 5. fig. 7 (1852).

Chili.

Genus 12. *EPINEPHELE*, Hübner.

Typical Species *Epinephele Janira*.

1. *Epinephele Eudora*.

♀. *Papilio* (Nymph. Gemm.) *Eudora*, Esper, *Schmett.* i. pt. 1. tab. 45. *Suppl.* xxi. fig. 1; ♂ ♀, pt. 2. tab. 69. cont. 19. figs. 1, 2 (1777).

Papilio Eudora, Fabricius, *Ent. Syst.* iii. pt. 1. p. 243. n. 755 (1793); Hübner, *Eur. Schmett.* i. pl. 36. figs. 163, 164 (♀ var.) (1805); also of Ochsenheimer.

Satyrus Eudora, Godart, *Enc. Méth.* ix. p. 541. n. 164 (1819); *Lép. de France*, ii. p. 128. n. 48, pl. 18. figs. 1-3 (1822).

Papilio Eudorus, Herbst, *Natursyst. Schmett.* viii. p. 163, tab. 205. figs. 5-8 (1796).

Papilio Lycaön, Müller, *Naturforsch.* iii. tab. 2. figs. d-f, vi. p. 17 (1784?).

Papilio Jurtina, Hübner, *Eur. Schmett.* i. pl. 36. fig. 160 (1805).

Germany.

B.M.

Var. ♀. Alia supra pallidioribus, anticis omnino ochreo flavescens, fascia anticarum quoque magis distincta: alis subtus multo pallidioribus.

Sweden.

B.M.

2. *Epinephele Rhamnusia*.

Papilio (Hipparchia) Rhamnusia, *Freyer, Neuere Beiträge*, v. p. 125. n. 817, tab. 457. figs. 2, 3 (1845).

Erebia Rhamnusia, *Herrich-Schäffer, Eur. Schmett.* i. figs. 377, 378 (♀), 427, 428 (♂) (1845).

Satyrus Rhamnusia, *Westwood in Gen. Diurn. Lepid.* p. 391. n. 33 (1851).

Mount Ætna (Freyer).

3. *Epinephele pulchella*.

Epinephele pulchella, *Felder, Reise der Fregatte Novara*, iii. tab. 69. fig. 16 (1867).

Eastern Himalayas (Felder).

Allied to *E. Eudora*.

4. *Epinephele pulchra*.

Epinephele pulchra, *Felder, Reise der Fregatte Novara*, iii. p. 491. n. 848 (1867).

Eastern Himalayas (Felder).

"Nearly allied to *E. pulchella*, Feld., but of the size of *E. Eudora*, Fabr."

5. *Epinephele Davendra*.

Epinephele Davendra, *F. Moore, Proc. Zool. Soc. London*, p. 502. n. 94, pl. 30. fig. 7 (1865).

Spiti and Tibet (Moore).

6. *Epinephele Cheena*.

Epinephele Cheena, *F. Moore, Proc. Zool. Soc. London*, p. 501. n. 93, pl. 30. fig. 6 (1865).

Kunawur (Moore).

7. *Epinephele Hispulla*.

♂ ♀. Papilio (Nymph. Gemm.) Hispulla, *Esper, Schmett.* i. pt. 2. tab. 119, cont. 74. figs. 1, 2 (1777).

♂ ♀. Papilio Hispulla, *Hübner, Eur. Schmett.* i. pl. 116. figs. 593-596 (1805).

Malta, South Europe.

♂ ♀, B.M.

8. *Epinephele pallescens*, sp. nov. Pl. III. fig. 3.

Alæ supra pallide olivaceo-fuscae, ciliis albidis: anticæ maris fascia interna basali cinerascens; margine externo paulum fuscescens, puncto subapicali nigro: feminæ macula disco-cellulari duabusque elongatis et puncto discalibus serie directa aurantiacis; ocello subapi-

cali nigro magno pupillis duobus oblique positis (inferiore minore), fulvoque irregulariter cincto: posticæ maris et feminæ undulatæ.

Alæ subtus fere velut in *Ida*, lineis mediis forma distantibus.

Exp. alar. unc. ♂ 2, ♀ 2½.

Persia.

♂ ♀, B.M.

9. *Epinephele Janira*.

♂. *Papilio Janira*, *Linnaeus*, *Syst. Nat.* i. p. 475 (1758), ii. p. 774. n. 156 (1766); also of *Fabricius* (1793).

Papilio (Nymph. Gemm.) *Janira*, *Esper*, *Schmett.* i. pt. 1. tab. 10. figs. 1, 2 (1777).

Papilio (*Hipparchia*) *Janira* (hermaphr.), *Freyer*, *Neuere Beiträge*, v. p. 138. n. 832, tab. 464. fig. 4 (1845).

Satyrus Janira, *Godart*, *Enc. Méth.* ix. p. 539. n. 163 (1819); also of *Zetterstedt* (*Ins. Lap.*) and *Westwood*.

Hipparchia Janira, *Stephens*, *Ill. Havst.* i. p. 59. sp. 9 (1828); also of *Westwood* and *Humphreys*.

Papilio Janirus, *Herbst*, *Natursyst. Schmett.* viii. p. 168. n. 101, tab. 206. figs. 7, 8, tab. 207. figs. 1, 2 (1796).

♀. *Papilio Jurtina*, *Linnaeus*, *Syst. Nat.* i. p. 475 (1758), ii. p. 774. n. 155 (1766); *Hübner*, *Eur. Schmett.* i. pl. 36. figs. 161, 162 (1805).

England (British Coll.), Germany.

B.M.

Var. JANIRULA.

Papilio (Nymph. Gemm.) *Janirula*, *Esper*, *Schmett.* i. pt. 2. tab. 113. cont. 68. fig. 1 (1777).

England.

♂, B.M.

This insect, though probably only a small variety of *Janira*, is in form sufficiently distinct for a species. The front wings are much more produced than in true *Janira*, so that it has somewhat the appearance of *E. Coctei*, a species placed by Westwood in the genus *Erebia*; the coloration of the underside is also somewhat similar to that of the latter species.

10. *Epinephele* ? *Servilia*.

Neomænas Servilia, *Wallengren* in *Kongl. Vet. Akad. Forhandl.* p. 78 (1858), and sp. descr. *Wien. ent. Monatschr.* iv. p. 36. n. 13 (1860). ? *Neomænas Cocti*, *Guér.*

Valparaiso.

11. *Epinephele monachus*.

Satyrus monachus, *Blanchard* in *Gay's Fauna Chilena*, vii. p. 35. n. 5 (1852).

Satyrus tristis, *ibid.* lam. 3. fig. 1 (1852).

Chili.

Allied to *E. Coctei*. The latter species is figured on the same plate and has been carelessly named *janirioides*. The figures of these species are not good, and only represent the upperside of the wings.

12. *Epinephele Pales*.

Satyrus Pales, *Philippi*, *Linnaea Entomologica*, xiv. p. 268. n. 5 (1860).

Chili (San Juan).

According to Philippi this is closely allied to *Coctei*; it seems only to differ in the absence of one or two points upon the underside of the hind wings; but not having seen the insect, it appears better to retain it as a distinct species.

13. *Epinephele Limonias*.

Satyrus Limonias, *Philippi*, *Linn. Ent.* xiv. p. 268. n. 6 (1860).

Chili (Valdivia).

Possibly another form of *Coctei*; it appears, however, to have greater claims to distinction than the preceding, the hind wings being more distinctly dentated with a ferruginous abbreviated band and with a reddish violet gloss.

14. *Epinephele Dryas*.

Epinephele Dryas, *Felder*, *Reise der Fregatte Novara*, iii. p. 492. n. 851 (1867).

Chili (Felder).

"Nearly allied to *Limonias*, but larger, the front wings with the costal, the hind wings with the inner margin longer, the outer margin of the front wings less waved."

15. *Epinephele Valdiviæ*.

Epinephele Valdiviæ, *Felder*, *Reise der Fregatte Novara*, iii. p. 493. n. 852 (1867).

Chili, Valdivia (Felder).

"In form of wing most like *E. Flora*" (*Hipparchia Flora*, antè), but considerably larger.

16. *Epinephele poliozona*.

Epinephele poliozona, *Felder*, *Reise der Fregatte Novara*, iii. p. 493. n. 853 (1867).

Chili, Valdivia (Felder).

A little larger than the male of *Coctei*; the wings entire, the outer margin of the front wings straight.

17. *Epinephele Coctei*.

Satyrus Coctei, *Guérin, Voyage Coquille, Zool. Ins.* pt. 2. div. 1. p. 281 (1832); *Mag. de Zool. Ins.* pl. 11 (1839).

Erebia Coctei, *Westwood in Gen. Diurn. Lepid.* p. 380. n. 52 (1851).

Var. Satyrus tristis, *Guérin, Voy. Coq. Zool.* p. 281.

Argynnis tristis, *ibid. Atlas, Ins.* n. 15. fig. 5 (1832).

Erebia tristis, *Westwood in Gen. Diurn. Lepid.* p. 380. n. 54 (1851).

Satyrus janirioides, *Blanchard in Gay's Fauna Chilena*, lam. 3. fig. 2 (1852).

Chili.

B.M.

18. *Epinephele Telmessia*.

Epinephele Telmessia, *Herrich-Schäffer, Eur. Schmett.* i. figs. 479-482 (1843-45).

Hipparchia Telmessia, *Zeller, Isis* (1847).

Satyrus Telmessia, *Westwood in Gen. Diurn. Lepid.* p. 391. n. 37 (1851).

Syria.

B.M.

19. *Epinephele nemyroides*.

Satyrus nemyroides, *Blanchard in Gay's Fauna Chilena*, vii. p. 33. n. 1. lam. 2. figs. 6-7 (1852); also of *Philippi* (1860).

Chili.

20. *Epinephele janirioides*.

Satyrus janirioides, *Blanchard in Gay's Fauna Chilena*, vii. p. 34. n. 2. lam. 2. fig. 8 (1852). *S. janirioides* on plate.

Chili.

The last two species are peculiar for the angular form of the front wings. The former somewhat resembles *Telmessia* above and *Ida* below; the latter appears to come near to *Tithonius*.

21. *Epinephele Narica*.

Papilio Narica, *Hübner, Eur. Schmett.* i. pl. 139. figs. 704-707 (1805).

Papilio (Hipparchia) Narica, *Freyer Neuere Beiträge*, v. p. 137. n. 831, tab. 464. figs. 2, 3 (1845).

Satyrus Narica, *Boisduval, Icon. Hist. Léop.* p. 212. n. 6, pl. 42. figs. 3-5 (1832); also of *Duponchel*.

South Ural.

22. *Epinephele Amaryllis*.

Papilio Amaryllis, *Geoffroy, Hist. Ins.* ii. p. 52. n. 20 (1764).

Satyrus Amaryllis, *Godart, Léop. de France*, i. p. 154. n. 51, pl. 7. fig. 2 (1821).

Papilio Amarillis (*sic*), *Ernst, Pap. d'Europe*, t. 1. p. 123, pl. 27. fig. 53 a-e (1779).

Papilio Tithonius, *Linnaeus, Mant.* i. p. 537 (1771).

Hipparchia Tithonus (*sic*), *Stephens, Ill. Haust.* i. p. 58. sp. 8 (1828).

Satyrus Tithonus, *Westwood in Gen. Diurn. Lepid.* p. 391. n. 36 (1851).

Papilio Herse, *Denis, Wien. Verz. fam. F.* p. 320. n. 8, 9 (1775).

Papilio pilosellæ, *Fabricius, Syst. Ent.* ii. p. 497 (1775).

Papilio (Nymph. Gemm.) *Phædra*, *Esper, Schmett.* i. pt. 1. tab. 9. fig. 1, tab. 28. *Suppl.* iv. fig. 3 (1777).

England (British Coll.), France.

B.M.

23. *Epinephele* *Ida*.

Papilio (Nymph. Gemm.) *Ida*, *Esper, Schmett.* i. pt. 2. tab. 92. cont. 42. fig. 2, tab. 102. cont. 57. fig. 3 (1777).

Papilio Ida, *Borkhausen, Naturgesch. eur. Schmett.* i. p. 105. n. 47 (1788).

Satyrus Ida, *Godart, Enc. Méth.* ix. p. 543. n. 167 (1819); *Lép. de France*, ii. p. 130. n. 49, pl. 18. figs. 4, 5 (1822).

Papilio Actæa, *Lang, Verz.* ii. p. 23. n. 165 (1789).

France, Malta, Germany.

B.M.

24. *Epinephele* *Roxane*.

Epinephele Roxane, *Felder, Reise der Fregatte Novara*, iii. tab. 69. figs. 12, 13 (1867).

Eastern Himalayas (Felder).

Allied to *E. Wagneri* and *E. Ida*. According to Dr. Felder, it is a local form of *E. Davendra* of Moore.

25. *Epinephele* *Pasiphaë*.

Papilio (Nymph. Gemm.) *Pasiphaë*, *Esper, Schmett.* i. pt. 2. tab. 67. cont. 17. fig. 4, tab. 97. cont. 52. fig. 1 (1777).

Papilio Pasiphaë, *Borkhausen, Naturgesch. eur. Schmett.* i. p. 97. n. 36 (1788).

Satyrus Pasiphaë, *Boisduval, Sp. Gén. Ind.* p. 32. n. 256 (1840).

Satyrus Bathseba, *Godart, Enc. Méth.* ix. p. 543. n. 108 (1819); *Lép. de France*, ii. p. 132. n. 50, pl. 18. figs. 6, 7 (1822).

Germany, France, South Europe.

B.M.

Var. *Papilio Bathseba*, *Fabricius, Ent. Syst.* iii. pt. 1. p. 235. n. 733 (1793). *Posticæ* subtus fascia alba, ocellis quatuor minusve.

Barbary (Fabricius).

- Var. Papilio Salome*, *Fabricius, ibid.* p. 238. n. 744 (1793).
Anticæ ocello unipupillato.
 Gibraltar (*Fabricius*).

26. *Epinephele Wagneri*.

Satyrus Wagneri, *Herrich-Schäffer, Eur. Schmett.* i. figs. 311–313 (1843–45).

Hipparchia Wagneri, *E. Doubleday, List Lep. Brit. Mus. Suppl.* p. 32 (1847).

Persia, Europe.

B.M.

♂. *Var. Satyrus Mandane*, *Kollar, Denkschr. k. Akad. Wissenschaft. Wien*, i. p. 52. sp. 8 (1850).

South Persia.

The description of this variety agrees entirely with specimens of *Wagneri* with which I have compared it.

Subgenus XENICA, *Westwood*.

Typical Species *Xenica Abeona*.

1. *Xenica Abeona*.

Papilio Abeona, *Donovan, Ins. New Holl.* i. pl. 22. fig. 1 (1805).
Satyrus Abeona, *Godart, Enc. Méth.* ix. p. 490. n. 72 (1819);
 also of *Boisduval*.

Hipparchia ? *Abeona*, *E. Doubleday, List Lep. Brit. Mus.* i. p. 134 (1844).

Lasiommata ? (*Xenica*) *Abeona*, *Westwood in Gen. Diurn. Lepid.* p. 387. n. 24 (1851).

Heteronympha Abeona, *Wallengren, Nov. Gen. Lepid. Kongl. Vet. Akad. Forhandl.* p. 78 (1858).

Oreas marmorea Zelinde, *Hübner, Exot. Schmett.* i. pl. 92. figs. 1, 2 (1806).

Tisiphone Zelinde, *Hübner, Verz. bek. Schmett.* i. p. 60, n. 574 (1816).

New Holland.

B.M.

2. *Xenica Joanna*.

Enodia (misprinted *Euodia*) *Joanna*, *Butler, Ann. & Mag. Nat. Hist.* 3rd ser. xvii. p. 286 (1866).

Enodia Joanna, *ibid.* xviii. pl. 4. fig. 8 (1867).

Australia.

B.M.

Genus 13. TISIPHONE, Westwood * (1851).

Typical Species *Tisiphone Hercyna*.1. *Tisiphone Hercyna*.

Tisiphone Hercyna, Hübner, *Samml. exot. Schm.* Bd. ii. pl. 86. figs. 3, 4 (1806); also of Westwood & Hewitson (1851).

Satyrus Anosia, Godart, *Enc. Méth.* ix. p. 453. n. 36 (1819).

Venezuela, Brazil, Rio Janeiro, New Granada, Mexico. B.M.

Genus 14. LEPTONEURA, Wallengren (1857).

Typical Species *Leptoneura Clytus*.1. *Leptoneura Clytus*.

Papilio Clytus, Linnaeus, *Syst. Nat.* ii. p. 768. n. 124 (1766); Cramer, *Pap. Exot.* i. pl. 86. figs. C, D (1779); also of Wulfen (1786) and Herbst (1796).

Satyrus Clytus, Godart, *Enc. Méth.* ix. p. 525. n. 132 (1819).

Erebia? *Clytus*, E. Doubleday, *List Lep. Brit. Mus.* i. p. 128 (1844).

Erebia Clytus, Westwood in *Gen. Diurn. Lepid.* p. 380. n. 43 (1851).

Leptoneura Clytus, Wallengren, *Lep. Rhop. Caffrar. in Kongl. Svenska Vetensk.* p. 31 (1857); also of Trimen (1866).

Dira Clyte, Hübner, *Verz. bek. Schm.* p. 60. n. 573 (1816).

Var. *Papilio* (Nymph. Gemm.) *Clytus*, Esper, *Schmett.* i. pt. 2. tab. 66. cont. 16. figs. 4. & ? 2, 3 (1777).

Cape of Good Hope, Knysna.

B.M.

Var. *Papilio Tisiphone*, Fabricius, *Gen. Ins. Mant.* p. 263 (1776?); *Ent. Syst.* iii. pt. 1. p. 214. n. 671 (1793); also of *Engramelle* (1780) and Borkhausen (1788).

Cape of Good Hope.

B.M.

2. *Leptoneura Mintha*.

Dira Mintha, Hübner, *Zuträge*, figs. 851, 852 (1806).

Erebia Mintha (? *Clytus*, var.), Westwood in *Gen. Diurn. Lepid.* p. 380. n. 48 (1851).

Leptoneura Mintha (? *Clytus*, var.), Trimen, *Rhop. Afric. Austr.* p. 193. n. 109 (part.) (1866).

South Africa.

? B.M.

The specimen which represents this species in the National Col-

* The type of Hübner's genus *Tisiphone* is the *Abeona* of Donovan, afterwards made into a subgenus by Westwood, under the name of *Xenica*, and still later united with *Merope* by Wallengren to form his genus *Heteronympha*.

lection does not quite agree with Hübner's figure, and may be distinct.

3. *Leptoneura* Cassus.

Papilio Cassus, *Linnaeus, Syst. Nat.* ii. p. 768. n. 125 (1766); *Cramer, Pap. Exot.* iv. pl. 314. figs. C, D (1782); also of *Herbst* (1796).

Satyrus Cassus, *Godart, Enc. Méth.* ix. p. 526. n. 133 (1819).

Erebia Cassus (? *Clytus* ♀)*, *Westwood in Gen. Diurn. Lepid.* p. 380. n. 44 (1851).

Leptoneura Cassus, *Trimen, Rhop. Afric. Austr.* ii. p. 195. n. 110 (1866).

Erebia ? *Clytus* (synonym), *E. Doubleday, List Lep. Brit. Mus.* i. p. 128 (1844).

Madagascar, interior of South Africa.

B.M.

4. *Leptoneura cassina*, sp. nov. Pl. II. fig. 12.

Alæ supra velut in *Casso*, minores et angulo anali anticarum magis rotundato, costaque brevior.

Alæ anticæ subtus velut in *Casso*, ocellis autem confluentibus majoribus extus albido marginatis; area anali quoque cinerascens, strisque distinctioribus: posticæ albido-cinereæ, area basali, fascia discali et margine externo fusciscentibus; lineis velut in *Casso* sed distinctis; ocellis magis elongatis, majoribus, nigris, albido cinctis.

Exp. alar. unc. 2.

South Africa.

B.M.

Closely allied to *Cassus*, which it more nearly resembles on the upper than the under surface of the wings.

Genus 15. *EREBIA*, *Dalman* (1816).

Typical Species *Erebia Ligea*, Sp. 18.

Section 1. *OREINA*, *Westwood* (1840).

Typical Species *Oreina Cassiope*.

1. *Oreina Epiphron*.

Papilio Epiphron, *Knoch, Beitrag*, iii. tab. 6. fig. 1 (1781-83).

Satyrus Epiphron, *Godart, Lép. de France*, ii. p. 113, pl. 16. figs. 3, 4 (1822).

Erebia Epiphron, *Herrich-Schäffer, Eur. Schmett.* figs. 92-94 (1843-45).

* Mr. Trimen has taken both sexes of this species, proving it to be distinct from *Clytus*. It is the type of Hübner's genus *Ypthima*; but it would create too great a commotion to rename the insects generally associated with that genus.

Papilio Egea, *Borkhausen, Naturgesch. eur. Schmett.* i. p. 77. n. 16, ii. p. 202. n. 16 (1788).

Papilio (*Hipparchia*) *Egea*, *Freyer, Neuere Beiträge*, vi. p. 133, tab. 567. fig. 1 ♂, 2 ♀ (1852).

Papilio *Ianthé*, *Hübner, Eur. Schmett.* i. pl. 44. fig. 202 (1805).

Papilio Cassiope, *Hübner, Eur. Schmett.* i. pl. 123. figs. 628, 629 (1805).

Erebia Cassiope, var.?, *Boisduval, Sp. Gén. Ind.* p. 26. n. 195 (1840).

England (British Coll.), Hercynia, Harz.

B.M.

In Knoch's figure the ocelli of the hind wings are subapical, and nearer to the margin than in the figures by Hübner and Godart; *O. Cassiope* is the commoner form of this species.

Var. CASSIOPE.

Papilio Cassiope, *Fabricius, Mant. Ins.* pt. 2. p. 42. n. 417 (1787); *Ent. Syst.* iii. pt. 1. p. 238. n. 742 (1793).

Papilio (*Hipparchia*) *Cassiope*, *Freyer, Neuere Beiträge*, i. p. 37, tab. 20. figs. 1, 2 (1833).

Hipparchia Cassiope, *Stephens, Ill. Haust.* i. p. 63, pl. 8. figs. 1-3 (1828).

Oreina Cassiope, *Westwood & Humphrey's Brit. Butterf.* p. 78, pl. 22. figs. 9, 10 (1840).

Erebia Cassiope, *Westwood in Gen. Diurn. Lepid.* p. 377. n. 1 (1851).

Papilio (*Nymph. Gemm.*) *Melampus*, *Esper, Schmett.* i. pt. 2. p. 131, tab. 78. cont. 28. (nec 31 et 103) fig. 2 (1777).

Papilio Melampus, *Herbst, Natursyst. Insekt.* viii. p. 186. n. 109 (1794).

Papilio Alcyone (part.), *Borkhausen, Naturgesch. eur. Schmett.* i. p. 96. n. 356 (1789).

Var. *Erebia Melampus*, *Newman, Zoologist*, ii. p. 729 (1844).

Var. *Papilio Mnemon*, *Haworth, Ent. Trans.* 1st ser. i. p. 332 (1812).

Melampias Rhodia, *Hübner, Verz. bek. Schmett.* n. 611 (1816).

Papilio Æthiops minor, *Villars, Ent.* ii. p. 37 (1834).

Var. *Erebia Nelamus*, *Boisduval, Sp. Gén. Ind.* p. 26. n. 195 (1840).

Var.?. *Papilio Cassiope*, *Hübner, Eur. Schmett.* i. pl. 123. figs. 626, 627 (1805).

England (British Coll.), Russia, Sweden.

B.M.

2. *Oreina Arete*.

Papilio Arete, *Fabricius, Mant. Ins.* ii. p. 42. n. 418 (1787); *Ent. Syst.* iii. pt. 1. p. 238. n. 743 (1793); *Hübner, Eur. Schmett.* i. pl. 50. figs. 231, 232 (1805).

Erebia Arete (Blandina, var. ?), *Westwood in Gen. Diurn. Lepid.* p. 377. n. 3 (1851).

Papilio Aretus, *Herbst, Natursyst. Insekt.* viii. p. 199. n. 117 (1796).

Papilio Claudina, *Borkhausen, Naturgesch. eur. Schmett.* i. p. 204. n. 161 (1788).

Southern Alps.

Professor Westwood has queried this species as a variety of *Blandina*, but it seems to be much more nearly allied to *Epiphron*. M. Chavignerie has figured the typical *Melampus*.

3. *Oreina Pharte*.

Papilio Pharte, *Esper, Schmett.* i. pt. 2. p. 17, tab. 120, cont. 75. fig. 3 ♂, 4 ♀ (1777); *Hübner, Eur. Schmett.* i. figs. 491-494 (1805).

Papilio (*Hipparchia*) *Pharte*, *Freyer, Neuere Beiträge*, i. p. 37, tab. 20. fig. 3 (1833).

Satyrus Pharte, *Godart, Enc. Méth.* ix. p. 536. n. 156 (1819).

Erebia Pharte, *Boisduval, Icon. Hist. Lép.* p. 180, pl. 35. figs. 1, 2 (1832).

Var. Satyrus Pharte, *Duponchel, Lép. de France*, i. p. 218, pl. 34. figs. 1, 2 (1832).

Var. Erebia Euryale, *Herrich-Schäffer, Eur. Schmett.* i. fig. 101 (1843-45).

Tyrol.

Coll. Salvin.

Var. ??. *Herrich-Schäffer, Eur. Schmett.* i. fig. 95.

Germany.

B.M.

Esper describes this species as chiefly differing from *Melampus* in the absence of black spots upon the red band. It appears to be much more closely allied to *Cassiope*, the front wings being more pointed at the apex than in *Melampus*.

4. *Oreina Kefersteinii*.

Erebia Kefersteinii, *Herrich-Schäffer, Eur. Schmett.* i. figs. 617, 618 (1843-45); *Eversmann, Bull. Soc. Nat. Mosc.* p. 610 (1851); also of *Westwood* (1852).

Papilio Theano, *Tauscher, Mém. Soc. Nat. Mosc.* i. p. 207, tab. 13. fig. 1.

Hipparchia (*Erebia*) *Theano*, *Eversmann, Bull. Mosc.* p. 611 (1851).

Siberia.

Near *Pyrenaica* (H.-Sch.), but the front wings ferruginous above.

5. *Oreina ? Stubbendorffii*.

Erebia Stubbendorffii, *Ménétriés in Bull. Acad. St. Pétersb.* v. p. 262 (1847).

Siberia.

6. *Oreina Melampus*.

Papilio Melampus, *Fuessly, Verz. Schw. Ins.* p. 31. n. 604, tab. 1. fig. 6 (1775).

Papilio (Hipparchia) Melampus, *Freyer, Neuere Beiträge*, i. p. 35, tab. 19. figs. 1, 2 (1833).

Satyrus Melampus, *Godart, Enc. Méth.* ix. p. 536. n. 155 (1819); *Lép. de France*, ii. p. 114. n. 42, pl. 16. figs. 5, 6 (1822).

Erebia Melampus, *Boisduval, Icon. Hist. Lép.* p. 178. n. 16, pl. 35. figs. 5, 6 (1832).

Papilio (Nymph. Gemm.) ætherius, *Esper, Schmett.* i. pt. 2. tab. 122. cont. 77. fig. 4, nec 3 (1777).

Papilio Alcyone (part.), *Borkhausen, Naturgesch. eur. Schmett.* i. p. 96. n. 356 (1789).

Erebia Arete, *Bellier de la Chavignerie, Ann. Soc. Ent. France*, ser. 3, tome vi. p. 485, pl. 2. fig. 2, ♂ ♀ (1858).

Germany, Switzerland, Spain, Sweden.

B.M.

Var. PYRENAICA.

Erebia Cassiope, *var. Pyrenaica*, *Herrich-Schäffer, Eur. Schmett.* i. figs. 535-538 (1843-45).

Tyrol.

Coll. Salvin.

Var. MNESTRA, sp. ??

Papilio Mnestra, *Esper, Schmett.* i. pt. 2. tab. 120. cont. 75. figs. 5, 6 (1777).

Papilio (Hipparchia) Mnestra, *Freyer, Neuere Beiträge*, i. p. 164. n. 165, tab. 91. fig. 3 ♂ (1833).

Satyrus Mnestra, *Godart, Enc. Méth.* ix. p. 533. n. 149 (1819).

Erebia Mnestra, *Herrich-Schäffer, Eur. Schmett.* i. fig. 96 (1843-45).

Germany, Sweden.

B.M.

Var. Papilio Mnestra, *Hübner, Eur. Schmett.* i. pl. 106. figs. 540-543 (1805).

Papilio (Hipparchia) Mnestra, *Freyer, Neuere Beiträge*, i. p. 36. n. 31, tab. 19. fig. 3, ♂ (1833).

Satyrus Mnestra, *Duponchel, Lép. de France*, i. p. 220. n. 77, pl. 34. figs. 3, 4 (1832).

Erebia Mnestra, *Boisduval, Icon. Hist. Lép.* p. 176. n. 15, pl. 35. figs. 1-4 (1832).

Mountains of Switzerland.

Var. ERIPHYLE.

Papilio (Hipparchia) Eriphyle, *Freyer, Neuere Beiträge*, ii. p. 150. n. 325, tab. 187. figs. 3, 4 (1836).

Erebia Cassiope, var. *Eriphile*, *Westwood in Gen. Diurn. Lepid.*
p. 377. n. 1 (1851).

Europe (locality not marked).

B.M.

Var. IANTHE.

Papilio Ianthæ, *Hübner, Eur. Schmett. i.* pl. 122. figs. 624, 625
(1805).

Erebia Cassiope, var. *Ianthæ*, *Westwood in Gen. Diurn. Lepid.*
p. 377. n. 1 (1851).

Germany.

B.M.

There are specimens intermediate between the last two forms
from the Tyrol in the Collection of Mr. Osbert Salvin.

7. *Oreina Pyrrha.*

Papilio Pyrrha, *Denis, Wien. Verz.* p. 167. fam. F. n. 9 (1775);
Hübner, Eur. Schmett. i. p. 35. n. 51, pl. 50. figs. 235, 236
(1805).

Papilio (Hipparchia) Pyrrha, *Freyer, Neuere Beiträge*, i. p. 58.
n. 53, tab. 31. figs. 3, 4 (1833).

Satyrus Pyrrha, *Godart, Lép. de France*, ii. p. 108. n. 38, pl. 15.
figs. 3, 4 (1822).

Erebia Pyrrha, *Westwood in Gen. Diurn. Lepid.* p. 377. n. 8
(1851).

Var. Papilio Maccabæus, *Herbst, Natursyst. Insekt.* viii. p. 185.
n. 108, (*Maccabæus*) tab. 209. figs. 5, 6 (1796).

Erebia Pyrrha, var. *Machabæus*, *Westwood* (1851).

Var. Papilio Erina, *Fabricius, Mant. Ins.* ii. p. 41. n. 415
(1787).

Var. ? Papilio (Nymph. Gemm.) Maurisius, *Esper, Schmett. i.*
pt. 2. tab. 113. cont. 68. figs. 4, 5 (1777).

Russia, Sweden.

B.M.

Var. MANTO (not Erebia Manto).

♀. *Papilio (Nymph. Gemm.) Manto*, *Esper, Schmett.* tab. 70.
figs. 2, 3 (1777).

Papilio Manto, *Herbst, Natursyst. Insekt.* viii. p. 194, tab. 210.
figs. 8, 9, tab. 211. figs. 1, 2 (1796).

Papilio Pyrrha, *Hübner, Eur. Schmett. i.* pl. 120. fig. 616
(1805).

Papilio (Hipparchia) Pyrrha, *Freyer, Neuere Beiträge*, i. tab. 91.
fig. 4 (1833).

♂ ♀. Tyrol (Coll. Salvin). ♀. Germany.

B.M.

We have no female specimens from Northern Europe.

Var. MÆRENS.

Erebia mærens, *Westwood in Gen. Diurn. Lepid.* p. 380. n. 40 (1851).

Erebia tristis, *Herrich-Schäffer, Eur. Schmett.* i. figs. 387-390 (1843-45).

Sweden.

B.M.

8. *Oreina Pluto.*

Papilio (Nymph. Gemm.) *Pluto*, *Esper, Schmett.* i. pt. 2. tab. 121, cont. 76. fig. 1 (1777).

Papilio Morio, *Giorna, Calend. Entom. Torino*, p. 102. n. 24 (1791).

Papilio petrosus, *De Prunner, Lep. Ped.* p. 71 (1798).

Erebia Pyrrha, var., *Westwood in Gen. Diurn. Lepid.* p. 377. n. 8 (1851).

Russia.

B.M.

This species is entirely black brown, and not the same as *Pyrrha*.

Var. TISSIPHONE.

Papilio (Nymph. Gemm.) *Tissiphone*, *Esper, Schmett.* i. pt. 2. tab. 122. cont. 77. fig. 5 (1777).

Papilio Cæcilia, *Hübner, Eur. Schmett.* i. pl. 46. figs. 213, 214 (1805), nec *Esper*.

Erebia Cæcilia, *Herrich-Schäffer, Eur. Schmett.* i. figs. 584, 585 (1843-45).

Papilio Alecto, *Freyer, Neuere Beiträge*, i. p. 99. n. 89, tab. 49. fig. 3 (1833).

Russia.

B.M.

9. *Oreina glacialis.*

Papilio (Nymph. Gemm.) *glacialis*, *Esper, Schmett.* i. pt. 2. tab. 116. cont. 71. fig. 2 (1777).

Var. Papilio (Nymph. Gemm.) *Erynys*, *ibid.* tab. 121, cont. 76. fig. 4 [nec 5] (1777).

Var. Papilio Alecto, *Hübner, Eur. Schmett.* i. pl. 104. figs. 528, 529 (1805).

Papilio (Hipparchia) *Alecto*, *Freyer, Neuere Beiträge*, i. tab. 49. figs. 3, 4 (1833).

Erebia Alecto, *Boisduval, Icon. Hist. Lép.* p. 160. n. 7, pl. 32. figs. 4-7 (1832).

Germany.

B.M.

Esper's figure of *glacialis* appears to be somewhat too strongly defined, but there is little doubt that *Alecto* of *Hübner* is merely a variety of it.

Section 2. *EREBIA* proper.1. *Erebia Alecto*.

Papilio Alecto, *Hübner, Eur. Schmett.* i. pl. 101. figs. 515, 516 (1805).

Var. Erebia Scipio, *Boisduval, Icon. Hist. Lépid.* p. 152. n. 3, pl. 30. figs. 1, 2 (var. 3, 4) (1832).

Var. Satyrus Scipio, *Duponchel, Lép. de France, Suppl.* p. 243. n. 88, pl. 38. figs. 5, 6 (1832).

Satyrus Gorge, *Godart, Enc. Méth.* ix. p. 527. n. 136 (1819); *Lép. de France*, ii. p. 120. n. 45, pl. 14. figs. 3, 4 (1822); but not of *Esper*.

South France, Pyrenees.

B.M.

Var. NERINE.

Erebia Nerine, *Boisduval, Icon. Hist. Lépid.* p. 154. n. 4, pl. 31. figs. 6, 7 (1832).

Var. Papilio (Hipparchia) Nerine, *Freyer, Neuere Beiträge*, p. 26. n. 18, tab. 13. figs. 3, 4 (1833).

Var. ? Erebia Nerine, *Herrich-Schäffer, Eur. Schmett.* i. figs. 69, 70, 73, 74 (1843-45).

Spain.

B. M.

2. *Erebia Pirene*.

Papilio (Nymph. Gemm.) Pirene, *Esper, Schmett.* i. pt. 2. tab. 116. cont. 71. fig. 3 (1777).

Erebia Pyrene, *Westwood in Gen. Diurn. Lepid.* p. 378. n. 13 (1851).

Var. Papilio (Nymph. Gemm.) Persephone, *Esper, Schmett.* i. pt. 2. tab. 121. figs. 5, 6 (1777).

Var. Erebia Lefebvrei, *Herrich-Schäffer, Eur. Schmett.* figs. 88, 89 (1843-45); but not of *Boisduval*.

Germany, Switzerland, Sweden.

B.M.

Var. STYGNE.

Satyrus Stygne, *Godart, Enc. Méth.* ix. p. 534. n. 150 (1819); *Lép. de France*, ii. p. 100. n. 34, pl. 14. figs. 1, 2 (1821).

Erebia Stygne, *Herrich-Schäffer, Eur. Schmett.* i. figs. 90, 91 (1843-1845).

Prussia, British Columbia.

B.M.

3. *Erebia Kalinda*.

Erebia Kalinda, *F. Moore, Proc. Zool. Soc.* p. 501. n. 92, pl. 30. fig. 5 (1865).

North-west Himalayas (Moore).

This species seems to come best next to *Pirene*; the coloration, however, is very distinct.

4. *Erebia Maurus*.

Papilio Maurus, *Esper*, *Schmett.* i. pt. 2. tab. 107. cont. 62. figs. 3, 4, tab. 110. cont. 65. figs. 4 (1777).

Pyrenees.

B.M.

Var. LEFEBVREI.

Erebia Lefebvrei, *Boisduval*, *Icon. Hist. Lépid.* p. 164. n. 9, pl. 33. figs. 1, 2 (1832).

Papilio Pyrene, *Freyer*, *Neuere Beiträge*, i. p. 82. n. 76, tab. 43. fig. 2 (1833); but not of *Esper*.

Pyrenees.

B.M.

Var. NELO.

Papilio Nelo, *Hübner*, *Eur. Schmett.* i. pl. 45. figs. 105, 106 (1805).

Satyrus Melas, *Godart*, *Enc. Méth.* ix. p. 534. n. 151 (1819); *Lép. de France*, p. 102. n. 35, pl. 17. figs. 1, 2 (1822); but not of *Herbst*.

Erebia Lefebvrei, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 280–282 (1843–45); but not of *Boisduval*.

Var. Satyrus Lefebvrei, *Duponchel*, *Lép. de France*, i. p. 226. n. 80, pl. 35. figs. 3, 4 (1832).

Pyrenees.

B.M.

Var. MELAS.

Papilio Melas, *Herbst*, *Natursyst. Schmett.* viii. p. 191. n. 112, tab. 210. figs. 4–7 (1796).

Papilio (*Hipparchia*) *Melas*, *Freyer*, *Neuere Beiträg.* i. p. 98. n. 88, tab. 49. fig. 2, p. 118. n. 111, tab. 61. figs. 1, 2 (1833).

Erebia Melas, *Boisduval*, *Icon. Hist. Lépid.* p. 166. n. 10, pl. 33, figs. 3, 4 (1832).

Var. Erebia Melas, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 65–68, 467, 468 (1843–45).

Sweden.

B.M.

Var. a. ♂. *Alæ* subtus ocello subanali anticarum obsoleto, fascia transversali media posticarum distincta.

Russia.

B.M.

Var. b. ♂. *Alæ* minores, costa anticarum magis convexa, apicibus alarum subtus pallescentibus, aliter velut in præcedente.

Spain.

B.M.

Var. c. *Satyrus Melas*, *Duponchel*, *Lép. de France*, i. p. 245: n. 89, pl. 39. figs. 1-4 (1832).

Spain.

B.M.

5. *Erebia Arachne*.

Papilio Arachne, *Fabricius*, *Mant. Ins.* pt. 2. p. 41. n. 415 (1787); *Ent. Syst.* iii. pt. 1. p. 237. n. 740 (1793).

Satyrus Arachne, *Godart*, *Enc. Méth.* ix. p. 529. n. 140 (1819); *Lép. de France*, ii. p. 116. n. 43, pl. 16. figs. 7, 8 (1822).

* *Papilio Pronoe*, *Ochsenheimer*, *Schmett. v. Eur.* i. p. 290. n. 59 (1807).

Papilio (*Hipparchia*) *Pronoe*, *Freyer*, *Neuere Beiträge*, i. p. 137. n. 133, tab. 73. figs. 3, 4 (1833).

Papilio (*Hipparchia*) *Styx*, *Freyer*, *Neuere Beiträge*, i. p. 44. n. 212, tab. 121. fig. 4 (1833).

Germany.

B.M.

Local form. Alæ supra fasciis rufescentibus magis distinctis, lineis transversis, subtus magis ad margines approximantibus.

Switzerland.

B.M.

Var. *PITHO*.

Papilio Pitho, *Hübner*, *Eur. Schmett.* i. pl. 112. figs. 574 & 577, var. fig. 576 (1805); *Freyer*, *Stett. ent. Zeit.* p. 92 (1847).

Switzerland.

B.M.

Var. Anticæ oculo subanali, subtus area disco-cellulari rufescente; posticæ subtus violaceo-cinerascentes.

Tyrol.

Coll. Salvin.

6. *Erebia Epipsodea*, sp. nov. Pl. II. fig. 9.

Alæ supra forma et coloribus fere *Psodea* (Hübner.), ocellis autem posticarum et tertio anticarum cæcis, aliis subcæcis, fascia quoque apud apicem magis extensa.

Alæ anticæ subtus velut in *Psodea* sed magis rufescentibus: posticæ fuscæ cinereo roratæ, fascia angulari sat lata media rufo-fusca, illius *Disa* fere forma, sed ad angulum ani magis elongata; ocellis tribus nigris cæcis ferrugineo distincte cinctis: corpus nigro-fuscum; antennis supra nigris, subtus albidis, fulvo acuminatis.

Exp. alar. unc. $1\frac{1}{8}$.

Rocky Mountains (two specimens).

B.M.

This species may possibly belong to the *Disa* group; but as it bears a greater general resemblance to *Arachne*, it appears better to place it here.

* *Ochsenheimer* describes the true *Arachne*, but quotes *Hübner's* figure.

7. *Erebia Medea*.

- Papilio Medea*, Denis, *Wien. Verz.* p. 167. n. 7 (1775).
Papilio Medea, Hübner, *Eur. Schmett.* i. pl. 48. figs. 220–222 (1805).
Papilio (*Hipparchia*) *Medea*, Freyer, *Neuere Beiträge*, i. tab. 38. fig. 3, p. 109. n. 100, tab. 55. figs. 1, 2 (1833).
Papilio Blandina, Fabricius, *Mant. Ins.* ii. p. 41 (1787); *Ent. Syst.* iii. 1. p. 236. n. 736 (1793).
Hipparchia Blandina, Stephens, *Ins. Haustell.* i. p. 62. sp. 12 (1828).
Oreina Blandina, Westwood and Humphreys, *Brit. Butterf.* p. 77, pl. 23. figs. 5–10? (1857).
♀. *Papilio Medusa*, Borkhausen, *Naturgesch. Schmett.* i. p. 75. n. 15 (1788).
Erebia Medusa, Dalman in *Kongl. Vetensk.* p. 81. n. 9 (1816).
Papilio (*Hipparchia*) *Neoridas*, *ibid.* p. 110. n. 101, tab. 55. figs. 3, 4 (1833); but not of Boisduval.
♀. *Papilio Alcyone*, Stewart, *Elem. Nat. Hist.* 2nd ed. ii. p. 132 (1817).
Var. ♂. *Papilio Æthiops*, Herbst, *Natursyst. Schmett.* viii. p. 184. n. 107, tab. 209. figs. 3, 4 (1796).
Scotland (Coll. Brit.), Russia, Prussia. B.M.

8. *Erebia Neoridas*.

- Erebia Neoridas*, Boisduval, *Icon. Hist. Léop.* p. 148. n. 1, pl. 29. figs. 1–4 (1832).
Hipparchia Neoridas, Treitschke, *Schmett. v. Eur. Suppl.* x. p. 51 (1834).
Papilio (*Nymph. Gemm.*) *Pronoë*, Esper, *Schmett.* i. pt. 2. tab. 54. cont. 4. fig. 1 (1777).
Papilio Pronoë, Borkhausen, *Naturgesch. Schmett.* i. p. 98. n. 37 (1788).
Papilio Arachne, Hübner, *Eur. Schmett.* i. pl. 47. figs. 215, ? 216, 217 (var.) (1805).
Var. *Satyrus Neoridas*, Duponchel, *Lép. de France*, i. p. 234. n. 84, pl. 36. figs. 5, 6 (1832).
♀. *Hipparchia Stygne*, Fischer, *Entom. Russ.* i. p. 56; *Lep.* tab. 1. fig. 2 (1820).
Var. ? *Papilio Medea*, Hübner, *Eur. Schmett.* i. pl. 48. figs. 220–222 (1805).
Papilio (*Hipparchia*) *Medea*, Freyer, *Neuere Beiträge*, i. p. 72. n. 68, tab. 38. fig. 2 (1833).
Var. ? *Erebia Sedakovii*, Herrich-Schäffer, *Eur. Schmett.* i. figs. 591, 592 (1843–1845).
Germany, Pyrenees. B.M.

We have a German variety of this species, having the front wings ferruginous on the underside, as represented by Esper (*Nymph. Gemm. Pronoë*). Fischer's *Stygne* seems to be a variety of the female.

9. *Erebia Euryale*.

Papilio (Nymph. Gemm.) *Euryale*, *Esper*, *Schmett.* i. pt. 2. tab. 118. cont. 73. figs. 2, 3 (1777).

Satyrus Euryale, *Godart*, *Enc. Méth.* ix. p. 533. n. 148 (1819); *Lép. de France*, ii. n. 33, pl. 8. figs. 3, 4 (1822).

Papilio (*Hipparchia*) *Euryale*, *Freyer*, *Neuere Beiträge*, i. p. 163. n. 161, tab. 91. figs. 1, 2; var.?, tab. 184. figs. 908, 909 (1833).

Erebia Euryale, *Boisduval*, *Sp. Gén. Ind.* p. 28. n. 219 (1840).

Var. *Papilio Euryale*, *Hübner*, *Eur. Schmett.* i. pl. 158. figs. 789, 790 (♀) (1807).

Var. *Papilio Adyte*, *Hübner*, *Eur. Schmett.* i. pl. 151. figs. 759, 760 (1805).

Var. *Papilio* (*Hipparchia*) *Euryale*, *Freyer*, *Neuere Beiträge*, i. p. 117. n. 110, tab. 61. figs. 3, 4 (1833).

Erebia Euryale, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 97–100 (1843–45).

♂. *Hipparchia Ligea* (part.), *Stephens*, *Ill. Haust.* pl. 6. fig. 1 (1828).

Ile of Arran? (Coll. Brit.), Tyrol (Coll. Salvin), Northern Europe. B.M.

Var. *PHILOMELA*.

Papilio (Nymph. Gemm.) *Philomela*, *Esper*, *Schmett.* i. pt. 2. tab. 116. cont. 71. fig. 4 (1777).

Papilio Philomela, *Hübner*, *Eur. Schmett.* i. tab. 47. figs. 218, 219 (1805).

Switzerland.

B.M.

Note.—Hübner's figure of *Euryale* (pl. 158) is probably somewhat exaggerated; the underside colouring of the hind wings seems too grey to be natural. To judge by the males alone, it might appear that *Euryale* was only an extreme form of *E. Ligea*; specimens of the latter from the Tyrol seem only to differ in the form of the front wings, and in the more sharply defined orange band on the underside, but the females are very distinct, and the two species are not found to fly together; *Euryale* seems to prefer damp situations. See Ent. Month. Mag. (December 1867).

10. *Erebia Ligea*.

Papilio Ligea, *Linnaeus*, *Syst. Nat.* (1735) 12th ed. p. 772. n. 144 (1766); *Fauna Suecica*, 2nd ed. p. 275. n. 1050 (1761); *Hübner*, *Eur. Schmett.* i. tab. 49. figs. 225–227 (1805).

Papilio (*Hipparchia*) *Ligea*, *Freyer*, *Neuere Beiträge*, i. p. 125. n. 121, tab. 67 (1833).

Erebia Ligea, *Dalman in Kongl. Vetensk. Akad. Handl.* pp. 58 (generis typus) & 81 (1816).

Satyrus Ligea, *Godart*, *Enc. Méth.* ix. p. 532. n. 147 (1819); *Lép. de France*, p. 96. n. 32, pl. 13. figs. 1, 2 (1822).

♀. *Hipparchia Ligea*, *Stephens*, *Ill. Haust.* i. p. 61. sp. 11, pl. 6. figs. 2, 3 (1828).

Oreina Ligea, *Westwood & Humphrey's Brit. Butterf.* p. 77. sp. 1, pl. 23. figs. 1-4 (1841).

Papilio Alexis, *Linnaeus, Fauna Suecica*, 1st ed. p. 239 (1746).

Papilio (Nymph. Gemm.) *Alexis*, *Esper, Schmett.* i. pt. 1. tab. 44, *Suppl.* xxii. figs. 1, 2, pt. 2. tab. 54. fig. 2 (var.) (1777).

Tyrol (Coll. Salvin), Switzerland.

B.M.

Var. SEDAKOVII.

Hipparchia Sedakovii, *Eversmann, Bull. Soc. Imp. Nat. Mosc.* xv. tab. 1. figs. 5, 6 (1847).

Erebia Sedakovii, *Westwood in Gen. Diurn. Lepid.* p. 379. n. 30 (1851).

Germany.

B.M.

This is only a slight variation of *E. Ligea*.

Var. LIGEAE.

Alæ anticae supra ocellis quatuor, tertio minimo, subtus tribus solum.

Tyrol.

Coll. Salvin.

Var. EUMONIA.

Erebia Eumonia, *Ménétriés, Cat. Lép. Acad. St. Pétersb.* ii. p. 104. n. 710 (1857); *Bull. Acad.* t. xvii. p. 216. n. 14 (1859); *Schrank, Lep. Amur-Lande*, p. 34. n. 68, tab. 3. fig. 4 (1859).

Germany.

B.M.

11. *Erebia Melusina*.

Erebia Melusina, *Herrich-Schäffer, Eur. Schmett.* i. figs. 373, 374 (1843-45).

Europe.

12. *Erebia Bonellii*.

Papilio Bonellii, *Hübner, Eur. Schmett.* i. pl. 181. figs. 892-895 (1805).

Papilio (*Hipparchia*) *Bonellii*, *Freyer, Neuere Beiträge*, i. p. 137. n. 132, tab. 73. figs. 1, ? 2 (1833).

Erebia Evias, *Boisduval, Icon. Hist. Lép.* p. 156. n. 5, pl. 31. figs. 3-6 (1832).

Russia, Pyrenees.

B.M.

Var. EVIAS.

Erebia Evias, *Lefebvre, Ann. Soc. Linn. Paris*, pp. 488, 490, pl. 10 (1826); *Duponchel, Lép. de France*, i. p. 236. n. 85, pl. 37. figs. 1, 2 (1832).

? *Hipparchia Medea*, *Eversmann, Fauna Lepid. Volgo-Ural.* p. 33. n. 25 (1844).

Var. Papilio Pirena, *Hübner, Eur. Schmett.* i. pl. 48. figs. 223, 224 (1805).

Germany.

B.M.

13. *Erebia Hewitsonii*.

Erebia Hewitsonii, *Lederer, Wien. ent. Monatschr.* viii. p. 167, Taf. 3. figs. 6 & 7 (1864).

Imeretia.

Coll. Hewitson.

14. *Erebia Epistygne*.

Papilio Epistygne, *Hübner, Eur. Schmett.* i. pl. 173. figs. 855–858 (1805).

Erebia Epistygne, *Boisduval, Icon. Hist. Lép.* p. 158. n. 6, pl. 31. figs. 1, 2 (1832).

Satyrus Epistygne (*sic*), *Duponchel, Lép. France*, i. p. 228. n. 86, pl. 37. figs. 3–6 (1832).

Papilio (*Hipparchia*) *Stygne*, *Freyer, Neuere Beiträge*, i. p. 97. n. 87, tab. 49. fig. 1 (1833).

Var. Papilio Stygne, *Hübner, Eur. Schmett.* i. pl. 125. figs. 639, 640 (1805).

Germany.

B.M.

15. *Erebia Goante*.

♀. *Papilio* (*Nymph. Gemm.*) *Goante*, *Esper, Schmett.* i. pt. 2. tab. 116. cont. 71. fig. 1 (1777).

Papilio Goante, *Ochsenheimer, Schmett. v. Eur.* i. p. 293. n. 60 (1807).

Satyrus Goante, *Godart, Enc. Méth.* ix. p. 530. n. 141 (1819); *Lép. de France*, ii. p. 118. n. 44, pl. 17. figs. 3, 4 (♂) (1822).

♂. *Erebia Goante*, *Boisduval, Sp. Gén. Ind.* p. 28. n. 222 (1840).

Papilio Scæa, *Hübner, Eur. Schmett.* i. pl. 50. figs. 233, 234 (1805).

♂. *Erebia Gorgone*, *Herrich-Schäffer, Eur. Schmett.* i. figs. 75, 76.

♂. *Erebia melancholica*, *ibid.* figs. 276, 277 (1843–45).

Germany.

B.M.

16. *Erebia Gorgone*.

Erebia Gorgone, *Boisduval, Icon. Hist. Lép.* p. 150. n. 2, pl. 29. figs. 5–8 (1832).

Hipparchia Gorge (*var. Gorgone*), *Treitschke, Schmett. v. Eur. Suppl.* x. 1. p. 231 (1834).

Pyrenees.

B.M.

This is a very variable species, but it may be readily distinguished by the form of the central bands on the underside.

17. *Erebia Gorgophone*.

Erebia Gorgophone, *Bellier, Ann. Ent. Soc. France*, iii. ser. 4. p. 419, pl. 9. figs. 1, 2 (1863).

Erebia Nerine, *Duponchel, Lép. France*, p. 35. figs. 5, 6 (1832), but not of *Boisduval*.

Erebia Gorgone, *Herrich-Schäffer, Eur. Schmett.* i. figs. 469, 470 (1843-45).

Tyrol.

♂, B.M.

18. *Erebia Gorge*.

♂. *Papilio* (Nymph. Gemm.) *Gorge*, *Esper, Schmett.* i. pt. 2. tab. 119. cont. 74. figs. 4, 5 (1777).

Papilio (Hipparchia) *Gorge*, *Freyer, Neuere Beiträge*, i. p. 147. n. 146 (1833).

Erebia Gorge, *Boisduval, Sp. Gén. Ind.* p. 28. n. 222 (1840).

Papilio (Nymph. Gemm.) *Erynys*, *Esper, Schmett.* i. pt. 2. tab. 121. cont. 76. fig. 3 (1833).

♀. *Erebia Goante*, *Herrich-Schäffer, Eur. Schmett.* i. figs. 77-79.

Erebia Gorgone, *ibid.* figs. 283, 284 (1843-45).

♂ ♀. *Var. ?*. *Papilio Gorge*, *Hübner, Eur. Schmett.* i. pl. 99. figs. 502-505 (1805).

Var. *Papilio* (Hipparchia) *George* (sic), *Freyer, Neuere Beiträge*, i. tab. 79. fig. 3 (1833).

Tyrol (♀, Coll. Salvin), Switzerland, Germany.

♂, B.M.

19. *Erebia Tyndarus*.

Papilio (Nymph. Gemm.) *Tyndarus*, *Esper, Schmett.* i. 2. tab. 67. cont. 17. fig. 1 (1777).

Papilio Herse, *Borkhausen, Naturgesch. Eur. Schmett.* i. p. 94. n. 33 (1788).

Papilio Tyndarellus, *Herbst, Natursyst. Schmett.* viii. p. 135. n. 74, tab. 202. figs. 5, 6 (1796).

Mountains of Switzerland (Esper).

Var. *Ocellis anticarum majoribus, albo pupillatis.*

Sweden.

B.M.

Var. *Dromus.*

Papilio Dromus, *Fabricius, Ent. Syst.* iii. pt. 1. p. 224. n. 701 (1793).

Switzerland (one specimen).

B.M.

Var. CASSIOIDES.

Papilio cassioides, *Esper, Schmett.* i. pt. 2. tab. 103. cont. 58. figs. 2, 3 (1777).

Papilio Cleo, *Hübner, Eur. Schmett.* i. pl. 46. figs. 209–212 (1805).

Satyrus Dromus, *Godart, Enc. Méth.* ix. p. 528. n. 138 (1819); *Lép. de France*, ii. p. 122. n. 46; *ibid.* (as *Cleo*) pl. 17. figs. 5, 6 (1822).

Erebia Dromus, *Westwood in Gen. Diurn. Lepid.* p. 379. n. 38 (1851).

Papilio Tyndarus, *Ochsenheimer, Schmett. Eur.* i. p. 299. n. 63 (1807).

Pyrenees, Switzerland, Sweden.

B.M.

Var. NELEUS.

Papilio (*Hipparchia*) *Neleus*, *Freyer, Neuere Beiträge*, i. p. 148. n. 148, tab. 80. figs. 1, 2.

Papilio Tyndarus, *ibid.* figs. 3, 4 (1833).

Germany.

B.M.

20. *Erebia Ottomana*.

♂. *Erebia Dromus*, var. *Ottomana*, *Herrich-Schäffer, Eur. Schmett.* i. fig. 376.

♀? *ibid.* figs. 379, 380 (1843–45).

Hab. — ?

21. *Erebia Vesagus*.

♂. *Erebia Vesagus*, *Hewitson & Westwood in Gen. Diurn. Lepid.* p. 380. n. 59, pl. 64. fig. 3 (1851).

Rocky Mountains ?

Coll. Hewitson.

22. *Erebia Hispania*, sp. nov. Pl. II. fig. 7.

Alæ supra fere ut in *cassioide* sed magis elongatæ, fasciis valde distinctis, ocello geminato anticarum multo majore, pupillis oblique positus; antennarum clava distincte ochrea.

Alæ subtus velut in *Tyndaro*, ocello anticarum excepto.

Exp. alar. unc. 1½.

Spain (two specimens).

B.M.

23. *Erebia Manto*.

Papilio Manto, *Denis, Wiener Verz.* p. 169. fam. F. n. 15 (1775); *Hübner, Eur. Schmett.* i. pl. 101. figs. 512–514 (1805).

Papilio (*Hipparchia*) *Manto*, *Freyer, Neuere Beiträge*, i. p. 155. n. 156, tab. 85. figs. 1, 2 (1833).

Erebia Manto, *Dalman in Kongl. Vetensk.* p. 82. n. 11 (1816).
Satyrus Manto, *Godart, Léop. de France*, ii. p. 125. n. 47, pl. 17.
 figs. 7, 8 (1822).

Papilio (Nymph. Gemm.) *Lappona*, *Esper, Schmett.* i. pt. 2.
 tab. 108, cont. 63. fig. 3 (1777).

Papilio Zilia, *Borkhausen, Naturgesch. eur. Schmett.* ii. p. 209.
 n. 40, 41 (1789).

Switzerland. B.M.

Var. Alis majoribus.

Hungary. B.M.

Var. CASTOR.

Papilio (Nymph. Gemm.) *Castor*, *Esper, Schmett.* i. pt. 2. tab. 67.
 cont. 17. fig. 2 (1777).

Papilio Pandrosus, *Herbst, Natursyst. Schmett.* viii. p. 137. n. 75,
 tab. 202. figs. 7, 8 (1796).

Lapland. B.M.

Var. POLLUX.

Papilio (Nymph. Gemm.) *Pollux*, *Esper, Schmett.* i. pt. 2. tab. 67.
 cont. 17. fig. 3 (1777).

Papilio Aglauros, *Herbst, Natursyst. Schmett.* viii. p. 138. n. 76,
 tab. 203. figs. 1, 2 (1796).

Maniola Baucis, *Schrank, Fauna Boica*, ii. p. 177. n. 1307
 (1801).

Hab. — ? (one bad specimen). B.M.

24. *Erebia Sthennyo.*

Erebia Sthennyo, *Graslin, Ann. Ent. Soc. France*, p. 393. n. 1,
 pl. 10. figs. 1-3 (1850).

Pyrenees.

25. *Erebia Ocnus.*

Hipparchia Ocnus, *Eversmann, Bull. Soc. Imp. Nat. Mosc.*
 p. 538, tab. 8. figs. 5a, 5b (1843).

Erebia Ocnus, *Herrich-Schäffer, Eur. Schmett.* i. figs. 291, 292
 (1843-45).

Ural Mountains.

The last two species are nearly allied to but quite distinct from
Manto; neither of them are at present in the National Collection.

26. *Erebia mantoides*, sp. nov. Pl. II. fig. 6.

Alæ supra velut in *Mantone*; subtus maculis ocellaribus subapicalibus anticarum contiguus, majoribus; apicibus magis albicantibus:

posticæ coloribus *Mantonis*, lineis autem mediis forma tota distantibus, his late separatis, externa minus irregulari, angulum inter venulas secundam et tertiam medianas formante.

Exp. alar. unc. $1\frac{1}{4}$.

Lapland (two specimens).

B.M.

27. *Erebia Embla*.

Papilio Embla, *Thunberg, Diss. Ent. Nov. Ins. Suec.* pt. 2. p. 38, pl. 2. fig. 8 (1791); *Quensel, Act. Reg. Suec.* t. 9. figs. 1, 2 (1791).

Papilio (*Hipparchia*) Embla, *Freyer, Neuere Beiträge*, v. p. 55. n. 727, tab. 416. figs. 3, 4 (1845).

Satyrus Embla, *Zetterstedt, Ins. Lappon.* p. 903. n. 12 (1839).

Erebia Embla, *Dalman* (1816); *Herrich-Schäffer, Eur. Schmett.* i. figs. 382, 383 (1841).

Papilio Ethus, *Fabricius, Ent. Syst.* iii. pt. 1. p. 217. n. 680 (1793).

Var. Papilio Sophia, *Acerbi, Voy. au Cap Nord*, pl. 16. figs. 1, 2 (1804).

Var. Papilio Dioxippe, *Hübner, Eur. Schmett.* i. pl. 105. figs. 538, 539 (1805).

Satyrus Dioxippe, *Duponchel, Lép. de France*, i. p. 230. n. 72, pl. 36. figs. 1, 2 (1832).

Sweden, Russia.

B.M.

28. *Erebia Ero*.

Erebia Ero, *Bremer, Lep. Ost-Sibir. u. Amur. in Bull. Acad. Imp. Sci. St. Pétersb.* p. 468. n. 11 (1861).

Apfel Mountains.

This species seems, from the description, to come between *Embla* and *Disa*; it has the bipupillated ocellus of the former, but in other respects seems to be more nearly allied to the latter species.

29. *Erebia Disa*.

Papilio Disa, *Thunberg, Diss. Ent. Nov. Ins. Suec.* pt. 2. p. 37 (1791).

Papilio (*Hipparchia*) Disa, *Freyer, Neuere Beiträge*, v. p. 54. n. 726, tab. 416. figs. 1, 2 (1845).

Satyrus Disa, *Zetterstedt, Ins. Lappon.* p. 903. n. 13 (1839).

Erebia Disa, *Westwood in Gen. Diurn. Lepid.* p. 379 n. 31 (1851).

Papilio (*Nymph. Gemm.*) Gesion, *Esper, Schmett.* i. pt. 2. tab. 108. cont. 63. fig. 2 (1777).

Papilio Gesion, *Quensel in Act. Acad. Reg. Suec.* p. 272, pl. 9. figs. 5, 6 (1791).

Papilio Griela, *Fabricius, Ent. Syst.* iii. pt. 1. p. 236. n. 737 (1793); *Hübner, Eur. Schmett.* i. pl. 49. figs. 228, 229 (1805).

Satyrus Griela, *Godart, Enc. Méth.* ix. p. 531. n. 145 (1819).

Erebia Embla, *Boisduval, Icon. Hist. Lép.* p. 162. n. 8, pl. 32. figs. 1-3 (1832).

Satyrus Embla, *Duponchel, Lép. France*, i. p. 232. n. 73, pl. 36. figs. 3, 4 (1832).

Papilio Embla, var., *Ochsenheimer, Schmett. v. Eur.* i. p. 287. n. 58 (1807).

Var. ? *Papilio Stheno*, *Hübner, Eur. Schmett.* i. pl. 109. figs. 561, 562 (1805).

Lapland.

B.M.

This species appears to be quite distinct from *Embla*.

Var. *EDDA*.

Erebia Edda, *Ménétrics in Middendorff's Reise*, p. 58. n. 11, pl. 3. fig. 11 (1854).

Eastern Siberia.

This is evidently only a small variety of *E. Disa*.

Var. *MANCINUS*.

Erebia Mancinus, *Hewitson & Westwood in Gen. Diurn. Lepid.* p. 380. n. 58, pl. 64. fig. 2 (1851).

Rocky Mountains.

B.M.

Only a local form of *E. Disa*.

30. *Erebia Rossii*.

Hipparchia Rossii, *Curtis in Append. Ross's Narrative, N.W. Passage*, p. lxxvii, pl. A. fig. 7 (1835).

Erebia Rossii, *Westwood in Gen. Diurn. Lepid.* p. 380. n. 55 (1851).

Rocky Mountains.

B.M.

Curtis's *subhyalina* must, from the description, be a species of *Æneis*; it appears to come near *Æ. exitiosa*.

31. *Erebia Ceme*.

Papilio (Nymph. Gemm.) *Ceme*, *Esper, Schmett.* i. pt. 2. tab. 120. cont. 75. fig. 2 [nec 1] (1777).

Papilio Ceme, *Hübner, Eur. Schmett.* i. pl. 104. figs. 530, 531 [nec 532, 533] (1805).

Satyrus Ceme, *Godart, Enc. Méth.* ix. p. 537. n. 158 (1819); *Duponchel, Lép. de France*, i. p. 222, pl. 34. figs. 5, 6 (1832).

Erebia Ceme, *Westwood in Gen. Diurn. Lepid.* p. 378. n. 9 (1851).

Var. *Papilio* (Nymph. Gemm.) *Cæcilia*, *Esper, Schmett.* i. pt. 2. tab. 121. cont. 76 (1777).

Switzerland, Germany.

B.M.

32. *Erebia Medusa*.

Papilio Medusa, *Denis, Wien. Verz.* p. 167. fam. F. n. 10 (1775);
Hübner, Eur. Schmett. i. tab. 45. figs. 103, 104 (1805).

Satyrus Medusa, *Godart, Enc. Méth.* ix. p. 538. n. 161 (1819);
Lép. de France, ii. p. 110. n. 39, pl. 15. figs. 5, 6 [bad] (1822).

Erebia Medusa, *Westwood in Gen. Diurn. Lepid.* p. 378. n. 12 (1851).

Papilio Ligea, *Esper, Schmett.* i. pt. 1. tab. 7. fig. 2 (1777),
 but not of *Linnaeus*.

Var. Papilio (Hipparchia) Medusa, *Freyer, Neuere Beiträge*, i.
 p. 81. n. 75, tab. 43. fig. 1 (part)* (1833).

Var. Hipparchia Medusa, *Eversmann, Fauna Lep. Volgo-Ural.*
 p. 32. n. 22 (1844).

Erebia Hippomedusa, *Boisduval, Sp. Gén. Lép. Ind.* p. 27.
 n. 204 (1840).

? *Maniola Medea*, *Schrank, Fauna Boica*, ii. p. 173 (1801).

Germany.

B.M.

Var. Psodea.

♂. *Papilio Psodea*, *Hübner, Eur. Schmett.* i. tab. 98. fig. 497, ♂
 [nec 498, 499†] (1805).

Satyrus Psodea, *Godart, Enc. Méth.* ix. p. 537 (1819), but not
 of *Duponchel*.

♂. *Erebia Psodea*, *Herrich-Schäffer, Eur. Schmett.* figs. 165,
 166 (1843-45), but not of *Boisduval*.

Erebia Oeme, *Boisduval, Icon. Hist. Lép.* p. 174. n. 14, pl. 34.
 figs. 5-8 (1832).

Satyrus Oeme, *Duponchel, Lép. de France*, i. pl. 34. figs. 7, 8
 (1832).

Var. ♂. Papilio (Hipparchia) Oeme, *Freyer, Neuere Beiträge*, i.
 p. 57. n. 52, tab. 31. figs. 1, 2 (1833).

(♂) Polish Ukraine, Russia, (♀) Sweden.

B.M.

The sexes do not appear to differ much in this *Erebia*.

Var. ♂. Papilio (Hipparchia) Psodea, *Freyer, Neuere Beiträge*,
 ii. p. 44. n. 211, tab. 121. fig. 3 (1836).

Anticæ apice magis acuto, ocellis omnibus majoribus.

Sweden.

B.M.

Var. ?. Erebia Medusa, *Herrich-Schäffer, Eur. Schmett.* i. figs. 170
 (1843-45).

Var. EUMENIS.

Papilio (Hipparchia) Eumenis, *Freyer, Neuere Beiträge*, i. tab. 85.
 figs. 3, 4 (1833).

* The right hand figure = *Eumenis*, Freyer.

† In appearance more like *Erias*, but resembling *Medusa* on the underside.

Papilio Medusa [part.], *ibid.* tab. 43. fig. 1 (right hand figure).

Switzerland, Germany.

B.M.

This can scarcely be more than a well-marked variety of *Medusa* ♀; it has a greater number of ocelli, and the hind wings are sometimes paler below.

33. *Erebia Ceto*.

♂. *Papilio Ceto*, *Hübner, Eur. Schmett.* i. p. 34. n. 48, pl. 112. figs. 578, 579 (1805).

Papilio (*Hipparchia*) *Ceto*, *Freyer, Neuere Beiträge*, i. p. 69. n. 65, tab. 37. figs. 1, 2, ?? 3 (1833).

Satyrus Ceto, *Godart, Léop. France*, ii. p. 112. n. 40, pl. 16. figs. 1, 2 (1822).

Erebia Ceto, *Westwood in Gen. Diurn. Lepid.* p. 378. n. 10 (1851).

Papilio (*Hipparchia*) *Phorcys*, *Freyer, Neuere Beiträge*, i. p. 4, tab. 193. fig. 2 (1833).

Var. Papilio (*Hipparchia*) *Bubastis*, *ibid.* p. 71. n. 67, tab. 38. fig. 1 (1833).

Sweden.

B.M.

Var. ATRATA.

Papilio (*Nymph. Gemm.*) *atratus*, *Esper, Schmett.* i. pt. 2. tab. 104. cont. 59. fig. 1 (1777).

Switzerland.

B.M.

34. *Erebia Afer*.

Papilio (*Nymph. Gemm.*) *Afer*, *Esper, Schmett.* i. pt. 2. tab. 83. cont. 23. figs. 4, 5 (1777).

Papilio Afer, *Schneider, Syst. Besch.* p. 104. n. 44 (1787).

Papilio (*Hipparchia*) *Afer*, *Freyer, Neuere Beiträge*, i. p. 70. n. 66, tab. 37. fig. 4 (1833).

Hipparchia Afer, *Eversmann, Fauna Lepid. Volgo-Ural.* p. 33. n. 23 (1844).

Papilio Afra (*sic*), *Fabricius, Ent. Syst.* iii. pt. 1. p. 236. n. 738 (1793).

Satyrus Afra, *Godart, Enc. Méth.* ix. p. 530. n. 144 (1819).

Erebia Afra, *Boisduval, Icon. Hist. Léop.* p. 170. n. 12, pl. 34. figs. 1, 2 (1832).

Papilio Phegea, *Borkhausen, Naturgesch. eur. Schmett.* i. p. 101. n. 42 (1805), also of *Hübner*.

Var. ? Satyrus Afer, *Duponchel, Léop. de France*, i. p. 224. n. 79, pl. 35. figs. 1, 2 (1832).

Var. ? Satyrus Dalmata, *Godart, Enc. Méth.* ix. p. 530. n. 143 (1819).

Switzerland.

B.M.

35. *Erebia Parmenio*.

Papilio Parmenio, *Boeber, Mém. Soc. Imp. Nat. Mosc.* (1809).
Erebia Parmenio, *Herrich-Schäffer, Eur. Schmett.* i. figs. 421,
 422, 464, 466 (1843-45).
Satyrus Stirius, *Godart, Enc. Méth.* ix. p. 530. n. 142 (1819).
 Siberia, Russia. B.M.

36. *Erebia tristia*.

Erebia tristia, *Bremer, Lep. Ost-Sibir. u. Amur. in Bull. Acad.*
Imp. Sc. St. Pétersb. p. 467. n. 10 (1861).
Erebia Wanga, *Bremer, Lep. Ost-Sibir. u. Amur. Mém. Acad.*
Imp. Sci. St. Pétersb. viii. p. 20, pl. 2. fig. 1 (1865).
 Bureya Mountains.

Seems to be most nearly allied to *Cyclopius*; the ocellus of front wings with a grey instead of a yellow iris, and blending into the ground-colour; sometimes with a grey spot at the outer angle of the hind wings.

37. *Erebia Cyclopius*.

Erebia Cyclopius, *Herrich-Schäffer, Eur. Schmett.* i. figs. 607,
 608 (1843-45).
Hipparchia Cyclopius, *Eversmann in Bull. Soc. Imp. Nat. Mosc.*
 p. 590, tab. 14. figs. 3A, 3B (1844).
Satyrus Cyclopius, *Westwood in Gen. Diurn. Lepid.* p. 389. n. 5
 (1851).
 Irkutsk.

38. *Erebia fasciata*, sp. nov. Pl. II. fig. 8.

Alæ supra fuscae, anticæ velut in *discoidalis* rufescente.

Alæ subtus pallidiores, anticæ rufo-ferrugineæ, apice albicante, fasciis tribus subfuscis fusco-ferrugineo limitatis, prima basali confusa indistincta, secunda media lata plerumque utrinque undulata, ad costam latiore, tertia marginali: posticæ cinereo-albidæ, fasciis tribus distinctis rufo-fuscis (rarius cinereo-ferrugineis fusco limitatis), basali confusa nebulari, secunda undata et utrinque plerumque dentata, tertia marginali intus minime undulata: corpus nigrum, antennis nigris (rarius ferrugineis).

Exp. alar. unc. 2-2½.

Arctic America (9 specimens).

B.M.

There seems to be no published description exactly suited to this species; and it has therefore been necessary to describe it. It may belong to the *Manto* group of *Erebia*.

39. *Erebia* ? *discoidalis*.

Hipparchia discoidalis, *Kirby in Fauna Boreali-Amer.* p. 298.
 n. 416, pl. 3. figs. 2, 3 (1837).

Erebia discoidalis, Westwood in *Gen. Diurn. Lepid.* p. 380.
n. 41 (1851).

Arctic America.

B.M.

Genus 16. PSEUDONYMPHA, Wallengren (1857).

Typical Species *Pseudonympha Hippiæ*.

1. *Pseudonympha Hippiæ*.

Papilio Hippiæ, Cramer, *Pap. Exot.* iii. pl. 302. figs. C, D
(1782).

Erebia? *Hippiæ*, E. Doubleday, *List Lep. Brit. Mus.* i. p. 128
(1844).

Erebia Hippiæ, Westwood in *Gen. Diurn. Lepid.* p. 380. n. 46
(1851); also of *Trimen* (1866).

Pseudonympha Hippiæ, Wallengren, *Lep. Rhop. Caffror. in
Kongl. Svenska Vetensk.* p. 32. n. 1 (1857).

Papilio Hippius, Herbst, *Natursyst. Insect.* viii. p. 156, tab. 204.
figs. 5, 6 (1796).

Satyrus Cassius, ♀, Godart, *Enc. Méth.* ix. p. 526. n. 134
(1819).

Cape of Good Hope.

B.M.

2. *Pseudonympha Hyperbius*.

Papilio Hyperbius, Linnæus, *Mus. Utr. Reg.* p. 257. n. 76 (1764);
Syst. Nat. ii. p. 769. n. 130 (1766); Cramer, *Pap. Exot.* ii.
pl. 168. figs. E, F (1779); also of *Wulfen* (1786), *Fabricius*
(1787), and *Herbst* (1796).

Satyrus Hyperbius, Godart, *Enc. Méth.* ix. p. 527. n. 135
(1819).

Erebia Hyperbius, Westwood in *Gen. Diurn. Lepid.* p. 380.
n. 47 (1851); also of *Trimen* (1866).

Pseudonympha Hyperbius, Wallengren, *Lep. Rhop. Caffror. in
Kongl. Svenska Vetensk.* p. 32. n. 3 (1857).

Cape Town, Afghanistan?

B.M.

3. *Pseudonympha Sabacus*.

Erebia Sabacus, *Trimen*, *Rhop. Afric. Austral.* ii. p. 200. n. 114,
pl. 4. fig. 1 (1866).

Cape Colony, Knysna.

B.M.

I cannot bring myself to believe that the two insects figured as *Sabacus* are mere variations of one species. The position of the ocelli is quite different in the white-veined form; and the central strigæ, which appear to constitute the most constant character in the present family, are different in outline.

4. *Pseudonympha Cassius*.

Satyrus Cassius ♂, *Godart, Enc. Méth.* ix. p. 526. n. 134 (1819).

Erebia? *Cassius, E. Doubleday, List Lep. Brit. Mus.* i. p. 128 (1844).

Erebia Cassius, Westwood in Gen. Diurn. Lepid. p. 380. n. 45 (1851); also of *Trimen* (1866).

Papilio Hyperbicus ♂, *Cramer, Pap. Exot.* ii. pl. 168. figs. C, D (1779); also of *Herbst* (1796).

Pseudonympha hyperbioides, Wallengren, Lep. Rhop. Caffror. in Kongl. Svenska Vetensk. p. 20 (1857).

Knysna, Delagoa Bay, Cape of Good Hope.

B.M.

5. *Pseudonympha Natalii*.

Satyrus Natalii, Boisduval, App. Voy. de Deleg. p. 594. n. 84 (1847).

Erebia Natalii, Westwood in Gen. Diurn. Lepid. p. 380. n. 49 (1851); also of *Trimen* (1866).

Port Natal.

6. *Pseudonympha Trimenii*, sp. nov.

Erebia Sabacus, var. *a*, *Trimen, Rhop. Afric. Austral.* ii. p. 201. n. 114, pl. 4. fig. 2 (1866).

Near Cape Town (Trimen), South Africa.

B.M.

7. *Pseudonympha Tamatavæ*.

Satyrus Tamatavæ, Boisduval, Fauna Ent. de Madag. p. 60, pl. 8. figs. 6, 7 (1834).

Erebia? *Tamatavæ, E. Doubleday, List Lep. Brit. Mus.* App. p. 32 (1847).

Erebia Tamatavæ, Westwood in Gen. Diurn. Lepid. p. 380. n. 51 (1851).

Madagascar.

B.M.

8. *Pseudonympha? Panda*.

Satyrus Panda, Boisduval, App. Voy. de Deleg. p. 594. n. 85 (1847).

Erebia Panda, Westwood in Gen. Diurn. Lepid. p. 380. n. 50 (1851); *Hopffer in Peter's Reise nach Mosamb. Ins.* pl. 25. figs. 1, 2 (1862); also of *Trimen* (1866).

Amazoulu (Boisd.), Inhambane (Hpffr.), Damaraland (Trim.).

Genus 17. HOMÆONYMPHA, *Felder* (1867).Typical Species *Homœonympha Pusilla*.1. *Homœonympha pusilla*.*Homœonympha pusilla*, *Felder, Reise der Fregatte Novara*, iii.
p. 487. n. 841 (1867).Chili (*Felder*).Genus 18. TETRAPHLEBIA, *Felder* (1867).Typical Species *Tetraphlebia Germainii*.1. *Tetraphlebia Germainii*.*Tetraphlebia Germainii*, *Felder, Reise der Fregatte Novara*, iii.
p. 487. n. 842 (1867).Chili (*Felder*).2. *Tetraphlebia? plumbeola*, sp. nov. Pl. II. fig. 11.Alæ supra olivaceo-fuscæ, forma generis *Leptoneuræ*.

Anticæ subtus pallidiores, area basali, costa et margine interno exceptis, rufo-ferruginea; hac a linea angulata et biarcuata discali fusca tenui limitata; linea submarginali subintegra alteraque marginali ab atomis plumbeo-albis apud apicem interrupta fuscis; ciliis olivaceis: posticæ olivaceæ venis albicantibus; fascia media lata irregulari arcuata fusca, de costa apud basim ad angulum interno-analem currente, hac nigro limitata et squamis ochraceis utrinque marginata, punctis sex ocellaribus regulariter positis, tertio et quarto albis, aliis nigris ochraceo cinctis fusco circumcinctis et partim minutissime pupillatis; linea submarginali interrupta nigra extus plumbeo-albido marginata; ciliis olivaceis.

Exp. alar. unc. 1½.

Port Famine.

B.M.

Genus 19. FAUNULA, *Felder* (1867).Typical Species *Faunula Leucoglène*.1. *Faunula Leucoglène*.*Faunula Leucoglène*, *Felder, Reise der Fregatte Novara*, iii.
p. 488. n. 843 (1867).Chili (*Felder*).

Rather larger than *Erebia Gorge*; the anal angle and apical margin of the hind wings more strongly produced than in any hitherto known *Erebia*.

Genus 20. *STYGNUS*, *Felder* (1867).Typical Species *Stygnus humilis*.1. *Stygnus humilis*.*Stygnus humilis*, *Felder, Reise der Fregatte Novara*, iii. p. 489.
n. 844 (1867).Chili, Valdivia (*Felder*).A small species, with somewhat the habit of a *Lymanopoda*.Genus 21. *NEOSATYRUS*, *Wallengren* (1858).Typical Species *Neosatyryus Ambiorix*.1. *Neosatyryus Ambiorix*.*Neosatyryus Ambiorix*, *Wallengren in Wien. ent. Monatschr.* iv.
p. 36. n. 14 (1860).

Valparaiso.

Genus 22. *CALLEREBIA*, *Butler* (1867).Typical Species *Callerebia scanda*.1. *Callerebia scanda*.*Erebia scanda*, *Kollar in Hügel's Kaschmir*, iv. pt. 2. p. 452,
tab. 17. figs. 3, 4 (1844); also of *Westwood and Moore*.*Callerebia scanda*, *Butler, Ann. & Mag. Nat. Hist.* ser. 3.
xx. p. 217. sp. 1 (1867).Var. *a.* *Butl. Ann. & Mag.* (1867).Var. *b.* *Erebia Annada*, *F. Moore, Cat. Lep. Mus. East-Ind.*
Comp. i. p. 226. n. 475 (1857).*Callerebia Annada*, *Butl.*, and var. *b.* pl. 4. fig. 8 (1867).Var. *c.* *Butl. Ann. & Mag.* var. *c.* pl. 4. fig. 9 (1867).

Cashmere, North India.

B.M.

2. *Callerebia Nirmala*.*Erebia Nirmala*, *F. Moore, Proc. Zool. Soc. London*, p. 501.
sp. 91 (1865).*Callerebia Nirmala*, *Butler, Ann. & Mag. Nat. Hist.* ser. 3.
xx. sp. ? 3 (1867).North-western Himalayas (*Moore*).*Callerebia scanda* is exceedingly variable both in form and pattern. Some specimens have the front wings rounded, whilst others have them slightly falcate at the apex. Some have the hind wings

nearly entire, whilst others have them more strongly sinuated and produced at the anal angle into a well-marked lobe.

The ocelli vary much in number, form, and position; the commoner form (*Annada*) being represented by two subanal ocelli and three subapical white dots. The subanal ocelli are sometimes blind, and frequently placed in an oblique position. The discal white dots sometimes form the pupils of ocelli. We have one specimen in which the third dot has become an ocellus, and two in which the second and third dots are replaced by perfect ocelli: in both cases the colouring of the hind wings is uniformly brown. I have seen a specimen of *Annada*, named by Mr. Moore, in Mr. Bates's Collection.

Genus 23. CALISTO (*Hübner*), *Westwood* (1851).

Typical Species *Calisto Zangis*.

1. *Calisto Zangis*.

♀. *Papilio Zangis*, *Fabricius*, *Syst. Ent.* p. 486 (1775); also of *Herbst*.

Hipparchia Zangis, *Hübner*, *Verz. bek. Schmett.* p. 57. n. 545 (1816).

Satyrus Zangis, *Godart*, *Enc. Méth.* ix. p. 525. n. 129 (1819).

Calisto Zangis, *Hewitson & Westwood*, *Gen. Diurn. Lepid.* pl. 66. fig. 5, p. 399. n. 1 (1851).

Papilio Agnes, *Cramer*, *Pap. Exot.* iv. pl. 325. figs. A, B (1782).

West Indies.

B.M.

2. *Calisto Herophile*.

Calisto Herophile, *Hübner*, *Zuträge Exot. Schmett.* figs. 269, 270 (1806); also of *Doubleday* and *Westwood*.

Honduras.

♀, B.M.

3. *Calisto Hysius*.

Satyrus Hysius, *Godart*, *Enc. Méth.* ix. p. 525. n. 131 (1819).

Calisto Hysius, *Westwood* in *Gen. Diurn. Lepid.* p. 399. n. 3 (1851).

North America? (*Godart*), var. *St. Domingo*.

B.M.

Our specimens agree with *Godart's* description in every particular, excepting the red colouring of the outer margins. The Museum specimens only have the margins a little paler in colouring.

4. *Calisto Archebates*.

Satyrus Archebates, *Ménétriés*, *Bull. Soc. Imp. Nat. Mosc.* v. (1832); and *Nouv. Mém. Soc. Nat. Mosc.* iii. p. 38 (1834).

Euptychia Archebates, *Westwood in Gen. Diurn. Lepid.* p. 374. n. 25 (1851); and *Bull. Zool. Proc.* (1866).

Antilles, Haiti.

Coll. Salvin.

There is a copy of the 'Nouveaux Mémoires' in the Library of the Linnean Society. Mr. Salvin has also pointed out that the paper is merely a reprint of that published in 1832 in the 'Bulletin of the Imperial Society.'

Genus 24. ZIPAETIS, *Hewitson* (1863).

Typical Species *Zipaetis Saitis*.

1. *Zipaetis Saitis*.

Zipaetis Saitis, *Hewitson, Exot. Butterf.* p. 100, pl. 50. fig. 6 (1863).

Neilgherries (Hewitson).

B.M.

2. *Zipaetis Scylax*.

Zipaetis Scylax, *Hewitson, Exot. Butterf.* p. 100, pl. 50. fig. 7 (1863).

Sylhet (Hewitson).

B.M.

Genus 25. PARAMECERA, gen. nov.

Alis omnibus supra dense pilosis: anticis costa convexa angulari; margine externo leviter sinuato; margine interno paulum undulato; vena prima disco-cellulari concava, brevi; secunda longiore subobliqua; vena mediana angulariter irregulari; venis ad basin bene tumidis: posticis costa convexiuscula, margine externo bene sinuato, margine interno ad angulum ani valde excavato; venis velut in *Euptychia*, magis autem regularibus.

Corpus lanare, breve; caput oculis cirratis, palpis magnis valde pilosis, articulo apicali angulariter projecto; antennis brevibus, clava subpiriformi, subtus excavata.

Coloribus generis *Ameccera*, Butl., cellis autem discoidalibus magis ad illas *Euptychia*, Hübn., approximantibus.

This genus bears a great general resemblance to the genus *Ameccera*. It, however, differs entirely in the form of the discoidal cells, the form of the hind wings, and the palpi; these are somewhat similar to the palpi of *Mycalesis*, of which a figure is given in the 'Annals and Magazine of Natural History,' Jan. 1867. It is allied to *Euptychia*, but the hind-wing cell is more regular, which brings the genus nearer to *Heteronympha*. The antennæ also are shorter, and more like those of *Ameccera*; and the densely hairy upper surface of the wings separates it from all the rest of the *Satyridæ*.

1. *Paramecera Xicaque* (*Neonympha Xicaque*, Reakirt, antè, p. 39).
Pl. II. fig. 5.

♂. Alis supra griseo-flavidis, area basali olivaceo tinctis: anticæ costa obscuriore, fascia discali interna cinerea obliqua cellam cingente; margine externo fusco; ocellis duobus subapicalibus cæcis nigris, secundo multo minore: posticæ area anali flavescente, margine apicali fusco, anali ferrugineo, bilineatis; ocellis quinque discalibus cæcis nigris, secundo et quarto majoribus: corpus cinereo-fuscum; antennis albo annulatis, præ flavescentibus.

Alæ subtus clariore: anticæ flavæ, area basali obscuriore ad basin fusca; fasciis duabus mediis ferrugineis, externa apud angulum analem angulis alternis irregulari; lineis duabus marginalibus fusco-ferrugineis, interna undata, externa subintegra; margine ipso nigro; ciliis albo variegatis; ocellis duobus subapicalibus nigris, flavo iridatis, albo pupillatis et fusco circumcinctis, primo subgeminato, secundo multo minore: posticæ area basali fusca, albedo rorata; area apicali rosea; fascia lata media valde irregulari fusca, tota atomis ochreis rorata, ferrugineo utrinque marginata; lineis duabus marginalibus undulatis, interna ad apicem nigrescente et valde sinuata, ferrugineis; ocellis sex discalibus nigris, albo pupillatis, flavo cinctis et fusco circumcinctis; ciliis fuscis albedo variis: corpus fusco-cinereum.

Exp. alar. unc. $1\frac{1}{4}$.

Mexico.

♂, B.M.

Genus 26. *IDIONEURA*, *Felder* (1867).

Typical Species *Idioneura erebioides*.

1. *Idioneura erebioides*.

Idioneura erebioides, *Felder*, *Reise der Fregatte Novara*, iii. p. 474. n. 813 (1867).

New Granada (*Felder*).

The size of *Erebia Epiphron*, which it resembles in the form of its wings. Dr. *Felder* places this insect near the genus *Lymanopoda*, but states that the form of the hind-wing cell is similar to that of *Heteronympha Merope*.

Genus 27. *HETERONYMPHA* (part.), *Wallengren* (1858).

Typical Species *Heteronympha Merope* *.

1. *Heteronympha Merope*.

♀. *Papilio Merope*, *Fabricius*, *Ent. Syst.* iii. pt.1. p. 99. n. 306 (1793); *Donovan*, *Ins. New Holland*, pl. 28. fig. 2 (1805).

* As this is the type of the genus *Hipparchioides*, the latter will have to be rejected, though *Wallengren*'s description will not suit the genus as it now stands.

- Satyrus Merope*, *Godart, Enc. Méth.* ix. p. 500. n. 80 (1819);
also of *Boisduval* (1832-35).
Lasiommata Merope, *E. Doubleday, List Lep. Brit. Mus.* i. p. 134
(1844); also of *Westwood* (1851).
Heteronympha Merope, *Wallengren, Nov. Gen. Lepid. Kongl.*
Wet. Akad. Forhandl. p. 78 (1858).
Hipparchioides Merope, *Butler, Ann. & Mag. Nat. Hist.* xix.
p. 126. n. 1 (1867).
♂ ♀. *Oreas nubila* *Cenomais*, *Hübner, Exot. Schmett.* i. pl. 94.
figs. 1-4 (1806).
Minois *Cenomais*, *Hübner, Verz. bek. Schmett.* p. 58. n. 552
(1816).
♂. *Satyrus Archemor*, *Godart, Enc. Méth.* ix. p. 500. n. 82
(1819).
Australia. ♂ ♀, B.M.

Var. Minor et obscurior.

Tasmania.

♂ ♀, B.M.

2. *Heteronympha Duboulayi*.

Hipparchioides Duboulayi, *Butler, Ann. & Mag. Nat. Hist.*
(March 1867).

Champion Bay.

B.M.

3. *Heteronympha Philerope*.

♂. *Satyrus Philerope* [part.], *Boisduval, Voy. de l'Astrolabe,*
Entom. pt. 1. p. 147 (1832-35); also of *Guérin, Voy. de la*
Favorite, Suppl. pl. 3. fig. 2, and p. 16 (1839).

♂. *Lasiommata Philerope* [part.], *Westwood in Gen. Diurn.*
Lepid. p. 387. n. 19 (1851).

♀. *Hipparchioides Philerope*, *Butler, Ann. & Mag. Nat. Hist.*
xix. p. 126. n. 2, pl. 3. fig. 2 (1867).

Australia.

♂, Coll. Hewitson; ♀, B.M.

4. *Heteronympha Banksii*.

Hipparchia Banksii, *Leach, Zool. Miscell.* i. p. 10 (1814).

Lasiommata Banksia, *E. Doubleday, List Lep. Brit. Mus.* i. p. 134
(1847); also of *Westwood* (1851).

Hipparchioides Banksia, *Butler, Ann. & Mag. Nat. Hist.* xix.
p. 126. n. 3 (1867).

Satyrus Gelanor, *Godart, Enc. Méth.* ix. p. 498. n. 73 (1819);
also of *Boisduval* (1832-35), but not *P. Gelanor*, *Cramer*.

New Holland.

♂, B.M.

5. *Heteronympha mirifica*.

♀. *Lasiommata mirifica*, *Butler, Ann. & Mag. Nat. Hist.* xvii.
p. 286 (1866).

Hipparchioides mirifica, *Bull. ibid.* xix. p. 127. n. 4, pl. 3. fig. 1 (1867).

Hab. — ?

♀, B.M.

Genus 28. *CALLITÆRA*, gen. nov.*

1. *Callitæra Menander*.

Papilio Menander, *Drury, Ill.* iii. pl. 38. fig. 3 (1782).

Papilio Andromeda, *Fabricius, Ent. Syst.* (nec *Syst. Ent.*) iii. pt. 1. p. 184. n. 569 (1793).

Satyrus Andromeda, *Godart, Enc. Méth.* ix. p. 485. n. 18 (1819).

Hætera Andromeda (part.), *Westwood in Gen. Diurn. Lepid.* p. 365. n. 2 (1851); *Butler, Proc. Zool. Soc.* pl. 3. fig. 2 (1866).

Panama.

B.M.

2. *Callitæra Aurora*.

Hætera Pireta, *Cram.?*, var. *Aurora*, *Felder, Wien. ent. Monatschr.* vi. p. 126. n. 149 (1862).

Hætera Aurora, *Butler, Proc. Zool. Soc.* p. 42. pl. 3. figs. 7, 8 (1866).

Bogota, Ega, Para, Peru.

B.M.

There are several undescribed species of this group in the collections of Messrs. Salvin and Bates, and one or two in the National Collection.

3. *Callitæra Esmeralda*.

Hætera Esmeralda, *E. Doubleday, Ann. & Mag. Nat. Hist.* xvi. p. 306 (1845); *Hewitson & Westwood, Gen. Diurn. Lepid.* p. 365. n. 3 (1851).

Para.

B.M.

4. *Callitæra Harpalyce*.

Hætera Harpalyce, *Butler, Proc. Zool. Soc.* p. 42. pl. 3. fig. 6 (1866).

Upper Amazons.

B.M.

5. *Callitæra Andromeda*.

Papilio Andromeda, *Fabricius, Syst. Ent.* (nec *Ent. Syst.*) p. 467. n. 107 (1775).

Cithærias Andromeda, *Hübner, Verz. bek. Schmett.* p. 53. n. 496 (1816).

♂. *Papilio Philis*, *Cramer, Pap. Exot.* iv. pl. 387. fig. E (1782).

Cithærias Cissa, *Hübner, Verz. bek. Schmett.* p. 53. n. 495 (1816).

♂. *Hætera pellucida*, *Butler, Proc. Zool. Soc.* p. 41. pl. 3. fig. 5 (1866).

Surinam (Cramer), Cayenne (Butler).

B.M.

* *Alis subpiriformibus, hyalinis; venis omnibus regularibus, bene separatis; cella anticarum perlonga, posticarum ei Heteronymphe simillima.*

Genus 29. *HÆTERA* (sic), *Fabricius* (1807).Typical Species *Hætera* *Piera*.1. *Hætera* *Piera*.

Papilio *Piera*, *Linnaeus*, *Mus. Lud. Utr.* p. 220. n. 39 (1764);
Syst. Nat. p. 754. n. 52 (1766); *Clerck*, *Icones*, tab. 36. fig.
 43 (1759-64).

Oreas dubia *Piera*, *Hübner*, *Samml. exot. Schmett.* i. pl. 82.
 figs. 1-4 (1806).

Cithærias *Piera*, *Hübner*, *Verz. bek. Schmett.* p. 53. n. 494
 (1816).

Satyrus *Piera*, *Godart, Enc. Méth.* ix. p. 483. n. 17 (1819).

Hætera *Piera*, *Westwood in Gen. Diurn. Lepid.* p. 365. n. 1
 (1851).

Papilio *Piera* (sic), *Sulzer, Gesch. Ins.* tab. 15. fig. 5.

♀. *Papilio Anacardii*, *Fabricius, Syst. Ent.* p. 467. n. 105
 (1775); and (part.), *Linnaeus, Mus. Lud. Utr.* p. 236. n. 55
 (1764); *Merian, Ins. Surin.* t. 16. fig. sup. sinistr. (1705).

Papilio *Piera*, *Cramer, Pap. Exot.* iv. pl. 291. figs. C-E (1782).
Surinam (*Merian, Cramer*), *Brazil* ? B.M.

Linnaeus quotes *Madame Merian's* figure of this insect as a representation of *Salamis Anacardii*; *Fabricius*, in the 'Systema Entomologiæ,' not only repeats the error, but alters the description to suit *Merian's* representation: they both quote *Clerck's* figure, which well illustrates the *Linnaean* species.

Para, *Peru*.

B.M.

Local form. ? *H. PIERA* of *Boisduval*.

Hætera *Piera*, *Boisduval, Sp. Gén. Lép.* pl. 13 (9 B). fig. 4
 (1836).

Alæ posticæ fascia integra haud fulvo tincta.

Brazil, *Ega*.

B.M.

Our copy of *Boisduval's* work is uncoloured, but his figure has quite the character of this form.

Local form. *H. DIAPHANA*.

Hætera diaphana, *Lucas, in Ramon de la Sagra's Hist. de l'Île de Cuba (Insectes)*, p. 587 (1857).

Cuba (*Lucas*).

This is probably identical with the preceding form from *Brazil* and *Ega*; at any rate the description is applicable to it.

Local form. *H. HYMENÆA*.

Hætera hymenæa, *Felder, Reise der Fregatte Novara*, iii. p. 459.
 n. 771 (1867).

Bahia (*Felder*).

The description of this form agrees with that of *diaphana*; and it

is more than probable that the two are identical, although from different localities.

Var. NEGRA.

Hætera Piera, *var. Negra*, *Felder, Wien. ent. Monatsch.* vi. p. 127. n. 150 (1862).

Rio Negro (Felder).

2. *Hætera Macleannania.*

Hætera Macleannania, *Bates, Ent. Month. Mag.* iii. p. 180 (1866).

Panama.

Colls. Salvin and B.M.

We have a species from Bogota and Para, intermediate between this and the typical form of *Piera*, which will probably be shortly described by Mr. Salvin.

3. *Hætera Hypæsia.*

Hætera Hypæsia, *Hewitson, Trans. Ent. Soc. London*, 2nd ser. ii. p. 247, pl. 23.

Bogota.

B.M.

This species varies slightly as regards the position of the discocellular veins of the hind wings.

Genus 30. *PIERELLA*, *Westwood* (1851).

Typical Species *Pierella Nereis*.

1. *Pierella Nereis.*

Papilio Nereis, *Drury, Ill.* iii. pl. 35. figs. 2, 3 (1782).

Cithærias Nereis, *Hübner, Verz. bek. Schmett.* p. 53. n. 497 (1816).

Satyrus Nereis, *Godart, Enc. Méth.* ix. p. 483. n. 16 (1819);

Lucas, Lep. Exot. p. 146, pl. 8 A. fig. 1 (1835).

Hætera (Pierella) Nereis, *Westwood in Gen. Diurn. Lepid.* p. 365. n. 4 (1851).;

Brazil &c.

B.M.

2. *Pierella Hyceta.*

Hætera Hyceta, *Hewitson, Exot. Butterf.* ii. p. 83. pl. 42. fig. 1 (1859).

Nauta (Amazons).

Type, B.M.

3. *Pierella Latona.*

Hætera Latona, *Felder, Reise der Fregatte Novara*, iii. p. 459. n. 770 (1867).

Bogota (Felder).

According to Felder, this species would be intermediate between *Hyceta* and *Lamia*.

4. *Pierella Rhea*.

Papilio Rhea, *Fabricius, Syst. Ent.* p. 467. n. 106 (1775).

Pieris Rhea, *Hübner, Verz. bek. Schmett.* p. 53. n. 498 (1816).

Demerara, Rio Joneiro.

B.M.

Subsp. P. LAMIA.

Papilio (Nymph. Phal.) *Lamia*, *Sulzer, Gesch. Ins.* p. 145. tab. 18. fig. 1 (1776).

Hætera Lamia, *Felder in Wien. ent. Monatschr.* vi. p. 126. n. 148 (1862).

Papilio Dindymene, *Cramer, Pap. Exot.* iii. tab. 198. figs. F, G (1782).

Satyrus Dindymene, *Godart, Enc. Méth.* ix. p. 482. n. 14 (1819).

Hætera Dindymene, *Westwood in Gen. Diurn. Lepid.* p. 365. n. 5 (1851).

Bogota, Peru, and (var.) Para.

B.M.

5. *Pierella Luna*.

Papilio Luna, *Fabricius, Ent. Syst.* iii. pt. 1. p. 109. n. 336 (1793).

Satyrus Luna, *Godart, Enc. Méth.* ix. p. 482. n. 15 (1819).

Hætera Luna, *Westwood in Gen. Diurn. Lepid.* p. 365. n. 6 (1851).

Demerara.

B.M.

Subsp. H. Luna, *Hewitson, Exot. Butterf.* ii. p. 83, pl. 42. fig. 3 (1859).

Nicaragua.

B.M.

This is a very constant and well-marked form, and will probably be named in Mr. Salvin's forthcoming monograph of this group of Butterflies.

6. *Pierella Astyoche*.

Hætera (sic) *Astyoche*, *Erichson in Schomburgk's Reisen in Britisch-Guiana*, Theil iii. p. 599 (1848).

Hætera Larymna, *Hewitson and Westwood in Gen. Diurn. Lepid.* p. 365. n. 7, pl. 62. fig. 1 (1851).

British Guiana and Para.

B.M.

Var. Hætera Astyoche, var., *Hewitson, Exot. Butterf.* ii. p. 84, pl. 42. fig. 5 (1859).

Peru, Nauta, Para.

B.M.

7. *Pierella ocreata*.*Pierella ocreata* (species hand descripta).

Panama.

Colls. Salvin and B.M.

This beautiful species will most likely be shortly described by Mr. Salvin; it is allied to *helvina* of Hewitson.

8. *Pierella helvina*.

Hætera helvina, *Hewitson, Exot. Butterf.* ii. p. 84, pl. 42. fig. 4 (1859).

New Granada.

B.M.

There are several beautiful species allied to *helvina* in Mr. Salvin's Collection.

9. *Pierella Lena*.

Papilio Lena, *Linnaeus, Syst. Nat.* ii. p. 784. n. 206 (1766); figured by *Rösel, Ins. Belust.* Theil iv. tab. 10. figs. 3, 4 (1761).

Pieris Lena, *Hübner, Verz. bek. Schmett.* p. 53. n. 499 (1816).

Satyrus Lena, *Godart, Enc. Méth.* ix. p. 481. n. 13 (1819); but not of *Lucas*.

Hætera Lena, *Westwood in Gen. Diurn. Lepid.* p. 365. n. 8 (1851).

Para, Demerara, Rio Janeiro.

B.M.

Var. Hætera Lena, var., *Hewitson, Exot. Butterf.* ii. p. 83, pl. 42 fig. 2 (1859).

Rio Janeiro, Peru.

B.M.

Var. Hætera Lena, var. *Braziliensis*, *Felder, Wien. ent. Monatschr.* vi. p. 126. n. 147 (1862).

Rio Negro (Felder).

10. *Pierella Dracontis*.

Pieris Dracontis, *Hübner, Verz. bek. Schmett.* p. 53. n. 500 (1806).

Hætera Dracontis, *Westwood in Gen. Diurn. Lepid.* p. 365. n. 9 (1851).

Papilio Lena, *Cramer, Pap. Exot.* iv. tab. 291. figs. A, B (1782).

Oreas dubia Lena, *Hübner, Exot. Schmett.* Band i. tab. 83. figs. 1-4 (1806).

Hætera Lena, *Hewitson in Gen. Diurn. Lepid.* pl. 62. fig. 2 (1851).

Para.

B.M.

Var. Satyrus Lena, *Lucas, Lep. Exot.* pl. 79. fig. 3 (1835).

Santa Lucia.

B.M.

11. *Pierella Hortona*.

Hætera Hortona, *Hewitson, Trans. Ent. Soc. London*, ii. p. 246,
new ser. pl. 23. fig. 1.

Ega (Amazons). B.M.

Local form? Forma fasciisque diversa."

Nauta. B.M.

12. *Pierella Hortensia*.

Hætera Hortensia, *Felder, Wien. ent. Monatschr.* Band vi. p. 125.
n. 145 (1862).

Rio Negro (Felder).

There are still one or two races of the *Hortona* group undescribed, and which seem almost sufficiently distinct to constitute species.

Genus 31. ANCHIPHLEBIA, gen. nov.

Typical Species *Anchiphlebia Archæa*.

Alæ forma elongatæ: anticæ costa regulariter convexa; apice convexo, rarius acuminato; margine externo obliquo undato, margine interno velut in *Euplicea*, maribus convexo, fœminibus subrecto, maribus subtus plaga pectinatim cirrata: posticæ costa maribus valde convexa, margine externo subintegro obliquo, angulo anali subquadrato; margine interno paulum convexo.

Alæ venis anticarum superiore disco-cellulari valde obliqua, inferiore subconvexa; posticarum prima et secunda subcostalibus ad origines mari valde approximatis et subparallelis; superiore disco-cellulari elongata paulum concava; inferiore brevior obliqua: corpus lanare, oculis subnudis, antennis tenuibus, palpis vix porrectis.

1. *Anchiphlebia Archæa*. (♂ neuration, Pl. V. fig. 3.)

Antirrhæa Archæa, *Hübner, Samml. exot. Schmett.* ii. pl. 81.
figs. 3, 4 (1806); also of *E. Doubleday* (1844).

Hætera (Antirrhæa) Archæa, *Westwood in Gen. Diurn. Lepid.*
p. 366. n. 12 (1851).

Satyrus Girondius, *Godart, Enc. Méth.* ix. p. 484. n. 19 (1819).

(♂) Bahia, (♂ ♀) Brazil. B.M.

2. *Anchiphlebia Hela*.

Antirrhæa Hela, *Felder, Wien. ent. Monatschr.* vi. p. 173. n. 144
(1862); *Nov. Voy.* iii. tab. 66. figs. 5, 6 (1867).

Rio Negro (Felder), Peru, St. Paulo. B.M.

Satyrus Betro of Godart, placed by Westwood in this group, is a species of *Taygetis*.

3. *Anchiphlebia Taygetina*, sp. nov. Pl. V. fig. 2.

Alæ supra ochraceæ, area basali obscuriore et a linea discali obliqua fusca limitata; margine externo fuscescente: anticæ ocellis, tribus ovalibus nigris, apicali minore, albo pupillatis: posticæ area anali obscurius fuscescente; maculis duabus vel quatuor violaceis albo punctatis discalibus; margine interno fusco: corpus olivaceo-fuscum.

Alæ subtus pallidius ochraceæ, aliquando roseo tinctæ; lineis duabus mediis, ad costam anticarum divergentibus, fuscis; externa albido extus marginata; litura interjecta anticis discoidali; linea submarginali undata indistincta fusca; punctis alis ambabus quinque discalibus serie angulari positis; margine externo paulum fuscescente: corpus ochreo-albidum.

Exp. alar. unc. $2\frac{1}{2}$ –3.

Ega.

♂ ♀, B.M.

Genus 32. ANTIRRHÆA*, Westwood (1851).

Typical Species *Antirrhæa Philoctetes*.1. *Antirrhæa Philoctetes*.

♀. *Papilio Philoctetes*, Clerck, *Icones*, tab. 30. fig. 3 (1759); also of *Linnaeus* (1766), *Fabricius* (1775), &c.; ♂ ♀, *Cramer*, *Pap. Exot.* i. pl. 20. figs. A–C (1779).

Satyrus Philoctetes, *Godart*, *Enc. Méth.* ix. p. 480. n. 11 (1819). *Antirrhæa Philoctetes*, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 121 (1844).

Hætera (*Antirrhæa*) *Philoctetes*, *Westwood in Gen. Diurn. Lepid.* p. 365. n. 10 (1851).

Pieris Philoctetes, *Hübner*, *Verz. bek. Schmett.* p. 53. n. 501 (1816).

♂. *Broken?* *Papilio Philoctetes*, *Drury*, *Ill.* ii. pl. 1.

Demerara, Surinam, Tapajos.

♂ ♀, B.M.

2. *Antirrhæa Morna*.

Papilio Morna, *Fabricius*, *Syst. Ent.* p. 500. n. 245 (1775).

Satyrus Morna, *Godart*, *Enc. Méth.* ix. p. 481. n. 12 (1819).

Hætera (*Antirrhæa*) *Morna*, *Westwood in Gen. Diurn. Lepid.* p. 366. n. 11 (1851).

India (*Fabricius*), West Indies (*Westwood*).

Professor Westwood suggests that this is very likely a mutilated specimen of *Philoctetes*, in which case it would be represented well by *Drury's* figure, *Ill.* ii. pl. 1.

* *Hübner's Antirrhæa* was only used in the 'Sammlung,' not in his 'Verzeichniss,' and appears not to have been characterized.

3. *Antirrhaea Philaretos* (*Philopœmen*, var.?).

Antirrhaea Philaretos, *Felder*, *Wien. ent. Monatschr.* vi. n. 136 (1862).

Bogota.

B.M.

4. *Antirrhaea Philopœmen*.

Antirrhaea Philopœmen, *Felder*, *Wien. ent. Monatschr.* vi. n. 137 (1862); *Nov. Voy.* iii. p. 459. n. 766, tab. 66. figs. 3, 4 (Sept. 1867).

Bogota (Felder), Ega, St. Paulo.

♂ ♀, B.M.

5. *Antirrhaea Geryon*.

Antirrhaea Geryon, *Felder*, *Wien. ent. Monatschr.* vi. n. 139 (1862); *Nov. Voy.* iii. p. 459. n. 768, tab. 67. figs. 1, 2 (Sept. 1867).

Bogota (Felder).

B.M.

6. *Antirrhaea Lindigii*.

Antirrhaea Lindigii, *Felder*, *Wien. ent. Monatschr.* vi. n. 138 (1862); *Nov. Voy.* iii. p. 459. n. 767, tab. 66. figs. 1, 2 (1867).
Antirrhaea casta, *Bates*, *Ent. Month. Mag.* i. p. 179. n. 73 (1864).

Bogota (Felder), Guatemala [Vera Paz] (Bates). Coll. Salvin.

7. *Antirrhaea* ? *Miltiades*.

Papilio Miltiades, *Fabricius*, *Ent. Syst.* iii. pt. 1. p. 66. n. 205 (1793).

Satyrus Miltiades, *Godart*, *Enc. Méth.* ix. p. 358. n. 26 (1819).

Nymphalis ? *Miltiades*, *Westwood in Gen. Diurn. Lepid.* p. 310. n. 43 (1850).

Hætera ?? (*Antirrhaea*) *Miltiades*, *Westwood*, *ibid.* p. 366. n. 14 (1851).

Charaxes ? *Miltiades*, *Butler*, *Proc. Zool. Soc.* p. 639. n. 68 (1865).

Hab. — ?

I have compared the description of this species with most of the tailed Diurnal Lepidoptera, but I can find none to suit it.

Genus 33. ZOPHOËSSA, *Westwood* (1851).

Typical Species *Zophoëssa Sura*.

1. *Zophoëssa Sura*.

Zophoëssa Sura, *Hewitson & Westwood in Gen. Diurn. Lepid.* p. 362. n. 1, pl. 61. fig. 1 (1851).

Sylhet.

B.M.

2. *Zophoëssa Yama*.

Zophoëssa Yama, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*
i. p. 221. n. 458 (1851).

Bootan, North India (Moore), Sylhet.

B.M.

3. *Zophoëssa Goalpara*.

Zophoëssa Goalpara, *F. Moore, Proc. Zool. Soc. London*, p. 768
(1865).

? North-east Bengal (Moore).

4. *Zophoëssa Baladeva*.

Zophoëssa Baladeva, *F. Moore, Proc. Zool. Soc. London*, p. 769,
pl. 61. fig. 5 (1865).

North-east Bengal (Moore).

Genus 34. *ZETHERA* *, *Felder* (1861).

Typical Species *Zethera Pimplea*.

1. *Zethera Pimplea*.

Cynthia Pimplea, *Erichson in Meyden's Reise um die Erde*,
Theil iii. p. 405 (280). n. 5, tab. 1. fig. 5 (1834).

Diadema ? *Pimplea*, *Westwood in Gen. Diurn. Lepid.* p. 281.
n. 23 (1850).

Zethera Pimplea, *Felder in Nov. Act. Acad. Cæsar. Leop. Carol.*
Germ. Nat. Curios. 3rd Dec. tom. viii. p. 26. n. 61 (June
1861).

Luzon (Philippines).

B.M.

Var. *AGANIPPE*.

Zethera Aganippe, *Felder, Wien. ent. Monatschr.* vii. n. 71
(1863).

Bourias, Locban (Felder).

2. *Zethera Parnassia*.

Zethera Parnassia, *Felder, Wien. ent. Monatschr.* vii. n. 70
(1863). ? *Aberr. Pimplea* (*Feld.*).

Luzon (Felder).

* *Felder's* genus *Zethera* was published one month earlier than the *Amechania* of *Hewitson*; I attach no importance to the fact of its having been previously used as a MS. name; on the contrary, I should prefer to reject it on that account, as there can be no doubt that MS. names are the plague of students in entomology.

3. *Zethera Musa*.

Zethera Musa, *Felder, Wien. ent. Monatschr.* v. p. 301. n. 15 (1861).

Luzon, Mindanao (Felder), Philippines.

B.M.

4. *Zethera incerta*.

Amecania incerta, *Hewitson, Exot. Butterf.* ii. p. 87, pl. 44 (1861).

Tondano (Celebes).

Coll. Wallace.

5. *Zethera hestioides*.

Zethera hestioides, *Felder, Wien. ent. Monatschr.* v. p. 302. n. 17 (1861).

Mindanao (Felder), Philippines.

B.M.

This species is most nearly allied to the genus *Orinoma* as regards pattern; most of the markings on *O. Damaris* can be seen in a modified form upon *Z. hestioides*.

Genus 35. *ORINOMA**, *Westwood* (1851).

Typical Species *Orinoma Damaris*.

1. *Orinoma Damaris*.

Orinoma Damaris, *G. R. Gray, Lep. of Nepal*, p. 14, tab. 7. figs. 2, 2 a (1846); *Hewitson in Gen. Diurn. Lepid.* p. 369. n. 1, pl. 63. fig. 3 (1851); *Butler, Ann. & Mag. Nat. Hist.* 3rd ser. xix. pl. 2. figs. 4, 4 a [structure] (1867).

Satyrus? *Orinoma*, *E. Doubleday, List Lep. Brit. Mus.* i. p. 122 (1844).

Nepal, North India.

B.M.

Genus 36. *ANADEBIS*, *Butler* (1867).

Typical Species *Anadebis Himachala*.

1. *Anadebis Himachala*.

Theope (*Mycalasis*?) *Himachala*, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i. p. 234 (1857).

Ethepe Himachala, *F. Moore, Proc. Zool. Soc. London*, p. 770 (1865).

* The MS. generic name *Orinoma* was applied first by Mr. E. Doubleday, and adopted by Mr. G. R. Gray in his 'Insects of Nepal'; he also uses the generic name *Pieris*. This genus only differs from *Zethera* in the relative length of the disco-cellular nervelets.

Anadebis Himachala, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 51,
pl. 2. figs. 1, 1a, 1b (1867).

Neorina sita, *Felder, Wien. ent. Monatschr.* iii. p. 403 (1859).

Darjeeling (Moore), Sylhet.

B.M.

Genus 37. *NEORINA*, *Westwood* (1851).

Typical Species *Neorina Hilda*.

1. *Neorina Hilda*.

Neorina Hilda, *Hewitson & Westwood in Gen. Diurn. Lepid.*
p. 370. n. 1 (1851); also of *Moore* (1857).

Darjeeling (Moore), East Indies.

B.M.

2. *Neorina Crishna*.

Cylo Crishna, *Westwood in Gen. Diurn. Lepid.* p. 361. n. 9
(1851); also of *Moore* (1857).

Java (Moore), North India.

B.M.

3. *Neorina Lowii*.

Cylo Lowii, *Hewitson & Westwood in Gen. Diurn. Lepid.* p. 361.
n. 8, pl. 61. fig. 4 (1851).

Borneo, Sumatra.

B.M.

Genus 38. "COELITES", *Westwood* (1851).

Typical Species *Cœlites Nothis*.

1. *Cœlites Nothis*.

Cœlites Nothis, *Hewitson & Westwood in Gen. Diurn. Lepid.*
p. 368. n. 1, pl. 66. fig. 2 (1851).

East India (Westwood).

2. *Cœlites Epiminthia*.

Cœlites Epiminthia, *Westwood in Gen. Diurn. Lepid.* p. 368. n. 2
(1851); also of *Moore* (1857).

Borneo (Westwood), Sumatra, and (var.) Borneo.

B.M.

3. *Cœlites vicinus*.

Cœlites vicinus, *Felder, Reise der Fregatte Novara*, iii. p. 499.
n. 865 (1867).

Celebes (Felder).

Dr. Felder thinks this may be a local form of *Epiminthia*; his description, however, seems to indicate a very distinct species. In

Epiminthia there are five ocelli in the front wings, the second and fifth largest, the third smallest; in the hind wings there are also five, the first and last larger. None of the ocelli could, strictly speaking, be described as black; they are olivaceous, with a yellow iris edged outwardly with dark brown, and they have a diagonal white pupil irrorated with black atoms; these ocelli, moreover, all stand upon a pale violet ground,—which, I should think, can scarcely answer to the description “*cincturis confluentibus, in ocellis quatuor inferioribus extus dilatatis.*”

4. *Coelites euptychioides*.

Coelites euptychioides, *Felder, Reise der Fregatte Novara*, iii. p. 499. n. 866 (1867).

Borneo (Felder).

B.M.

According to Felder, “Agreeing with *Epiminthia* (Westw.) in the form of its wings; in the colouring of the upperside resembling the species of *Neonympha* of the group *Euptychia*, especially *Chloris* of Cramer.

“The wings are shorter than in *C. vicinus*, and the apical area broader.”

Var. HUMILIS.

Coelites humilis, *Butler, Ann. & Mag. Nat. Hist.* ser. 3. xx. p. 403, pl. 8. figs. 5, 6, pl. 9. fig. 2 (1867).

Malacca.

Coll. Roberts.

A local form of Felder's *C. euptychioides*.

Genus 39. NEOPE, *Butler* (1867).

Typical Species *Neope Bhadra*.

1. *Neope Bhadra*.

Lasiommata? (*Enope*) *Bhadra*, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i. p. 227. n. 478 (1857).

Neope Bhadra, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 166. n. 1 (1867).

Darjeeling, East Indies.

B.M.

2. *Neope Pulaha*.

Lasiommata? (*Enope*) *Pulaha*, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i. p. 227. n. 477 (1857).

Enope Pulaha, *F. Moore, Proc. Zool. Soc.* p. 499. n. 80 (1865).

Bootan, India.

B.M.

3. Neope dendrophilus.

Satyrus dendrophilus, *Trimen, Tr. Ent. Soc. Lond.* 3rd ser. i. p. 399 (1862).

Debis dendrophilus, *Trimen, Rhop. Afric. Austral.* ii. p. 191. n. 108, pl. 3. fig. 8 (1866).

Bashee River, Kaffraria (Trimen).

I have seen specimens of this species in Mr. Hewitson's Collection. It evidently belongs to the genus *Neope*.

4. Neope Moorei.

Neope Moorei, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 166. n. 3, pl. 4. fig. 7 (1867).

East Indies.

♀, B.M.

This is evidently a local representative of *N. Goschkevitschii*.

5. Neope Goschkevitschii.

Lasiommata Goschkevitschii, *Ménétriés, Enum. Corp. Anim. Lepid.* pt. 2. p. 121. n. 1298, tab. 10. fig. 4 (1857). *L. Goschkevitschii* (*sic*), *Felder, Wien. ent. Monats.* vi. p. 28 (1862).

Neope Pulaha, *Butler, Proc. Linn. Soc.* ix. p. 56. n. 21 (1866).

Neope Japonica, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 167. n. 4. (1867).

Japan, Hakodadi.

♂ ♀, B.M.

6. Neope Bremeri.

Lasiommata Bremeri, *Felder, Wien. ent. Monatschr.* vi. p. 28 (1862).

Near Ning-po.

Allied to *Goschkevitschii*.

7. Neope Muirheadii.

Lasiommata Muirheadii, *Felder, Wien. ent. Monatschr.* vi. p. 28 (1862).

Tse-Kiang (Felder), Shanghai.

B.M.

Allied to the preceding species, but with wider and more delicate wings; the front wings with the apex less produced, and the discoidal cell not reaching beyond the middle of the costa.

Var. SEGONAX.

Debis Segonax, *Hewitson, Exot. Butterf.* iii. p. 74, pl. 37. fig. 5 (1862).

China (Hewitson).

Genus 40. LETHE, *Hübner* (1816).Typical Species *Lethe Europa*.1. *Lethe Portlandia*.

Papilio Portlandia, *Fabricius, Ent. Syst.* iii. pt. 1. p. 103. n. 319 (1793).

Satyrus Portlandia, *Boisduval & Leconte, Léop. Am. Sept.* pl. 58 (1833).

Debis Portlandia, *Westwood in Gen. Diurn. Lepid.* p. 360. n. 9 (1851).

Oreas marmorea Andromacha, *Hübner, Samml. exot. Schm.* Band i. pl. 89. figs. 1, 2 (1805).

United States.

B.M.

2. *Lethe Europa*.

♀. *Papilio Europa*, *Fabricius, Syst. Ent.* p. 500 (1775).

♂ ♀. *Oreas marmorea Europa*, *Hübner, Samml. exot. Schm.* Band i. pl. 90. figs. 1-4 (1805).

Lethe Europa, *Hübner, Verz. bek. Schm.* p. 56. n. 534 (1816).

Satyrus Europa, *Godart, Enc. Méth.* ix. p. 478. n. 6 (1819).

Debis Europa, *Westwood in Gen. Diurn. Lepid.* p. 359. n. 1 (1851).

Papilio Beroë, *Cramer, Pap. Exot.* i. pl. 79. figs. C, D (1779).

Papilio Arete, *Cramer, Pap. Exot.* iv. pl. 313. figs. E, F (1782).

Sylhet, *Penang*, *Punjaub*, *Hong-Kong*, *Java*. ♂ ♀, B.M.

3. *Lethe Drypetis*.

Debis Drypetis, *Hewitson, Exot. Butterf.* iii. p. 76, pl. 38. figs. 11, 12 (1863).

Var. ? *China*, *Northern India*, *Himalayas*, *Landoor*.

♂ ♀, B.M.

4. *Lethe arcuata*, sp. nov. Pl. II. fig. 3.

♂. *Alæ* costa anticarum multo magis convexa, margine externo magis concavo, fascia submedia obsoleta; ciliis posticarum fuscescentibus; aliter velut in *L. Europa*.

Anticæ subtus fascia subbasali latiore, discali abbreviata subtriangulari et costali; ocellis majoribus; aliter velut in *Europa*; posticæ ocellis magis elongatis; apicali tantum unipupillato; fascia media medio latiore; aliter velut in *Europa*.

Exp. alar. unc. 2½.

Celebes.

B.M.

5. *Lethe Darena*.

Debis Darena, *Felder, Reise der Fregatte Novara*, iii. p. 498. n. 862, tab. 68. figs. 4, 5 (1867).

Java (Felder).

Closely allied to *Drypetis*, Hewits. (♀).

6. *Lethe Dyrta*.

Debis Dyrta, *Felder, Reise der Fregatte Novara*, iii. p. 497. n. 860 (1867).

Bengal (Felder).

"Is the size of *D. Rohria* (Fabr.). The margin of the wings is more regularly scalloped between the nervures than in the allied *D. Drypetis* (Hew.), and the outer margin of the front wings not concave."

7. *Lethe embolima*.

Debis embolima, *Butler, Ent. Month. Mag.*, iii. p. 77. n. 3 (1866).

Ceylon.

♂, B.M.

This species is remarkable from the fact of its having no central pale band or subapical spots on the upperside of the front wings, which makes it appear to belong to the ♂ *Samio* group. It is evidently closely allied to *Drypetis*.

8. *Lethe Neelgherriensis*.

♂. *Satyrus* (Cyllo) *Neelgherriensis* and *Neelgherriensis*, *Guérin in Delessert's Voyage dans l'Inde*, pt. 2. p. 74, pl. 21. figs. 1, 1a (1843).

Debis *Neelgherriensis*, *Westwood in Gen. Diurn. Lepid.* p. 533. n. 360 (1852).

♀. *Debis Neelgherriensis* (sic), *Hewitson, Exot. Butterf.* iii. p. 75, pl. 38. fig. 6 (1863).

♂ ♀. *Debis Neelgherriensis* (sic), *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i. p. 217. n. 447 (1857).

(♂) Ceylon, (♀) East Indies.

B.M.

9. *Lethe Daretis*.

♀. *Debis Daretis*, *Hewitson, Exot. Butterf.* iii. p. 75, pl. 38. figs. 7, 8 (1863).

Ceylon.

♀, B.M.

10. *Lethe Dynsate*.

♀. *Debis Dynsate*, *Hewitson, Exot. Butterf.* iii. p. 76, pl. 38. figs. 9, 10 (1863).

Ceylon.

♀, B.M.

11. *Lethe Rohria*.

Papilio Rohria, *Fabricius, Mant. Ins.* ii. p. 45 (1788).

Satyrus Rohria, *Godart, Enc. Méth.* ix. p. 479 (1819).

Debis Rohria, *Westwood in Gen. Diurn. Lepid.* p. 360. n. 4 (1851); also of *Moore* (1857).

Satyrus Isana, *Kollar in Hügel's Kaschmir*, iv. pt. 2. p. 448, tab. 16. figs. 3, 4 (1844).

Debis Isana, *Westwood in Gen. Diurn. Lepid.* p. 360. n. 3 (1851).

Northern India, Nepaul, Sylhet, Java.

♂ ♀, B.M.

The species *P. Penthera* of Fabricius (queried by Westwood as a form of *Debis*) seems to me to be identical with (*Elymnias*) *Dusara* of Horsfield.

12. *Lethe Nicetas*.

Debis Nicetas, *Hewitson, Exot. Butterf.* iii. p. 78, pl. 39. figs. 17, 18 (1863).

East India.

Coll. Hewitson.

13. *Lethe Verma*.

Satyrus Verma, *Kollar in Hügel's Kaschmir*, iv. pt. 2. p. 447, tab. 16. figs. 1, 2 (1844).

Debis Verma, *Westwood in Gen. Diurn. Lepid.* p. 360. n. 5 (1851); also of *Moore* (1857).

Northern India, Nepaul, Sylhet.

B.M.

14. *Lethe Sidonis*.

Debis Sidonis, *Hewitson, Exot. Butterf.* iii. p. 77, pl. 39. fig. 16 (1863).

Sylhet.

B.M.

15. *Lethe Visrava*.

Debis Visrava, *F. Moore, Proc. Zool. Soc. London*, p. 768, pl. 41. fig. 4 (1865).

North-eastern Bengal (Moore).

16. *Lethe Hyrania*.

Satyrus Hyrania, *Kollar in Hügel's Kaschmir*, iv. pt. 2. p. 449. n. 7, tab. 17. figs. 1, 2 (1844).

Debis Hyrania, *Westwood in Gen. Diurn. Lepid.* p. 360. n. 6 (1851).

Himalaya, Simla (Kollar).

17. *Lethe Whitelyi*.

Lethe Whitelyi, *Butler, Ann. & Mag. Nat. Hist.* xx. p. 403,
pl. 9. fig. 8 (1867).

Nagasaki, North Japan.

B.M.

18. *Lethe Diana*.

Debis Diana, *Butler, Proc. Linn. Soc.* ix. p. 55. n. 20 (1866);
Ann. & Mag. Nat. Hist. 3rd ser. xix. pl. 2. figs. 2, 2a (1867).

Hakodadi, North Japan.

B.M.

19. *Lethe Dinarbas*.

♂. *Debis Dinarbas*, *Hewitson, Exot. Butterf.* iii. p. 77, pl. 39.
fig. 15 (1863).

Northern India.

B.M.

20. *Lethe scanda*.

♂. *Debis scanda*, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*
p. 218. n. 451 (1857).

♀. ? *D. Nada*, *F. Moore, ibid.* n. 452 (1857).

Darjeeling (Moore), Northern India. Coll. Ind. Mus. and ♂, B.M.

21. *Lethe Bhairava*.

♀. *Debis Bhairava*, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*
p. 217. n. 450 (1857).

Northern India.

Coll. Ind. Mus. and ♀, B.M.

22. *Lethe Anysis*.

♂. *Debis Anysis*, *Hewitson, Exot. Butterf.* iii. p. 73, pl. 37.
fig. 1 (1862).

East India (Hewitson).

Except in the form of the hind wings, this species seems identical
with *Bhairava*. May it not be the male of that insect?

23. *Lethe Latiaris*.

♀. *Debis Latiaris*, *Hewitson, Exot. Butterf.* iii. p. 74, pl. 37.
fig. 4 (1863).

Sylhet.

♂, B.M.

The male of this species differs, in several particulars, from the
female. The hind wings are less angulated; the hind wings have a
discal hairy streak above, and the central bands below are placed
slightly nearer together.

24. *Lethe Arcadia*.

♀. *Papilio Arcadia*, *Cramer, Pap. Exot.* ii. pl. 116. figs. D, E (1779).

Temenis Arcadia, *Hübner, Verz. bek. Schmett.* p. 34. n. 279 (1816).

Debis Arcadia, *Westwood in Gen. Diurn. Lepid.* p. 359. n. 2 (1851); also of *Moore* (1857).

♀. *Papilio Minerva*, *Fabricius, Ent. Syst.* iii. pt. 1. p. 95. n. 295 (1793).

Satyrus Minerva, *Godart, Enc. Méth.* ix. p. 499. n. 78 (1819).

Satyrus ? *Minerva*, *Westwood in Gen. Diurn. Lepid.* p. 392. n. 53 (1851).

♂. *Satyrus Caumas*, *Godart, Enc. Méth.* ix. p. 479. n. 7 (1819).

Java.

♂, *Horsf. Coll.*

The male of this species has a discal patch of hair on the upper-side of all the wings.

25. *Lethe Sicelis*.

Debis Sicelis, *Hewitson, Exot. Butterf.* iii. p. 73, pl. 37. fig. 3 (1862).

Japan.

B.M.

26. *Lethe Syrcis*.

Debis Syrcis, *Hewitson, Exot. Butterf.* iii. p. 77, pl. 39. figs. 13, 14 (1863).

North China.

B.M.

27. *Lethe Samio*.

♂. *Debis Samio*, *Hewitson & Westwood in Gen. Diurn. Lepid.* p. 360. n. 8, pl. 61. fig. 3 (1851).

East India.

♂, B.M.

28. *Lethe Purana*.

♀. *Debis Purana*, *Felder, Wien. ent. Monatschr.* iii. p. 401 (1859).

Hab. — ?

Is not this the female of *L. Samio* ?

29. *Lethe Sinorix*.

Debis Sinorix, *Hewitson, Exot. Butterf.* iii. p. 78, pl. 39. figs. 19, 20 (1863).

East India.

Coll. *Hewitson*.

30. *Lethe Vindhya*.

Debis Vindhya, *Felder, Wien. ent. Monatschr.* iii. p. 401 (1859).

Assam (Felder).

31. *Lethe Manthara*.

Debis Manthara, *Felder, Reise der Fregatte Novara*, iii. p. 497. n. 861 (1867); *Butler, Ann. & Mag. Nat. Hist.* xx. p. 403, pl. 9. fig. 9 (1867).

Java.

B.M.

"Without doubt a local form of *Mekara* (Moore). The wings of the male are narrower, the front wings in the costal margin, the hind wings in the inner margin, longer than in *Mekara*." It also differs in the paler colouring of the upperside (which in the hind wings shows five instead of four black discal spots), in the straighter discal line below, the more slender and better-defined marginal lines, and the distinct whitish ocelli.

32. *Lethe Mekara*.

♂. Debis Mekara, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i. p. 219. n. 454 (1857).

Darjeeling (Moore), Northern India. Coll. Ind. Mus. & ♂ ♀, B.M.

33. *Lethe Kansa*.

♂ ♀. Debis Kansa, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i. p. 220. n. 456 (1857).

Darjeeling (Moore).

Coll. Ind. Mus.

34. *Lethe Negrito*.

♂ ♀. Debis Negrito, *Felder, Wien. ent. Monatschr.* v. n. 97 (1861).

♂. Debis Isabella, *Butler, Proc. Zool. Soc. London*, p. 41. n. 4, pl. 3. fig. 4 (1866).

Luzon, Burias, Locban (Felder), Philippines.

♂, B.M.

35. *Lethe Chandica*.

♂ ♀. Debis Chandica, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i. p. 219. n. 455 (1857).

East Indies.

Coll. Ind. Mus. and ♂, B.M.

Genus 41. PARARGE, Hübner (1816).

Typical Species *Pararge Aegeria* (*Egeria*).1. *Pararge Aegeria*.

Papilio Aegeria, *Linnaeus*, *Syst. Nat.* ii. p. 771. n. 143 (1766); also of *Fabricius* (1793), *Lewin* (1795), *Hübner* (1805), *Ochsenheimer* (1807).

♀. *Papilio* (Nymph. Gemm.) *Aegeria*, *Esper*, *Europ. Schmett.* i. tab. 7. fig. 1 (1777).

Satyrus Aegeria, *Godart*, *Enc. Méth.* ix. p. 504. n. 89 (1819).
Var., *Vollenhoven*, *Tijdschr. v. Ent.* viii. pl. 2 (1865).

Hipparchia Aegeria, *Stephens*, *Ill. Haust.* i. p. 54 (1828-35); also of *Eversmann* (1844).

Lasiommata Aegeria, *Westwood & Humphreys*, *Brit. Butterf.* p. 65, pl. 17. figs. 7-10 (1840); also of *E. Doubleday* (1844), *Westwood & Hewitson* (1851), *Butler* (1867).

Pararge Egeria (sic), *Hübner*, *Verz. bek. Schmett.* p. 59. n. 568 (1816).

♂. *Satyrus Xiphia*, *Duponchel*, *Lép. de France*, i. pl. 46. figs. 5, 6 (1832).

England, Germany.

♂ ♀, B.M.

Var. *TIRCIS*.

♀. *Satyrus Tircis*, *Godart*, *Lép. de France*, i. p. 163, pl. 8a. fig. 1 (1821).

Lasiommata Tircis, *Butler*, *Ann. & Mag. Nat. Hist.* xix. p. 162. n. 2 (1867).

France.

Var. *MEONE*.

Papilio (Nymph. Gemm.) *Meone*, *Esper*, *Europ. Schmett.* i. tab. 95. cont. 50. fig. 1 (1777).

Papilio Meone, *Cramer*, *Pap. Exot.* iv. pl. 314. figs. E, F (1782); also of *Hübner* (1805), *Ochsenheimer* (1807).

Hipparchia Meone, *Wagner*, *Reisen in der Regentsch. Algier*, *Atlas*, tab. 9 (1841).

Lasiommata Meone, *Butler*, *Ann. & Mag. Nat. Hist.* xix. p. 162. n. 3 (1867).

Satyrus Xiphia ♂, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 84, 85 (1843-45).

Lasiommata Aegeria, var., *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 135 (1844); also of *Westwood* (1851).

(♂ ♀) Germany, (var. ♀) Italy.

B.M.

2. Pararge Xiphia.

Papilio Xiphia, *Fabricius*, *Syst. Ent.* p. 492. n. 215 (1775);
Ent. Syst. iii. pt. 1. p. 95. n. 294 (1793).

Satyrus Xiphia, *Godart*, *Enc. Méth.* ix. p. 509. n. 96 (1819);
 also of *Herrich-Schäffer* (1843-45).

Lasiommata Xiphia, *E. Doubleday*, *List Lep. Brit. Mus.* i. p. 135
 (1844); also of *Westwood* (1851) and *Butler* (1867).

Pararge Xyphia (*sic*), *Hübner*, *Verz. bek. Schmett.* p. 59. n. 569
 (1816).

(♂ ♀) *Madeira*, (♀) *Teneriffe*.

B.M.

3. Pararge Roxelana.

Papilio Roxelana, *Cramer*, *Pap. Exot.* ii. pl. 161. figs. C, D ♂,
 E, F ♀ (1779); also of *Fabricius* (1793), *Hübner* (1805),
Ochsenheimer (1807).

Dira Roxelana, *Hübner*, *Verz. bek. Schmett.* p. 60. n. 570
 (1816).

Satyrus Roxelana, *Godart*, *Enc. Méth.* ix. p. 504. n. 88 (1819);
Lép. de France, i. pl. 30. figs. 1-4 (1832).

Papilio (*Hipparchia*) *Roxelana*, *Freyer*, *Neuere Beiträge*, i. p. 1,
 tab. 25. figs. 1, 2 (1833).

Lasiommata Roxelana, *E. Doubleday*, *List Lep. Brit. Mus.* i.
 p. 136 (1844); also of *Westwood* (1851) and *Butler* (1867).

(♂ ♀) *Smyrna*, (♂ ♀) *Crete*, (♂) *Athens*, (♂) *Syria*. B.M.

4. Pararge Climene.

Papilio (*Nymph. Gemm.*) *Climene*, *Esper*, *Schmett.* i. pt. 2. p. 165,
 tab. 85. cont. 35. figs. 1-3 (1777).

Papilio Climene, *Hübner*, *Samml. europ. Schmett.* i. figs. 165,
 166 (♀ var.) (1805); also of *Ochsenheimer* (1807).

Satyrus Climene, *Godart*, *Enc. Méth.* ix. p. 541. n. 165 (1819);
 also of *Duponchel* and *Boisduval* (1832).

Hipparchia Climene, *Eversmann*, *Fauna Lepidopt. Volgo-Ural.*
 p. 28. n. 13 (1844).

Lasiommata Climene, *Butler*, *Entomologist* (1867).

Papilio Clymene (*sic*), *Fabricius*, *Ent. Syst.* iii. pt. 1. p. 242. n. 753
 (1793) [*nec Erymanthia*].

Satyrus Clymene, *Herrich-Schäffer*, *Eur. Schmett.* i. figs. 102,
 103, 472, 473 (1843-45).

Lasiommata Clymene, *Westwood in Gen. Diurn. Lepid.* p. 386.
 n. 1 (1851).

Lasiommata Clymenus, *E. Doubleday*, *List Lep. Brit. Mus.* i.
 p. 136 (1844).

Epinephele Synclimene, *Hübner*, *Verz. bek. Schmett.* p. 59. n. 567
 (1816).

Russia.

♂ ♀, B.M.

Pararga Clymene, var. *Caucasica* Roxandra, Nordmann, *Bull. Soc. Imp. Nat. Mosc.* p. 400, tab. 10. figs. 1-4 (1851).

Russia.

♂, B.M.

P. Erymanthia does not belong to this genus. It is probably a species of *Epinephele*.

5. *Pararge Epimenides*.

Lasiommata *Epimenides*, *Ménétriés in Schrenck's Amur-Lande*, p. 39. n. 75, tab. 3. figs. 8, 9 (1859).

Amoor Land.

6. *Pararge Schrenckii*.

Pronophila *Schrenckii*, *Ménétriés, Bull. Acad. St. Pétersb.* xvii. p. 215. n. 13 (1859); *Reise und Forsch. Amur-Lande*, p. 33. n. 65; and *P. Schrenckii* (sic), tab. 3. fig. 3 (1859).

Nemophila Schrenckii, *Bremer, Bull. Acad. Imp. Sci. St. Pétersb.* p. 469. n. 12 (part.) (1861).

Bureya Mountains.

Allied to *Epimenides*.

7. *Pararge Maackii*.

Lasiommata *Maackii*, *Bremer, Lep. Ost-Sibir. u. Amur. in Bull. Acad. Imp. Sci. St. Pétersb.* p. 468. n. 12 (1861).

Ussuri.

Closely allied to the preceding, but much smaller, the front wing more produced, and with three ocelli below; the hind-wing ocelli nearer the margin, and surrounded by grey irides.

8. *Pararge Ménétériésii*.

Satyrus Ménétériésii, *Bremer & Grey, Fauna des nordl. China's*, p. 8. n. 25 (1853); *Ménétriés, Enum. Corp. Anim. Lepid.* pt. 1. p. 47. n. 781, Taf. 6. fig. 4 (1855).

North China.

9. *Pararge Achine*.

Papilio Achine, *Scopoli, Ent. Carn.* p. 150. n. 433. fig. 433 (1763).

Papilio Dejanira, *Linnaeus, Syst. Nat.* ii. p. 774. n. 154 (1766); also of *Fabricius* (1793), *Hübner*, figs. 170, 171 (1805), *Ochsenheimer* (1807).

Papilio (Nymph. Gemm.) *Dejanira*, *Esper, Schmett.* i. p. 125, tab. 9. fig. 2 (1777).

Enodia Dejanira, *Hübner, Verz. bek. Schmett.* p. 61. n. 588 (1816).

- Satyrus Dejanira*, *Godart, Enc. Méth.* ix. p. 509. n. 96 (1819);
Lép. de France, i. pl. 8. fig. 1 (1821).
Hipparchia Dejanira, *Eversmann, Fauna Lepidopt. Volgo-Ural.*
 p. 30. n. 17 (1844).
Papilio (Hipparchia) Dejanira, *Freyer, Neuere Beiträge*, tab. 391
 (1845).
Lasiommata Dejanira, *Westwood in Gen. Diurn. Lepid.* p. 386.
 n. 10 (1851); also of *Butler* (1867).
 (♂ ♀) Polish Ukraine, (♂) Germany, (♂) South Europe. B.M.

10. *Pararge Deidamia*.

- Hipparchia (Pararge) Deidamia*, *Eversmann, Bull. Soc. Imp.*
Nat. Mosc. p. 617. n. 6 (1851).
Lasiommata Deidamia, *Westwood in Gen. Diurn. Lepid. Suppl.*
 p. 533. n. 386 (2) (1852).
 Irkoutzk, Eastern Siberia (*Eversmann*).

It is doubtful whether this is more than a local form of *Achine*.

11. *Pararge Canthus*.

- Papilio Canthus*, *Linnaeus, Syst. Nat.* (1748) 12th ed. p. 768.
 n. 129 (1766); also of *Fabricius* (1793).
Satyrus Canthus and Cantheus, *Godart, Enc. Méth.* ix. p. 493.
 n. 35, 36 (1819); *Boisduval & Leconte, Icon. Lép. Am. Sept.*
 pl. 60 (1833).
Neonympha Canthus, *Westwood in Gen. Diurn. Lepid.* p. 375.
 n. 13 (1851).
Papilio Eurydice, *Linnaeus, Amoen. Acad.* tom. vi. p. 406. n. 65
 (1763).
 Canada. B.M.

Genus 42. AMECERA, *Butler* (1867).

Typical Species *Ameccera Megæra*.

1. *Ameccera Megæra*.

- Papilio Megæra*, *Linnaeus, Syst. Nat.* ii. p. 771. n. 142 (1766);
 also of *Fabricius* (1793), *Lewin* (1795), *Hübner* (1805),
 and *Ochsenheimer* (1807).
P. Nymphalis Megæra, *Esper, Schmett.* i. p. 101, tab. 6. fig. 3
 (1777).
Dira Megæra, *Hübner, Verz. bek. Schmett.* p. 60. n. 571 (1816).
Satyrus Megæra, *Godart, Enc. Méth.* ix. p. 503. n. 87 (1819).
Hipparchia Megæra, *Stephens, Ill. Haust.* i. p. 55 (1828).
Lasiommata Megæra, *Westwood & Humphrey's Brit. Butterf.*
 p. 66, pl. 18. figs. 1-5 (1840); also of *E. Doubleday* (1844)
 and *Westwood & Hewitson* (1851).

Pararge Megæra, *Ström, Naturhist. Tidsk.* (1866).

Amecera Megæra, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 163. n. 1 (1867).

Papilio Mæra, *Borkhausen, Schmett.* i. p. 125 (1788).

England, Southern Europe.

B.M.

Var. Lyssa.

Papilio Lyssa, *Hübner, Eur. Schmett.* i. figs. 114–117 (1805); also of *Boisduval* (1832).

♀. Satyrus Lyssa, *Boisduval, Icon. Hist. Lép.* p. 222, pl. 44. figs. 4, 5 (1832).

Satyrus Megæra ♀, *Godart, Lép. de France*, i. pl. 7 secund. fig. 3 (1821). P. Megæra, var., *Esper, Eur. Schmett.* tab. 6. fig. 3 (1777).

Dalmatia.

B.M.

2. *Amecera Adrasta.*

Papilio Adrasta, *Hübner, Eur. Schmett.* i. figs. 836–839 (1805).

Papilio (Nymph. Gemm.) Megæra, var., *Esper, Schmett. Eur.* i. pt. 2. tab. 68. cont. 18. fig. 2 (1777).

(♂ ♀) Germany, (♀) Naples.

B.M.

This has been mistaken for a variety of *Mæra*; but it seems very evident that it is either a distinct species or a variety of *Megæra*. The male figured by Hübner is much darker than usual, and might almost be taken for the common Swiss type of *Mæra*; but a careful examination of the markings on the underside at once points out its distinctness. We have one pair of the insect agreeing with Hübner's figures in the coloration and pattern of the underside; but the upperside of our male specimen has more orange colouring in the front wings. Our female from Naples differs slightly from the German one, the band of the front wings being placed rather nearer to the outer margin.

Var. Differt ocello apicali unipupillato, fascia magis undata, posticis subtus brunnescentibus, ocellos tres supra habentibus.

Persia.

B.M.

This variety, at first sight, can scarcely be distinguished from *Megæra*.

3. *Amecera Paramegæra.*

Papilio Paramegæra, *Hübner, Eur. Pap.* i. pl. 170. figs. 842–844 (1805).

Papilio Tigelius, *Bonelli in Mem. Accad. Torino*, xxx. t. 1. fig. 2 (1826).

Satyrus Tigelius, *Boisduval, Icon. Hist. Lépid.* p. 224. n. 13, pl. 45. figs. 1-3 (1832); also of *Duponchel*.

Papilio (*Hipparchia*) *Tigelius*, *Freyer, Neuere Beiträge*, i. t. 68. fig. 1 (1833).

Lasionmata Tigelius, *Westwood in Gen. Diurn. Lepid.* p. 386. n. 7 (1851).

Amecera Tigelius, *Butler in Ann. & Mag. Nat. Hist.* xix. p. 163. n. 3 (1867).

Sardinia, Corsica.

♂ ♀, B.M.

4. *Amecera Xiphia*.

Satyrus Xiphia, *Boisduval, Icon. Hist. Lépid.* p. 223. n. 12, pl. 44. figs. 6, 7 (1832); but not the *Xiphia* of *Fabricius*.

Portugal, Spain, Teneriffe, Madeira (*Boisduval*).

This insect has nothing to do with the *Xiphia* of *Fabricius*. *M. Boisduval* suggests the possibility of its being a variety of *Meone*, which it neither resembles in structure, pattern, nor coloration. He says:—"Il a de très-grands rapports avec la variété *Meone* de l'*Ægeria*, et il serait bien possible qu'il ne fût lui-même qu'une variété plus prononcée."

5. *Amecera Mæra*.

Papilio Mæra, *Linnaeus, Syst. Nat.* ii. p. 771. n. 141 (1766); also of *Fabricius* (1793).

Maniola Mæra, *Schrank, Fauna Boica*, Band ii. p. 172. n. 388 (1801).

Satyrus Mæra, *Godart, Enc. Méth.* ix. p. 501. n. 86 (1819).

Lasionmata Mæra, *Westwood in Gen. Diurn. Lepid.* p. 386. n. 3 (1851).

Amecera Mæra, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 163. n. 4 (1867).

Italy, France.

♂ ♀, B.M.

♀. Var. *a*. Supra multo magis flavescens, ocello apicali ovali bipupillato: subtus lineis omnibus multo distinctioribus, lineis marginalibus distincte sinuatis. *Esper, Schmett.* ii. tab. 68. cont. 18. fig. 1 (1777).

South Europe.

♀, B.M.

♂. var. *b*. *Papilio* (*Nymph. Gemm.*) *Mæra*, *Esper, Eur. Schmett.* i. tab. 6. fig. 2 (1777).

France.

♂ ♀, B.M.

Sp. ?? ♂. *Papilio Mæra*, *Hübner, Eur. Schmett.* i. pl. 39. figs. 174, 175 (1805).

♀. *Papilio* (*Nymph. Gemm.*) *Mæra*, *Esper, Eur. Schmett.* i. pt. 2. tab. 68, cont. 18. figs. 3 (1777).

(♂ ♀) Italy, (♀) Switzerland.

B.M.

It is uncertain whether this is distinct from *Mæra* or no; it seems to be common all over Switzerland; and though a good many specimens have been taken, it does not seem to vary. It is, however, very like the true *Mæra*, from which it differs in its more robust form, darker colouring both above and below; in the form of the front-wing ocelli, in having only two distinct ocelli on the upperside of the hind wing, and in the less irregular central lines on the underside of the hind wings.

6. *Amecera Mærula*.

Lasiommata Mærula, *Felder, Reise der Fregatte Novara*, iii. p. 496. n. 858 (1867).

Eastern Himalayas (Felder).

Intermediate between *Mæra* and *Schakra*.

7. *Amecera Menava*.

♂ ♀. *Lasiommata Menava*, *F. Moore, Proc. Zool. Soc.* p. 499. n. 83, pl. 30. fig. 3 (1865).

Amecera Menava, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 163. n. 8 (1867).

♂. *Lasiommata mæroides*, *Felder, Reise der Fregatte Novara*, iii. tab. 69. fig. 1 (1867).

(♂ ♀) Middle Kunawur (Moore), (♂) Germany. B.M.

8. *Amecera Schakra*.

Satyrus Schakra, *Kollar in Hügel's Kaschmir*, p. 446, tab. 15. figs. 3, 4 (1848).

Lasiommata Schakra, *Westwood in Gen. Diurn. Lepid.* p. 387. n. 12 (1851).

Amecera Shakra (*sic*), *Butler, Ann. & Mag. Nat. Hist.* xix. p. 163. n. 7 (1867).

Himalayas.

♂, B.M.

9. *Amecera Maderakal*.

Satyrus Maderakal, *Güérin-Méneville, Voy. en Abyssin.* vi. p. 381, pl. 10. figs. 1-3 (1849); also of *Westwood* (1852).

Abyssinia.

♂, B.M.

This is a very distinct species; the hind wings are of a peculiar kite-like shape, and the front wings have the apex very acute.

10. *Amecera Hiera*.

♀. *Papilio Hiera*, *Hübner, Europ. Schmett.* i. pl. 39. fig. 176 (1805).

Satyrus Hiera, *Boisduval, Icones Hist. Léop.* p. 220. n. 10, pl. 44. figs. 1-3 (1832).

Lasiommata Hiera, *Westwood in Gen. Diurn. Lepid.* p. 386.
n. 4. (1851).

Amecera Hiera, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 163. n. 6
(1867).

Helvetia.

♂ ♀, B.M.

Var. *Alis magis abruptis, fasciis subtus distinctioribus.*

South Europe.

♂ ♀, B.M.

11. *Amecera Eversmanni*.

Hipparchia Eversmanni, *Eversmann in Bull. Soc. Imp. Nat. Mosc.*
pl. 2. figs. 3-6 (1847).

Lasiommata Eversmanni, *Westwood in Gen. Diurn. Lepid.*
p. 386. n. 11 (1851).

Amecera Eversmanni, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 163.
n. 5 (1867).

Dauria ?

12. *Amecera Baldiva*.

Lasiommata Baldiva, *F. Moore, Proc. Zool. Soc.* p. 499. n. 84,
pl. 30. fig. 4 (1865).

Amecera Baldiva, *Butler, Ann. & Mag. Nat. Hist.* xix. p. 163.
n. 9 (1867).

Genus 43. *IDIOMORPHUS*, *Doumet* (1861).

Typical Species *Idiomorphus Hewitsonii*.

1. *Idiomorphus Hewitsonii*.

Idiomorphus Hewitsonii, *Doumet, Rev. et Mag. de Zool.* p. 174,
pl. 5. figs. 2, 2 a, 2 b (1861).

Hab. — ?

2. *Idiomorphus Iocius*.

Idiomorphus Iocius, *Hewitson, Exot. Butterf.* iii. p. 96, pl. 48.
figs. 4, 5 (1865).

Old Calabar.

Coll. Hewitson.

This species somewhat resembles the preceding in colour; it, however, has no violet marginal band in the hind wings, and differs in form and other particulars. *I. Hewitsonii* is similar in outline to *Clerome Busiris* of Westwood from Sumatra, &c.

3. *Idiomorphus Italus*.

Idiomorphus Italus, *Hewitson, Exot. Butterf.* iii. p. 95, pl. 48.
figs. 1-3 (1865).

Old Calabar.

Coll. Hewitson.

Genus 44. BLETOGONA, *Felder* (1867).Typical Species *Bletogona Mycalesis*.1. *Bletogona Mycalesis*.*Bletogona Mycalesis*, *Felder, Reise der Fregatte Novara*, iii. tab. 68. figs. 6, 7 (1867).Celebes ? (*Felder*).Genus 45. MYCALESIS, *Hübner* (1816).Typical Species *Mycalesis Evadne*.1. *Mycalesis Evadne*.*Papilio Evadne*, *Cramer, Pap. Exot.* iii. pl. 222. figs. E, F (1782).*Mycalesis Evadne*, *Hübner, Verz. bek. Schmett.* p. 55. n. 521 (1816).*Satyrus Servatius*, *Godart, Enc. Méth.* ix. p. 525. n. 130 (1819).Sierra Leone, Africa (*Cramer*).

Closely allied to *Eusirus* of *Hopffer*, from which it chiefly differs in having a purplish gloss and no ocelli on the upperside, and a more distinct subbasal transverse streak on the underside.

2. *Mycalesis funebris*.*Satyrus funebris*, *Guérin, Icon. Règne Anim.* p. 488 (1844).*Mycalesis funebris*, *Westwood in Gen. Diurn. Lepid.* p. 394. n. 6 (1851).

Cazamanca, Senegal.

According to M. Guérin-Ménéville this species is allied to (*Servatius*) *Evadne*: it has ocelli near the margin of the hind wings on the upperside.

3. *Mycalesis Saftza*.*Mycalesis Saftza*, *Hewitson, Gen. Diurn. Lepid.* p. 394. n. 10, pl. 66. fig. 3 (1851); *Exot. Butterf.* iii. p. 80. pl. 40. fig. 4 (1862).

Africa, Congo, Port Natal.

B.M.

This insect seems to be very closely allied to *Eusirus*.

4. *Mycalesis Eusirus*.*Mycalesis Eusirus*, *Hopffer in Peters's Reise'n. Mossamb. Ins.* p. 393, pl. 25. figs. 3, 4 (1855).*Mycalesis Evenus*, var., *Trimen in Rhop. Afric. Austral.* p. 207. n. 120 & p. 208 (obs.) (1866).

Congo.

B.M.

Mr. Trimen, having received a single specimen of the variety *injusta*, has considered himself fully justified in sinking the species *Eusirus*, which he places as a synonym of the more recent *Evenus*; *injusta* does not, however, link these species, nor do any connecting forms appear to exist.

5. *Mycalesis Milyas*.

Mycalesis Milyas, *Hewitson, Exot. Butterf.* iii. p. 89, pl. 45.
fig. 34 (1864).

White Nile.

B.M.

Possibly a local form of the preceding species.

6. *Mycalesis nebulosa*.

Mycalesis nebulosa, *Felder, Reise der Fregatte Novara*, iii. p. 502.
n. 871 (1867).

Guinea (Felder).

"The wings are as broad as in *M. Safitza*, Hew.; but the costa is shorter, and the hind wing more produced at the anal angle." Described from a specimen larger than *M. Evenus*, Hopffer.

7. *Mycalesis Evenus*.

Mycalesis Evenus, *Hopffer, Peters's Reise n. Mossamb. Ins.* p. 394,
pl. 25. figs. 5, 6 (1855).

Var. ♂. *Mycalesis injusta*, *Wallengren, Lep. Rhop. Caffr. in Kongl. Svensk. Vetensk. Akad. Handl.* 2nd ser. p. 33. n. 1 (1857).

(*Injusta*) Port Natal, Cape of Good Hope.

B.M.

8. *Mycalesis Caffra*.

♂. *Mycalesis Caffra*, *Wallengren, Lep. Rhop. Caffr. in Kongl. Svensk. Vetensk. Akad. Handl.* 2nd ser. p. 34. n. 2 (1857).

♀. *Mycalesis Gambius*, *E. Doubleday, List. Lep. Brit. Mus.* i. p. 139 (1844).

Alis supra ocellis anticarum distinctioribus; subtus posticis ocellis punctiformibus ocellaribus, area basali pallidiore.

(♂ ♀) Knysna, (♀) Zoulu.

♂ ♀, B.M.

9. *Mycalesis Aramis*.

Mycalesis Aramis, *Hewitson, Exot. Butterf.* iii. p. 91, pl. 46.
figs. 46, 47 (1866).

Philippine Islands.

Coll. Hewitson.

Seems to be allied to the South African *M. Caffra*.

10. *Mycalosis Rhacotis*.

Mycalosis Rhacotis, Hewitson, *Exot. Butterf.* iii. p. 93, pl. 47. fig. 50 (1866).

Old Calabar.

Coll. Hewitson.

Closely allied to *Evenus* (var. *injuncta*): it is almost identical in pattern with *M. Blasius* of Fabricius (var. *Samba*).

11. *Mycalosis Sciathia*.

Mycalosis Sciathia, Hewitson, *Exot. Butterf.* iii. p. 94, pl. 47. fig. 55 (1866).

Old Calabar.

12. *Mycalosis saga*, sp. nov. Pl. III. fig. 1.

Alæ supra piceo-fuscae, apice anticarum et marginibus externis pallidioribus, linea submarginali undata nigra interrupta.

Alæ subtus velut in *Caffra*, sed obscuriores et apice pallidore: anticæ ocellis tribus punctiformibus subapicalibus oblique positis; unoque nigro parvo discali: posticæ ocellis septem, tertio, quarto et quinto fuscis, aliis nigris.

Exp. alar. unc. $2\frac{1}{2}$.

Sierra Leone.

B.M.

13. *Mycalosis vulgaris*, sp. nov. Pl. III. fig. 2.

Alæ supra velut in *Safitea* obscuriores, rarius macula subapicali diffusa ochracea.

Alæ subtus pallidiores, omnino lituratae, fascia lata media irregulari obscuriore, utrinque fusco limitata; ocellis velut in *Caffra*, inferiore autem anticarum discali minore; linea marginali et margine ipso fuscescentibus; ciliis griseis.

Exp. alar. unc. $1\frac{1}{2}$ –2.

Sierra Leone.

B.M.

14. *Mycalosis angulosa*, sp. nov. Pl. III. fig. 8.

Alæ supra, velut in *Saga*, piceo-nigræ, margine externo paulum dilutiore; sacculo parvo ad venæ anticarum submedianæ medium ochraceo; margine costali posticarum iridescente, maculam circumratam nigram post costæ medium ferente; cirris consuetis basalibus ochraceis.

Alæ subtus velut in specie præcedente scriptæ, sed magis violascentes, area basali obscuriore, apice subquadrato ochreo tincto, lineis omnibus magis nigrescentibus, externa posticarum media multo magis angulata; ocellis minoribus, partim cinereo roratis.

Exp. alar. unc. $1\frac{1}{2}$.

Sierra Leone.

Closely allied to the preceding species.

15. *Mycalesis auricruda*, sp. nov. Pl. III. fig. 6.

Alæ supra velut in *Sanao* piceo-nigræ, fascia autem anticis subapicali bimaculari abbreviata ochrea; costa posticarum pallide olivacea.

Alæ subtus fuscæ, area basali medioque anticarum disci ochreis, fusco lituratis, lineis mediis sinuatis et dentatis, interna valde indistincta, externa ad costam anticarum introrsum arcuata, posticarum obliqua minime toto concava; lineis duabus marginalibus nigris: antica ocellis duobus velut in *Eveno*: posticæ duobus apicalibus tribusque analibus velut in *Eveno* positis; his omnibus distinctis et late violaceo cinctis.

Exp. alar. unc. $1\frac{1}{8}$.

Ashanti.

B.M.

16. *Mycalesis Sanaos*.

Mycalesis Sanaos, Hewitson, *Exot. Butterf.* iii. p. 94, pl. 47. figs. 53, 54 (1866).

Old Calabar (Hewitson), Ashanti.

Var. ?, B.M.

17. *Mycalesis Elionas*.

Mycalesis Elionas, Hewitson, *Exot. Butterf.* iii. p. 91, pl. 46. figs. 41, 42 (1866).

Old Calabar (Hewitson).

This insect is in Mr. Hewitson's collection; it seems allied to the preceding species.

18. *Mycalesis Maianeas*.

Mycalesis Maianeas, Hewitson, *Exot. Butterf.* iii. p. 87, pl. 44. figs. 27, 28 (1864).

Malacca (Hewitson), Borneo.

B.M.

This, though an Indian insect, evidently belongs to the present group.

19. *Mycalesis Xeneas*.

♀. *Mycalesis Xeneas*, Hewitson, *Exot. Butterf.* iii. p. 92, pl. 46. fig. 48; ♂, p. 93, pl. 47. fig. 49 (1866).

Old Calabar.

Coll. Hewitson.

This species seems to be intermediate between the *Eusirus* and *Asochis* groups, the male being most like the former, whilst the female more nearly resembles the latter form.

20. *Mycalesis Asochis*.

Mycalesis Asochis, Hewitson, *Exot. Butterf.* iii. p. 92, pl. 46. figs. 46, 47 (1866).

Gaboon.

B.M.

In the Museum specimen of this species there are two nearly parallel central lines on the underside of the wings.

21. *Mycalesis Eliasius*.

Mycalesis Eliasius, *Hewitson, Exot. Butterf.* iii. p. 91, pl. 46. figs. 44, 45 (1866).

Congo.

B.M.

Allied to *Asochis*, which it seems to unite with the *Anapita* group.

22. *Mycalesis Dorothea*.

♀. *Papilio Dorothea*, *Cramer, Pap. Exot.* iii. pl. 204. figs. E, F (1782).

Neonympha Dorothea, *Hübner, Verz. bek. Schmett.* p. 65. n. 625 (1816).

♂. *Papilio Melusina*, *Fabricius, Ent. Syst.* iii. pt. 1. p. 240. n. 750 (1793).

Satyrus Melusina, *Godart, Enc. Méth.* ix. p. 496. n. 66 (1819).

Mycalesis? *Melusina*, *Westwood in Gen. Diurn. Lepid.* p. 394. n. 15 (1851).

Sierra Leone.

♂, B.M.

Var. RÆSACES.

♂ ♀. *Mycalesis Ræsaes*, *Hewitson, Exot. Butterf.* iii. p. 93, pl. 47. figs. 61, 62 (1866).

Old Calabar.

Coll. Hewitson.

Var. MIRIAM.

Papilio Miriam, *Fabricius, Ent. Syst.* iii. pt. 1. p. 242. n. 754 (1793).

Angola.

♂, B.M.

23. *Mycalesis Otreia*.

Papilio Otreia, *Cramer, Pap. Exot.* iv. pl. 314. figs. A, B (1782).

Mycalesis Otreia, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* p. 230. n. 486 (1857).

Mycalesis Polydecta, var.?, *Westwood in Gen. Diurn. Lepid.* p. 393. n. 1 (part.) (1851).

Java (Horsfield Coll.), (var.) Canara.

B.M.

Var. FRANCISCA.

Papilio Francisca, *Cramer, Pap. Exot.* iv. pl. 326. figs. E, F (1782). *Franciscus*, *Herbst* (1796).

Var. Satyrus Zopyrus, *Kollar in Hügel's Kaschmir*, p. 450. n. 8 (1842). *Zophyrus*, *Westwood* (1851).

China (Cramer), var. *Zopyrus*, North and East India.

B.M.

Var. VISALA.

Mycalesis Visala, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i. p. 230. n. 488 (1857).

Sylhet.

Coll. Ind. Mus. and B.M.

Chiefly differs from the typical *Otrea* in the more sharply angulated form of the front wings.

Var. PERSEUS.

Papilio Perseus *, *Fabricius, Syst. Ent.* p. 488. n. 199 (1775);
Butler, P. Z. S. p. 719. fig. 2 (1867).

Papilio Tabitha, *Fabricius, Ent. Syst.* iii. pt. 1. p. 243. n. 756 (1793).

North India.

B.M.

Var. CLERIMON.

Papilio Clerimon, *Fabricius, Ent. Syst.* iii. pt. 1. p. 217. n. 678 (1793).

Satyrus Clerimon, *Godart, Enc. Méth.* ix. p. 512. n. 105 (1819).

Lasiommata? *Clerimon*, *Westwood in Gen. Diurn. Lepid.* p. 387. n. 16 (1851).

Hab. — ?

This appears to be only another variety of *Otrea*.

24. *Mycalesis Igoleta*.

Mycalesis Igoleta, *Felder, Wien. ent. Monatschr.* v. (1861);
also in *Lep. Diagn.* n. 103.

Luzon (Felder).

According to Felder, possibly the Philippine form of *Polydecta*; he should rather say, of *Otrea*; for *Polydecta* seems to be allied to *Malsara* of Moore.

25. *Mycalesis Cæsonia*.

Mycalesis Cæsonia, *Wallengren in Wien. entom. Monatschr.* iv. p. 36. n. 12 (1860).

Manilla (Wallengren).

Probably closely allied to *Igoleta*, from which, however, it seems to be sufficiently distinct. Wallengren remarks that it is scarcely larger than *Justina*.

26. *Mycalesis Drusia*.

Papilio Drusia, *Cramer, Pap. Exot.* i. pl. 84. figs. C, D (1779).

* The type of this insect is in the Banksian Collection. Its chief peculiarity consists in the small size of the front-wing ocellus on the upperside.

Papilio Mineus, var.?, *Westwood in Gen. Diurn. Lepid.* p. 393.
n. 3 (syn.) (1851).

China (Cramer).

It seems improbable that this is a variety of *Mineus*, as the position of the central band and the ocelli of the underside are quite different. It is just possible that Cramer's figure is from a broken specimen, which might account for these differences; but there is still a distinctive character in the dark colour of the ocellus-irides and the different form of the marginal lines. Mr. Moore has considered some specimens of *Mineus* to represent *Drusia*.

27. *Mycalesis Malsarida*, sp. nov. Pl. III. fig. 14.

Alæ supra rufo-fuscae, area apicali anticarum minime dilutior; linea submarginali et margine externo obscurioribus undulatis, linea marginali et posticis intus altera ochraceis subdistinctis; costa posticarum ochracea, ad basin sericea.

Alæ subtus minime obscuriores, area basali fusco liturata; fascia media anticis abbreviata subintegra obliqua violaceo-plumbea; altera ocellos circumcingente tertiaeque submarginali violaceis; linea submarginali et margine ipso nigris, linea marginali fulva; his omnibus undulatis: anticæ ocellis tribus subapicalibus parvis (medio paulum majore) oblique positis unoque permagno discali: posticæ ocellis septem, secundo, tertio, sexto et septimo parvis, quinto maximo; his omnibus nigerrimis albo pupillatis et fulvo cinctis.

Exp. alar. unc. 2.

Sylhet.

B.M.

28. *Mycalesis Gotama*.

Mycalesis Gotama, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*
p. 232. n. 493 (1857).

? *Mycalesis Borealis*, *Felder, Reise der Fregatte Novara*, iii. p. 500.
n. 867 (1867).

China, Shanghai.

Coll. Ind. Mus. and B.M.

This is a very distinct species. It is related to *Mineus* of Linnaeus.

29. *Mycalesis Heri*.

Mycalesis Heri, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*
p. 223. n. 496 (1857).

Bootan.

Coll. Ind. Mus.

Seems to be allied to *Mineus*.

30. *Mycalesis Cepheus*.

Mycalesis Cepheus, *Butler, Ann. & Mag. Nat. Hist.* xx. p. 402,
pl. 9. figs. 3, 4 (1867).

Penang.

Coll. Roberts.

31. *Mycalesis Mineus*.

Papilio Mineus, *Linnaeus*, *Syst. Nat.* ii. p. 768. n. 126 (1766); also of *Fabricius*.

Satyrus Mineus, *Godart*, *Enc. Méth.* ix. p. 510. n. 97 (1819); also of *Boisduval*.

Mycalesis Mineus, *Westwood in Gen. Diurn. Lepid.* p. 393. n. 3 (1851)*.

* China (Hongkong).

♂ ♀, B.M.

Var. Alæ anticæ subtus ocello subapicali geminato.

China.

♂ ♀, B.M.

Var. JUSTINA.

Papilio Justina, *Cramer*, *Pap. Exot.* iv. pl. 326. fig. C (1782).

Mycalesis Justina, *Hübner*, *Verz. bek. Schmett.* p. 55. n. 524 (1816).

Sylhet.

♂, B.M.

32. *Mycalesis Justinella*, sp. nov. Pl. III. fig. 12.

Alæ supra velut in *Justina* (Cramer); ocello autem iride magis distincta.

Alæ subtus fascia discali latiore alba; ocello discali inferiore anticarum geminato: posticæ area interna late albido rorata; linea interna media posticis distincta, ocellisque albido nec ochreo circumcinctis.

Exp. alar. unc. 1½.

Philippine Islands.

B.M.

Possibly the Philippine representative of *Mineus*. It is most like the form *Justina* as regards the position of the bands and ocelli on the underside. Our specimen is rather smaller than *M. Gotama*.

33. *Mycalesis Polydecta*.

Papilio Polydecta, *Cramer*, *Pap. Exot.* ii. pl. 144. figs. E, F (1779).

Satyrus Polydecta, *Godart*, *Enc. Méth.* ix. p. 511. n. 101 (1819).

Mycalesis Polydecta, *Westwood in Gen. Diurn. Lepid.* p. 393. n. 1 (1851); *Butler*, *Ann. & Mag. Nat. Hist.* xx. p. 402, pl. 9. figs. 5, 6 (1867).

Tranquebar (Cramer), Borneo (Butler), (var.) Sumatra, Celebes.

B.M.

Cramer states that his specimen of this species was from Tranquebar. The Bornean specimens essentially agree with Cramer's figure; but there can be little doubt that his locality is correct.

* *M. Blasius* is a distinct species closely allied to, if not identical with, *M. Samba* of Moore. Westwood, however, makes it a synonym of *Mineus*.

34. **Mycalesis Mamerta.**

Papilio Mamerta, *Cramer, Pap. Exot.* iv. pl. 326. fig. D (1782).
China (Cramer).

Var. MAISARA.

Mycalesis Malsara, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*
p. 231. n. 492 (1851).

Himalayas.

B.M.

This may be considered only another form of *Mamerta*, which is possibly a variety of *Polydecta*, and even may be an extreme variety of *Mineus*, to which it is evidently allied.

35. **Mycalesis Orseis.**

Mycalesis Orseis, *Hewitson, Exot. Butterf.* iii. p. 89, pl. 45.
figs. 36, 37 (1864).

Sumatra.

Colls. Saunders and Hewitson.

36. **Mycalesis Jopas.**

Mycalesis Jopas, *Hewitson, Exot. Butterf.* iii. p. 86, pl. 43. fig. 24
(1862).

East India.

Coll. Hewitson.

Allied to *Mamerta*, and also to *Hesione*.

37. **Mycalesis Janardana.**

Mycalesis Janardana, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*
i. p. 234. n. 502 (1857).

Java.

Horsf. Coll.

Allied to *Megamede* of Hewitson and to *Mamerta* of Cramer.

38. **Mycalesis Megamede.**

Mycalesis Megamede, *Hewitson, Exot. Butterf.* iii. p. 84, pl. 42.
fig. 14 (1862).

Ternate.

B.M.

Allied to *Polydecta* and *Mamerta*.

39. **Mycalesis Nautilus.**

Mycalesis Nautilus, *Butler, Ann. & Mag. Nat. Hist.* xx. p. 402,
pl. 9. fig. 7 (1867).

Malacca, India.

♂ ♀, Coll. Roberts.

40. *Mycalesis Blasius*.

Papilio Blasius, *Fabricius, Ent. Syst. Suppl.* v. p. 426. n. 488-489 (1798); *Butler, P. Z. S.* p. 720. fig. 4 (1867).

East Indies (*Fabricius*), Philippines.

B.M.

Var. Mycalesis Samba, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* p. 233. n. 498 (1857). *Alis paulum majoribus.*

North India (*Moore*).

Coll. Ind. Mus.

Seems to be closely allied to *M. Blasius* of *Fabricius*, which has, however, three ocelli on the underside of the front wings.

Var. LALASSIS.

Mycalesis Lalassis, *Hewitson in Exot. Butterf.* iii. p. 89, pl. 45. fig. 35 (1864).

Gilolo (*Hewitson*), Gilolo?, North India, Ceylon, (var.) Java.

B.M.

Differs from *Blasius* and *Samba* in the distinct subanal spot of the front wings above, the fainter central band on the underside, also in having four instead of two distinct ocelli in the front wings.

41. *Mycalesis Semperi*, sp. nov. Pl. III. fig. 7.

Alæ supra olivaceo-fuscæ, area basali paulum pallescente: ocello alis ambabus subanali nigro, brunneo cincto et albo pupillato.

Alæ subtus velut in injusta fere scriptæ, linea autem discali magis ad marginem approximante, ocellis nec distincte violaceo-albido circumcinctis.

Exp. alar. unc. 2½.

Philippine Islands.

B.M.

42. *Mycalesis Madjicosa*, sp. nov. Pl. III. fig. 10.

Alæ supra velut in Ostrea, Westw. (Ostrea, Hübn.), ocellis autem anticarum vix iridatis, posticæque ocellis duobus analibus, externo minimo, interno permagno, similibus.

Alæ subtus fascia discali ad medium alarum magis approximante; linea submarginali ad marginem propiore, linea marginali tenuiore; margine has includente pallidior; aliter velut in Ostrea, var. (formæ communis).

Exp. alar. unc. 2½.

Madjico-Sima.

B.M.

43. *Mycalesis Mestra*.

Mycalesis Mestra, *Hewitson, Exot. Butterf.* iii. p. 79, pl. 40. figs. 2, 3 (1862).

East India, Deccan (*Hewitson*).

44. *Mycalesis Malsara*.

Mycalesis Malsara, *Hewitson, Exot. Butterf.* iii. p. 80, pl. 40.
figs. 5, 6 (1862).

Darjeeling (Hewitson).

Coll. Ind. Mus.

45. *Mycalesis Ostrea*.

Mycalesis Ostrea, *Westwood in Gen. Diurn. Lepid.* p. 394. n. 14
(1851).

Mycalesis Ostrea, *Hübner, Exot. Schmett. Beiträge*, figs. 79, 80
(1806); but not *P. Ostrea* of Cramer.

Sylhet.

B.M.

Var. Mycalesis Ostrea, *E. Doubleday, List Lep. Brit. Mus.* i. p. 140
(1844).

Sylhet, North India.

B.M.

46. *Mycalesis Sanatana*.

Mycalesis Sanatana, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*
i. p. 231. n. 489 (1857).

Darjeeling (Moore).

Coll. Ind. Mus.

47. *Mycalesis Perdiccas*.

Mycalesis Perdiccas, *Hewitson, Exot. Butterf.* iii. p. 84, pl. 42.
fig. 15 (1862).

Japan.

B.M.

48. *Mycalesis Oroatis*.

Mycalesis Oroatis, *Hewitson, Exot. Butterf.* iii. p. 90, pl. 45.
figs. 38, 39 (1864).

Java (Hewitson).

B.M.

Evidently allied to *Ostrea* of Hübner (not of Cramer).

49. *Mycalesis cinerea*.

Mycalesis cinerea, *Butler, Ann. & Mag. Nat. Hist.* xx. p. 401,
pl. 8. fig. 9 (1867).

Singapore.

Coll. Roberts.

50. *Mycalesis Hesione*.

Papilio Hesione, *Cramer, Pap. Exot.* i. pl. 11. figs. C, D (1779);
also of *Fabricius*.

Mycalesis Hesione, *Hübner, Verz. bek. Schmett.* p. 55. n. 525
(1816).

Satyrus Hesione, *Godart, Enc. Méth.* ix. p. 510. n. 98 (1819);
also of *Boisduval*.

Orsotrisena Hesione, Wallengren, *Nov. Gen. Lepid. Kongl. Vet. Akad. Forhandl.* p. 80 (1858).

Java, Sylhet.

B.M.

Var. DORIS.

Papilio Doris, Cramer, *Pap. Exot.* iv. pl. 362. fig. C (1782).

Papilio Medus, Fabricius, *Syst. Ent.* p. 488. n. 198 (1775).

China (Cramer), Borneo.

B.M.

Var. RUNEKA.

Mycalesis Runeka, F. Moore, *Cat. Lep. Mus. East-Ind. Comp.* p. 234. n. 501 (1857).

Assam (Moore), East India.

Coll. Ind. Mus. and B.M.

51. *Mycalesis mandata*.

Mycalesis mandata, F. Moore, *Cat. Lep. Mus. East-Ind. Comp.* p. 233. n. 499 (1857).

Mycalesis Gamaliba, Butler, *Ann. & Mag. Nat. Hist.* ser. 3. xix. pl. 2. figs. 3, 3a, 3b (1867).

Canara (Moore), Ceylon.

Coll. Ind. Mus. and B.M.

52. *Mycalesis Mandosa*, sp. nov. Pl. III. fig. 9.

Turbæ *Hesionice* maxima, supra velut præcedenti colorata; antice magis productæ, posticæ margine magis sinuato.

Alæ subtus obscuriores, margine externo squamis albidis rorato; ocellis velut in *Hesione* nec *mandata* positis; fascia media nivea lata, anticarum gradatim ad costam pyramidata, posticarum vero velut in *mandata*; ocellis anticarum fuscis valde indistinctis, posticarum nigris fusco cinetis et introrsum a lunulis plumbeis circumcinctis.

Exp. alar. unc. $2\frac{1}{8}$.

South India.

B.M.

53. *Mycalesis Nicotia*.

Mycalesis Nicotia, Hewitson, *Gen. Diurn. Lepid.* p. 394. n. 11, pl. 66. fig. 4 (1851); *Exot. Butterf.* iii. p. 79, pl. 40. fig. 1 (1862).

East India (Hewitson), South India.

B.M.

54. *Mycalesis Adolphe*.

Satyrus Adolphe, Guérin, *Souven. Voy. Inde*, p. 76 (1843).

Mycalesis Adolphe, Westwood in *Gen. Diurn. Lepid.* p. 394. n. 8 (1851).

Neelgherry Mountains.

According to Guérin very much like *Servatius* of Godart, but with-

out the violet gloss. From the description, it seems in some measure to resemble *Nicotia* of Hewitson, only the latter species has a large distinct subanal ocellus in the front wings instead of a white spot.

55. *Mycalesis Onatas*.

Mycalesis Onatas, *Hewitson, Exot. Butterf.* iii. p. 90, pl. 45. fig. 40 (1864).

Neelgherry Mountains.

Mr. Hewitson thinks this may be identical with the preceding. It is evidently closely allied to *Nicotia*.

56. *Mycalesis Martius*.

Papilio Martius, *Fabricius, Ent. Syst.* iii. pt. 1. p. 219. n. 686 (1793).

? *Satyrus Martius*, *Godart, Enc. Méth.* ix. p. 512. n. 102 (1819).

? *Mycalesis Martius*, *Westwood in Gen. Diurn. Lepid.* p. 393. n. 2 (1851).

Var. (also *anticæ subtus ocellis quinque*) *Mycalesis Moorei*, *Felder, Nov. Voy.* iii. tab. 67. fig. 9 (Sept. 1867).

Java.

(Typical form) B.M.

Fabricius appears to have counted the ocelli on his insect backwards. He describes the species as black above and unspotted; below the same colour, but with a paler margin, upon which, in the front wings, are four ocelli, in the hind wings seven; the first and third of the front, and the third of the hind wings larger. This description is exactly applicable to a species near the *Mamerta* of Mr. Moore's Catalogue.

57. *Mycalesis Gopa*.

Mycalesis Gopa, *Felder, Reise der Fregatte Novara*, iii. p. 501. n. 869 (1867).

Darjeeling (Felder).

Allied to *Nala*. Form of the hind wing the same as in *Diniche* of Hewitson, the *D. fuscum* of Felder.

58. *Mycalesis Nala*.

Mycalesis Nala, *Felder, Wien. ent. Monatschr.* iii (1859); also in *Lep. Fragm.* n. 46 (1859).

Mycalesis Mamerta, *Moore, Cat. Lep. East-Ind. Mus.* i. p. 231. n. 490 (1851).

Var. *Mycalesis Sudra*, *Felder, Reise der Fregatte Novara*, iii. p. 500. n. 868, tab. 67. fig. 10 (1867).

Java.

The true *Mamerta* of Cramer appears to me to be nearly allied to, if not a local form of, *Malsara* of Moore. It seems chiefly to differ from the latter in the size and number of the ocelli on the underside.

59. ***Mycalesis Anaxias*.**

Mycalesis Anaxias, *Hewitson, Exot. Butterf.* iii. p. 86, pl. 43. figs. 25, 26 (1862).

South India.

B.M.

60. ***Mycalesis Æthiops*, sp. nov.** Pl. III. fig. 11.

Alæ supra olivaceo-nigræ, ocellis velut in *Remulia*.

Alæ subtus area apicali violacea, ocellos velut in *Asophide* ♀ includente; area hac apud costam anticarum introrsum directa, fusco rorata; linea submarginali undata altera marginali undulata et margine ipso nigris; linea ad marginem quoque flava: posticæ area basali olivaceo tincta, a linea undulata intersecta.

Exp. alar. unc. $2\frac{1}{8}$.

New Guinea.

B.M.

61. ***Mycalesis Phidon*.**

Mycalesis Phidon, *Hewitson, Exot. Butterf.* iii. p. 84, pl. 42. fig. 16 (1862).

Aru Islands.

B.M.

62. ***Mycalesis Dorycus*.**

Satyrus Dorycus, *Boisduval, Voy. Astrolabe*, p. 152. n. 13 (1832).

Cænonympha? *Dorycus*, *Westwood in Gen. Diurn. Lepid.* p. 398. n. 21 (1851).

♀. *Mycalesis Dorycus*, *Hewitson, Exot. Butterf.* iii. pl. 41. figs. 7, 10 (*Doricus*), p. 81 (1862).

♂. *Satyrus Duponchellii*, *Gutrin, Voy. Coquille*, pl. 17. fig. 3 (1829-32).

♂. *Mycalesis Getulia*, *Felder, Wien. entom. Monatschr.* iii. p. 269 (1859).

Dorey (New Guinea).

B.M.

63. ***Mycalesis Mnasicles*.**

Mycalesis Mnasicles, *Hewitson, Exot. Butterf.* iii. p. 88, pl. 44. figs. 32, 33 (1864).

Sumatra (Hewitson).

This species must come near *Dorycus* of Boisduval.

64. Mycalesis Dexamenus.

Mycalesis Dexamenus, *Hewitson, Exot. Butterf.* iii. p. 83, pl. 42.
figs. 17, 18 (but not 19) (1862).

Celebes.

B.M.

65. Mycalesis Dinon.

Mycalesis Dinon, *Hewitson, Exot. Butterf.* iii. p. 88, pl. 44.
fig. 31 (1864).

Mycalesis Dexamenus (part.), *Hewitson, Ex. Butterf.* iii. pl. 42.
fig. 19 (1862).

Celebes.

B.M.

66. Mycalesis Deianira.

Mycalesis Deianira, *Hewitson, Exot. Butterf.* iii. p. 83, pl. 42.
fig. 13 (1862).

Mycalesis Dora, *Hewitson, Exot. Butterf.* iii. p. 88 (1864).

Tondano (Hewitson).

Coll. Wallace.

Closely allied to *Dexamenus* and *Dinon*, but differing in the sub-marginal lines of the underside. Mr. Hewitson altered the name from *Deianira* to *Dora*, as he considered the former too much like *Dejanira* *, the *i* and *j* being synonymous on the continent. *Dejanira*, however, belongs to another genus.

67. Mycalesis Mucia.

Mycalesis Mucia, *Hewitson, Exot. Butterf.* iii. p. 82, pl. 41.
figs. 11, 12 (1862).

Aru Islands.

B.M.

Allied to *Cyamites* of Boisduval. The female on the upperside is somewhat like *Dinon* of Hewitson.

68. Mycalesis Cyamites.

Satyrus Cyamites, *Boisduval, Vog. de l'Astrolabe*, pt. 1. p. 152
(1832).

Cænonympha Cyamites, *Westwood in Gen. Diurn. Lepid.* p. 398.
n. 22 (1851).

Satyrus Bazochii, *Gutrin, Voy. Coquille, Atlas, Ins.* pl. 14². fig. 3
(1832).

Dorey, New Guinea.

69. Mycalesis Messene.

Mycalesis Messene, *Hewitson, Exot. Butterf.* iii. p. 81, pl. 41.
figs. 8, 9 (1862).

Ternate.

♂, B.M.

* *Pararge Dejanira*.

70. **Mycalesis Shiva.**

Satyrus Shiva, *Boisduval, Voy. de l'Astrolabe, Entom.* pt. 1.
p. 149. n. 9 (1832).

Cænonympha Shiva, *Westwood in Gen. Diurn. Lepid.* p. 398.
n. 25 (1851).

Dory (New Guinea).

Coll. Hewitson.

This species is allied to *Messene*.

71. **Mycalesis Tagala.**

Mycalesis Tagala, *Felder in Wien. ent. Monatschr.* v. (1861);
also in *Lep. Diagn.* n. 101; *Nov. Voy.* iii. tab. 67. figs. 7, 8
(1867).

Bourias, Locban, Mindanao (Felder).

Seems to be allied to *Mycalesis Mucia* of Hewitson, but may come
into the *Maianaeas* group.

72. **Mycalesis Bisaya.**

Mycalesis Bisaya, *Felder in Wien. ent. Monatschr.* v. (1861);
also in *Lep. Diagn.* n. 102.

Bourias, Locban, Luzon (Felder).

According to Felder, it is quite possible that this may turn out to
be a variety of the preceding species.

73. **Mycalesis Mehadeva.**

Satyrus Mehadeva, *Boisduval, Voy. de l'Astrolabe, Entom.* pt. 1.
p. 151. sp. 12 (1832).

Cænonympha Mehadeva, *Westwood, Gen. Diurn. Lepid.* p. 398.
n. 23 (1851).

Dorey (New Guinea).

B.M.

In our specimen of this insect there is a seventh very minute anal
ocellus in the hind wings. It is allied to *Asopis* of Hewitson. The
specimens of *Mehadeva* in Mr. Hewitson's Collection have a greater
extent of black colour in the front wings than exists in our specimen.

74. **Mycalesis Remulia.**

Papilio Remulia, *Cramer, Pap. Exot.* iii. pl. 237. figs. F, G
(1782); but not of *Godart & Boisduval*.

Mycalesis? *Remulia*, *Westwood in Gen. Diurn. Lepid.* p. 394.
n. 17 (1851).

Java (Cramer), North Ceram.

B.M.

75. **Mycalesis Terminus.**

Papilio Terminus, *Fabricius, Ent. Syst.* iii. pt. 1. p. 220. n. 687
(1793); *Donovan, Ins. New Holl.* pl. 28. fig. 4 (1805).

Satyrus Terminus, *Godart, Enc. Méth.* ix. p. 501. n. 84 (1819).

Mycalesis? *Terminus*, *Westwood in Gen. Diurn. Lepid.* p. 394. n. 10 (1851); *Butler, P. Z. S.* p. 720. figs. 3, 3a (1867).

New Holland, Australia.

B.M.

Certainly not a variety of *Remulia* of Cramer, from which it differs in the position of the subanal ocellus of the fore wing, in the smaller and more regular ocelli, paler colouring, and waved submarginal line of the underside. The type of this species is in the Banksian Collection.

76. *Mycalesis Minois*.

Mycalesis Minois, *Hewitson, Exot. Butterf.* iii. p. 87, pl. 44. figs. 29, 30 (1864).

Timor.

77. *Mycalesis Asophia*.

Mycalesis Asophia, *Hewitson, Exot. Butterf.* iii. p. 85, pl. 43. figs. 20, 21 (1862).

(♂) Ternate, (♀?) Mysol.

B.M.

78. *Mycalesis Felderi*, sp. nov. Pl. III. fig. 5.

Alæ supra fuscae: anticæ apice velut in *Remulia* nigrescente; ocello permagno discali nigro albo pupillato et ferrugineo cincto; puncto subapicali minimo albo: posticæ ocellis septem, quinto maximo, secundo, tertio et septimo minimis, omnibus nigris albo pupillatis flavo cinctis et nigro circumcinctis; fascia submarginali et ciliis fuscis obscurioribus, margine ipso nigro.

Alæ subtus fere velut in Ita (Feld.), ocellis autem velut supra.

Exp. alar. unc. 2½.

Philippines.

B.M.

This may be only an extreme form of *Ita*.

79. *Mycalesis fuscum*.

Dasyomma fuscum, *Felder, Wien. ent. Monatschr.* iv. p. 401. n. 27 (1860).

Mycalesis Diniche, *Hewitson, Exot. Butterf.* iii. p. 85, pl. 43. fig. 23 (1862).

Malay peninsula (Felder), Singapore.

B.M.

80. *Mycalesis Sirius*.

*Papilio Sirius**, *Fabricius, Syst. Ent.* p. 488. n. 201 (1775); *Donovan, Ins. New Holl.* pl. 28. fig. 1 (1808).

* The type is in the Banksian Collection.

Satyrus Sirius, *Godart, Enc. Méth.* ix. p. 500. n. 83 (1819).
Cænonympha Sirius, *Westwood in Gen. Diurn. Lepid.* p. 398.
 n. 26 (1851).

New Holland, Australia.

♂, B.M.

Var. ZACHÆUS.

Papilio Zachæus, *Fabricius, Ent. Syst.* iii. pt. 1. p. 217. n. 679
 (1793).

Satyrus Zachæus, *Godart, Enc. Méth.* ix. p. 512. n. 103 (1819).
Lasiommata? *Zachæus*, *Westwood in Gen. Diurn. Lepid.* p. 387.
 n. 17 (1851).

Hab. — ? (Fabr.), Australia.

♂ ♀, B.M.

Var. DAIDIS.

Mycalesis Daidis, *Hewitson, Exot. Butterf.* iii. p. 85, pl. 43.
 fig. 22 (1862).

Mycalesis Manipa (*syn.*), *Felder, Lep. Diagn.* p. 46. n. 100, note.

Ceram (Hewitson), var. North Ceram.

B.M.

Another form of *M. Sirius*.

Var. MANIPA.

Satyrus Manipa, *Boisduval, Voy. de l'Astrolabe, Entom.* pt. 1.
 p. 150. n. 10 (1832).

Cænonympha Manipa, *Westwood in Gen. Diurn. Lepid.* p. 398.
 n. 24 (1851).

"Amboyna, Bourou, Offack," Australia.

B.M.

Scarcely differs from *Daidis*. Dr. Felder considers it identical with it.

81. *Mycalesis Ita.*

Mycalesis Ita, *Felder in Wien. ent. Monatsch.* v. (1861); also
 in *Lep. Diagn.* n. 100; *Nov. Voy.* iii. tab. 68. figs. 8, 9
 (1867).

Luzon, Philippines.

B.M.

82. *Mycalesis Itys.*

Mycalesis Itys, *Felder, Reise der Fregatte Novara*, iii. p. 903.
 n. 873 (1867).

Celebes (Felder).

83. *Mycalesis fraterna*, sp. nov. Pl. III. fig. 13.

Alæ supra velut in *Narcisso*, sed majores.

Alæ subtus pallidiores, lineis ocellisque magis distinctis, linea media
 externa posticarum leviter undata; ocello discali anticarum rufo cir-

cumcincto; ocellis posticarum majoribus, uno subapicali; aliter velut in *Narcisso*.

Exp. alar. unc. $1\frac{1}{4}$.

Madagascar.

B.M.

84. *Mycalesis Narcissus*.

Papilio Narcissus, *Fabricius, Ent. Syst. Suppl. v. p. 428. n. 672, 673 (1793).*

Satyrus Narcissus, *Godart, Enc. Méth. ix. p. 551. n. 181 (1819);*
also of *Boisduval*.

Ctenonympha Narcissus, *E. Doubleday, List Lep. Brit. Mus. App. p. 33 (1847).*

Mycalesis Narcissus, *Westwood in Gen. Diurn. Lepid. p. 394. n. 9 (1851).*

Mauritius.

B.M.

85. *Mycalesis Anapita*.

Mycalesis Anapita, *Moore, Cat. Lep. Mus. East-Ind. Comp. i. p. 232. n. 495 (1857).*

Borneo, India.

Coll. Ind. Mus. and, ♂ ♀, B.M.

86. *Mycalesis Patnia*.

♂. *Mycalesis Patnia*, *F. Moore, Cat. Lep. Mus. East-Ind. Comp. p. 232. n. 494 (1857).*

Ceylon.

Coll. Ind. Mus. and B.M.

87. *Mycalesis Junonia*, sp. nov. Pl. III. fig. 4.

♂ ♀. Alæ supra fuscae, margine pallido lineam nigram includente, ciliis griseis fusco variis: anticae ocellis duobus nigris albo pupillatis, superiore brunneo cincto; inferiore multo majore, insuper a lunula irregulari albida cincto: posticae ocellis uno duobusve indistinctis discalibus minimis.

Alæ subtus olivaceae, aliter fere velut in *Patnia* scriptae.

Exp. alar. unc. $1\frac{1}{8}$ – $1\frac{1}{4}$.

South India.

B.M.

This is clearly allied to *Patnia*, but quite distinct.

Genus 46. ERITES, *Westwood (1851).*

Typical Species *Erites Madura*.

1. *Erites Madura*.

Hipparchia Madura, *Horsfield, Cat. Lep. Mus. E.-I. C. tab. 5. figs. 8, 8 a (1829).*

Satyrus (*Erites*) *Madura*, *Westwood in Gen. Diurn. Lepid.* p. 392. n. 47 (1851).

Erites Madura, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i. p. 229. n. 484 (1857).

Java.

Horsf. Coll. and B.M.

2. *Erites elegans*, sp. nov. PL. II. fig. 4.

Alæ supra fuscæ subpellucidæ, marginibus obscurioribus: posticæ lineis duabus marginalibus ochreis; ocellis sex submarginalibus, quinto permagno, aliis minimis, nigris ochreo cinctis; fascia discali undata ochracea.

Alæ subtus margaritacæ, area basali et margine interno fusco liturata; fascia obliqua communi integra, altera discali, posticis undata, et tertia posticis inter ocellos pallidiore ochreis, fusco limitatis; lineis duabus (interna interrupta) marginalibus et margine ipso fuscis; margine has includente lacteo: anticæ fascia submarginali ocellos quinque subæquales ferente, his fuscis lacteo cinctis, et albo pupillatis: posticæ ocellis sex nigris, sexto indistinctissimo, quinto permagno, aliis ochreo cinctis argenteo pupillatis et ferrugineo-ochreo late circumcinctis, quinto pupilla permagna.

Exp. alar. unc. $2\frac{1}{4}$.

Borneo (two specimens).

B.M.

This is a very distinct and beautiful species.

Genus 47. *CÆNYRA*, *Hewitson* (1865).

Typical Species *Cænryra Hebe*.

1. *Cænryra Hebe*.

Ypthima Hebe, *Trimen, Trans. Ent. Soc.* 3rd ser. i. p. 280 (1864); *Rhop. Afric. Austral.* ii. p. 205. n. 118, pl. 4. fig. 3 (1866).

Var. Cænryra Hebe, *Hewitson, Trans. Ent. Soc.* 3rd ser. ii. p. 281 (1865).

Cænryra Corycia, *Hewits. ibid.* pl. 17. figs. 1, 2 (1865).

Bashee River (Trimen), Port Natal. Colls. Hewitson and B.M.

Genus 48. *XOIS*, *Hewitson* (1865).

Typical Species *Xois Sesara*.

1. *Xois Sesara*.

Xois Sesara, *Hewitson, Trans. Ent. Soc.* 3rd ser. ii. pt. 4. p. 282, pl. 17. figs. 3, 4 (1865).

Fiji Islands.

Colls. Hewitson and B.M.

Genus 49. *YPITHIMA*, Westwood [not Hübner*] (1851).

Typical Species *Ypthima* (Baldus) *Lisandra*.

1. *Ypthima Asterope*.

Hipparchia Asterope, Klug in Ehrenb. & Hempr. *Symb. Phys.* pl. 29. figs. 11-14 (1829-45).

Ypthima (sic) *Asterope*, Hewitson in *Trans. Ent. Soc. London*, 3rd ser. ii. p. 283. n. 1 (1865).

Var. *Ypthima Norma*, Westwood & Hewitson in *Gen. Diurn. Lepid.* p. 395. n. 5, pl. 67. fig. 1 (1851).

Java (B.M.), Syria [Natal and Calabar] (Hewitson), var. China. B.M.

There can be no doubt that Mr. Hewitson is right in sinking *Y. Norma* as a variety of *Asterope*.

2. *Ypthima Nareda*.

Satyrus Nareda, Kollar in Hugel's *Kaschmir*, p. 451. n. 12 (1848).

Ypthima Nareda, Hewitson in *Trans. Ent. Soc. Lond.* 3rd ser. ii. p. 284. n. 6, pl. 17. figs. 6, 7 (1865).

Northern India, (var.) Borneo. B.M.

Local form. *Ocello apicali anticarum iride magna*.

South and West Africa. B.M.

Var. *MOTSCHULSKII*.

Satyrus Motschulskii, Bremer, *Beitr. zur Fauna nördl. Chinas*, p. 8. n. 26 (1853); *Ménétriés, Enum. Corp. Anim.* pt. 1. p. 47. n. 782, t. 6. fig. 5 (1855).

Ypthima Motschulskiji (sic), Hewitson, *Trans. Ent. Soc. London*, 3rd ser. ii. p. 290. n. 17 (1865).

(Var.) Northern India. B.M.

3. *Ypthima Loryma*.

Ypthima Loryma, Hewitson, *Trans. Ent. Soc. London*, 3rd ser. ii. p. 289. n. 14, (*Ypthima*) pl. 19. figs. 16, 17 (1865).

Macassar (Hewitson), Celebes. B.M.

4. *Ypthima amphithea*.

Ypthima amphithea, *Ménétriés in Schrenck's Amur-Lande*, pl. 3. fig. 10, p. 41 (1859).

* The type of Hübner's *Ypthima* is the *cassus* of Linnæus, recently made into a genus by Wallengren. As it would create great confusion to alter the name of the present group, it is preferable to retain it as representing the *Ypthima* of Westwood.

Ypthima amphitheia, *Hewitson, Trans. Ent. Soc.* 3rd ser. ii. p. 289. n. 15 (1865).

Amoor, Sula.

Coll. Hewitson.

This species seems to differ very little from suffused specimens of *Nareda*. Mr. Hewitson, in his excellent monograph of the genus, remarks that it chiefly differs from *Loryma* in colour.

5. *Ypthima Pandocus*.

Ypthima Pandocus, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*

i. p. 235. n. 506 (1857); *Hewitson, Trans. Ent. Soc. Lond.* 3rd ser. ii. p. 290. n. 16, pl. 18. fig. 12 (1865).

Java.

♂, B.M.

Var. Anticæ ocello multo minus oblique : posticæ magis elongatæ ; subtus ocellis quinque, duobus apicalibus et tribus analibus, primo et quarto maximis.

Borneo.

♀, B.M.

This species links the *Nareda* and *Baldus* groups together.

6. *Ypthima Batesii*.

Ypthima Batesii, *Felder, Reise der Fregatte Novara*, iii. tab. 68.

figs. 10, 11 (1867).

Madagascar (Felder).

Allied to *Ypthima Pandocus* of Moore.

7. *Ypthima Sakra*.

Ypthima Sakra, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i.

p. 236. n. 508 (1857); *Hewitson, Trans. Ent. Soc. London*, 3rd ser. ii. p. 290. n. 18, pl. 18. fig. 18 (1865).

North India.

Coll. Hewitson.

8. *Ypthima Methora*.

Ypthima Methora, *Hewitson, Trans. Ent. Soc. London*, 3rd ser.

ii. p. 291. n. 19, (*Ypthima*) pl. 18. figs. 20, 21 (1865).

North India.

B.M. ? Coll. Hewitson.

9. *Ypthima Lisandra*.

Papilio Lisandra, *Cramer, Pap. Exot.* iv. tab. 293. figs. G, H (1782).

Ypthima Lysandra, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.*

i. p. 235. n. 504 (1857); also of *Hewitson* (1865).

China.

B.M.

Var. BALDUS.

Papilio Baldus, *Fabricius, Mant. Ins.* ii. p. 34. n. 356 (1787);
also of *Donovan* (figure very bad). Coll. Banks.

Satyrus Baldus, *Godart, Enc. Méth.* ix. p. 551. n. 184 (1819).

Ypthima Baldus, *Westwood in Gen. Diurn. Lepid.* p. 395. n. 1
(1851), type of genus.

Ypthima Baldus, *Hewitson, Trans. Ent. Soc. Lond.* 3rd ser. ii.
p. 286. n. 9 (1865).

India (*Fabricius*), Hongkong, China.

B.M.

The type specimen of this insect has a subapical ocellus on the upperside of the hind wings: there is not sufficient difference between it and *Lisandra* to constitute a species.

Var. LARA.

Papilio Lara, *Donovan, Nat. Repos.* ii. pl. 71 (1824).

Ypthima laroides, *Westwood in Gen. Diurn. Lepid.* p. 395.
n. 6 (1851).

South Africa, Cape of Good Hope (*Donovan* and *Westwood*).

This form seems more distinct than *Baldus* from *Lisandra*. As there are no African specimens of this species in the collection, it is impossible to say whether or no *Lara* is identical in all respects with it.

10. Ypthima Sepyra.

Ypthima Sepyra, *Hewitson, Trans. Ent. Soc. London*, 3rd ser.
p. 292. n. 22, (*Ypthima*) pl. 17. fig. 10 (1865).

Gilolo, Batchian.

Coll. Hewitson.

11. Ypthima Argus.

Ypthima Argus (*sic*), *Butler, Journ. Linn. Soc. (Zool.)* ix.
p. 56. n. 22.

Hakodadi, North Japan.

♂ ♀, B.M.

This species chiefly differs from *Lisandra* in the different ground-colour, the clearer ocelli, and in the addition of a curious ocelloid spot at the termination of the front-wing cell on the underside in the female.

12. Ypthima Itonia.

Ypthima Itonia, *Hewitson, Trans. Ent. Soc. London*, 3rd ser.
ii. p. 287. n. 11, (*Ypthima*) pl. 18. fig. 13 (1865).

White Nile.

B.M.

13. Ypthima Stellera.

Hipparchia Stellera, *Eschscholtz, Kotzeb. Voy.* pl. x. figs. 24a,
24b, p. 216 (1821).

- Ypthima Stellera*, *Hewitson, Trans. Ent. Soc. London*, p. 285. n. 8 (1865).
Var. Ypthima Sempera, *Felder, Wien. ent. Monatschrift*, iii. p. 125 (1863).
Manilla (Eschscholtz). Coll. Hewitson and B.M.
14. *Ypthima fasciata*.
Ypthima fasciata, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. p. 287. n. 12 (1865).
 Sarawak, Sumatra (Hewitson).
15. *Ypthima Chenu*.
Satyrus Chenu, *Gutrin-Méneville in Delessert's Souv. Voy. Ind.* pl. 21. fig. 2, p. 77 (1843).
 Australia. B.M.
16. *Ypthima Inica*.
Ypthima Inica, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. ii. p. 284. n. 5, (*Ypthima*) pl. 17. fig. 5 (1865).
 East India. B.M.
17. *Ypthima Arctous*.
Papilio Arctous, *Fabricius, Mant. Ins.* ii. p. 33 (1785).
Satyrus Arctous, *Godart, Enc. Méth.* ix. p. 552. n. 185 (1819); also of *Boisduval*.
Ypthima Arctous, *Westwood in Gen. Diurn. Lepid.* p. 395. n. 7 (1851).
Ypthima Arctous, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. p. 283. n. 2 (1865).
 Aru, Port Denison (Hewitson), New Holland.
 Coll. Banks and B.M.
- Var. ?? ARCTOIDES*.
Ypthima arctoides, *Hewitson in Trans. Ent. Soc. London*, p. 284. n. 3 (1865).
Papilio Arctous (*Irius in plate*), *Donovan, Ins. New Holland*, pl. 24. fig. 2 (1805).
 New Holland (Donovan).

The figure by Donovan must either be incorrect or not copied from the Fabrician type: he distinctly states that his insect is from the Banksian cabinet; and an examination of this type shows that it in no way differs from the usual form of *Arctous*. It seems tolerably evident that the extra ocellus in Donovan's figure is merely an addition of his own, in which case *arctoides* will become a synonym of *Arctous*.

18. Ypthima Philomela.

Ypthima Philomela, *Hübner, Zuträge Exot. Schmett.* figs. 83, 84 (1806); *Verz. bek. Schmett.* (part.) p. 63. n. 608 (1816); but not *Philomelus* of *Linnaeus*, p. 768.

Ypthima Philomela, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. ii. p. 284. n. 4 (1865).

North India.

B.M.

Var. Ocellis posticarum *subtus obsoletis.*

East Indies.

B.M.

19. Ypthima Ceylonica.

Ypthima Ceylonica, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. ii. p. 288. n. 13, (*Ypthima*) pl. 18. figs. 14, 15 (1865).

Ceylon.

B.M.

20. Ypthima Hyagriva.

Ypthima Hyagriva, *Moore, Cat. Lep. Mus. East-Ind. Comp.* p. 236. n. 507 (1857); *Hewitson, Trans. Ent. Soc. London*, 3rd ser. ii. p. 291. n. 20, (*Ypthima*) pl. 18. fig. 11 (1865).

Darjeeling.

Coll. Hewitson.

21. Ypthima Narasingha.

Ypthima Narasingha, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* p. 236. n. 509 (1857); *Hewitson, Trans. Ent. Soc. London*, p. 291. n. 21, (*Ypthima*) pl. 18. fig. 19 (1865).

Darjeeling.

Coll. Hewitson.

22. Ypthima Aphnius.

Satyrus Aphnius, *Godart, Enc. Méth.* ix. p. 551. n. 183 (1819).

Ypthima Aphnius, *Westwood in Gen. Diurn. Lepid.* p. 396. n. 10 (1851).

Ypthima Aphnius, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. p. 292. n. 23 (1865).

Var. ? Papilio Magus, *Fabricius, Ent. Syst.* iii. pt. 1. p. 223. n. 700 (1793).

Timor.

B.M.

It is better to regard *Tamatavæ* as a *Pseudonympha*. Its accidental resemblance to *Y. Aphnius* is not a sufficient reason for placing it in *Ypthima*; and the general colouring is more like that of the species of *Pseudonympha*. It seems nearly allied to *Cassius* of Godart.

Genus 50. ORESSINOMA, *Westwood* (1851).Typical Species *Oressinoma Typhia*.1. *Oressinoma Typhia*.*Oressinoma Typhia*, *Hewitson, & Westwood in Gen. Diurn. Lepid.* p. 372. n. 1, pl. 62. fig. 5 (1851).

Venezuela, Bolivia, New Granada, West Coast.

B.M.

Genus 51. ARGE, *Hübner* (1816)*.Typical Species *Arge Syllius* (*Psyche*).1. *Arge Galathea*.*Papilio Galathea*, *Linnaeus, Syst. Nat.* i. p. 474 (1758), ii. p. 772. n. 147 [12th ed.] (1766); *Hübner, Eur. Schmett.* i. pl. 41. figs. 183–185 (1805); also of *Scopoli, Denis, &c.**Papilio* (Nymph. Gemm.) *Galathea*, *Esper, Schmett.* i. pt. 1. tab. 7. fig. 3, tab. 25. *Suppl.* i. fig. 1 (1777).*Satyrus Galathea*, *Godart, Enc. Méth.* ix. p. 505. n. 91 (1819); *Lép. de France*, i. p. 165. n. 55, pl. 8. fig. 2 (1821).*Hipparchia Galathea*, *Stephens, Ill. Haust.* i. p. 57. sp. 7 (1828).*Arge Galathea*, *Boisduval, Sp. Gén. Lép. Ind.* p. 25, n. 185 (1840).*Maniola Galathea* (sic), *Schrank, Fauna Boica*, ii. pt. 1. p. 170. n. 1297 (1801).*Papilio Galathea* (sic), *Ochsenheimer, Schmett. v. Eur.* i. p. 242. n. 35 (1807).*Var. Papilio* (Nymph. Gemm.) *Galaxera*, *Esper, Schmett.* i. pt. 2. tab. 111. cont. 66. fig. 5 (1777).

Italy, France, England.

B.M.

*Var. PROCIDA.**Papilio Procida*, *Herbst, Natursyst. Schmett.* viii. p. 22. n. 6 (1796).*Arge Procida*, *Boisduval, Icon. Hist. Lép.* p. 131. n. 1, pl. 25. figs. 5, 6 (1832).*Papilio* (Hipparchia) *Galathea*, *var. Procida*, *Freyer, Neuere Beitrage*, iv. tab. 379, (*Galathea*) p. 157. n. 649 (1842).*Var. Papilio* (Nymph. Gemm.) *Galathea*, *var.*, *Esper, Schmett.* i. pt. 2. tab. 111. cont. 66. fig. 4 (1777) [B.M.].

Italy, Germany, South Europe.

B.M.

*Var. LEUCOMELAS.**Papilio* (Nymph. Gemm.) *leucomelas*, *Esper, Schmett.* i. pt. 2. tab. 81. cont. 31. figs. 1, 2 (1777).* The *Arge* of *Esper* is a specific, and not a generic name.

Papilio leucomelas, *Borkhausen, Naturgesch. Schmett.* i. p. 106. n. 49 (1788); *Hübner, Eur. Schmett.* i. pl. 102. figs. 517, 518 (1805).

Arge leucomelas, *Boisduval, Icon. Hist. Léop.* p. 133. n. 2, pl. 25. figs. 3, 4 (1832).

Papilio (*Hipparchia*) *leucomelas*, *Freyer, Neuere Beiträge*, v. p. 87. n. 767, pl. 433 (1845).

Papilio Liriope, *Cyrrilli Ent. Neap. Spec.* i. tab. 12. fig. 3 (1787).

Var. Arge Clotho?, *Lucas, Expl. Algér. Léop.* pl. 1. figs. 4, 4a (1849).

Pyrenees.

B.M.

2. *Arge meridionalis*.

Melanargia Halimede, *var. meridionalis*, *Felder, Lep. Chin. et Jap., in Wien. ent. Monatschr.* p. 27.

North China.

B.M.

This is much more worthy of a specific name than the *Halimede* of Ménétriés; it appears to be quite distinct from the latter, which seems to be only a race of *Lachesis*.

3. *Arge Lachesis*.

Papilio Lachesis, *Hübner, Beiträge Gesch. Schmett.* ii. pt. 3. p. 70, pl. 3. figs. 1, 2 (1786-90); also of *Herbst* and *Ochsenheimer*.

Arge Lachesis, *Boisduval, Sp. Gén. Léop. Ind.* p. 25. n. 183 (1840); also of *Doubleday* (1844).

Papilio (*Nymph. Gemm.*) *Arge* (*Nemausiaca*), *Esper, Schmett.* i. pt. 2. tab. 96. cont. 51. figs. 1, 2, and pt. 2. p. 15 (1777).

Var. Satyrus Lachesis, *Godart, Enc. Méth.* ix. p. 507. n. 92 (1819); *Léop. de France*, ii. p. 135. n. 51, pl. 19. figs. 1, 2 (1822); also of *Lefebvre*.

France.

B.M.

Var. HALIMEDE.

Arge Halimede, *Schrenck, Reise und Forsch. Amur-Lande*, Bd. ii. p. 37. n. 71, tab. 3. figs. 6, 7 (1859).

Amur Land.

Var. TITEA.

Hipparchia Titea, *Klug, Symb. Phys.* tab. 29. figs. 15-18 (1829-45).

Arge Titea, *Boisduval, Sp. Gén. Léop. Ind.* p. 26. n. 189 (1840).

Satyrus D'Arceti, *Lefebvre, Ann. Ent. Soc. France*, i. p. 89, pl. 2. fig. 1 (1832-34); *Duponchel, Léop. de France*, i. p. 174. n. 58, pl. 26. figs. 5, 6 (1832).

Hipparchia Teneates, *Ménétriés, Cat. rais. Ins. Caucas.* p. 255.

n. 1200 (1832); *Herrich-Schäffer, Eur. Schmett. i. figs. 423, 424* (1845).

Syria (Klug), Mount Lebanon (Lefebvre), Caucasus (Ménétriés).

A. Halimede and *Titea* must be looked upon as merely local forms of *A. Lachesis*; they differ about as much as *Galathea* and *Pro-cida* do.

4. *Arge Hylata*.

Hipparchia Hylata, *Ménétriés, Cat. rais. Ins. Caucas. p. 251. n. 1198* (1832); *Herrich-Schäffer, Eur. Schmett. i. figs. 425, 426* (1845).

Caucasus (Ménétr.), Persia.

B.M.

This species is exceedingly variable in pattern and coloration; it is not clear that it is more than a local form of *A. Larissa*.

5. *Arge Larissa*.

Papilio Larissa, *Hübner, Eur. Schmett. i. pl. 182. figs. 896-899* (1805).

Papilio (*Hipparchia*) *Larissa*, *Freyer, Neuere Beiträge, ii. p. 85, tab. 73. fig. 1* (1836).

Arge Larissa, *Boisduval, Icon. Hist. Léop. p. 136, pl. 28. figs. 4-6* (1832).

Dalmatia.

B.M.

Var. HERTA.

Papilio Herta, *Hübner, Eur. Schmett. i. pl. 183. figs. 900-903* (1805).

Arge Herta, *Boisduval, Icon. Hist. Léop. p. 134. n. 3, pl. 28. figs. 1-3* (1832); also of *Lefebvre* (1832-34).

North Europe.

B.M.

A local form of *A. Larissa*.

6. *Arge Iapygia*.

Papilio (*Nymph. Gemm.*) *Iapygia*, *Esper, Schmett. i. pt. 2. tab. 105. cont. 60. fig. 3, tab. 111. cont. 66. fig. 3* (1777).

Papilio Iapygia, *Cyrrilli Ent. Neap. Spec. i. tab. 3. fig. 5* (1787); *Herbst* (1796), as *Galathea*, *var.*

Papilio Arge, *Fabricius, Ent. Syst. iii. pt. 1. p. 239. n. 746* (1793).

Satyrus Arge, *Godart, Enc. Méth. ix. p. 507. n. 93* (1819).

Papilio Clotho, *Hübner, Eur. Schmett. i. pl. 42. figs. 190, 191* (1805); also of *Ochsenheimer*.

Arge Clotho, *Boisduval, Icon. Hist. Léop. p. 137. n. 5, pl. 25. figs. 1, 2* (1832); also of *Hewitson* (1851), but not of *Lucas*.

Var. Papilio (*Nymph. Gemm.*) *Arge* (*Russiae*), *Esper, Schmett. i. pt. 2. tab. 84. cont. 34. fig. 1, ? 2* (1777).

Papilio Suwarovius, *Herbst, Natursyst. Schmett.* viii. p. 13. n. 2, tab. 182. figs. 5, 6, ? 7 (1796).

Var. Papilio Atropos, *Hübner, Eur. Schmett.* i. pl. 43. figs. 192, 193 (1805).

Papilio Lyssianassa, *Dahl* (1823?).

Arge Lyssianassa, *Boisduval, Sp. Gén. Lép. Ind.* p. 25 (1840).

Arge Clotho, var. *Atropos*, *Bellier de la Chavignerie, Ann. Ent. Soc. France*, p. 677. n. 1 (1860).

Sicily (Bellier), South Europe, Germany.

B.M.

Var. CLEANTHE.

Arge Cleanthe, *Boisduval, Icon. Hist. Lép.* p. 139. n. 6, pl. 26. figs. 1-3 (1832).

Satyrus Clotho, var. *Cleanthe*, *Duponchel, Lép. de France*, i. p. 170, pl. 25. figs. 5, 6 (1832).

Pyrenees, South of France.

B.M.

Var. CLOTHO.

Satyrus Clotho, *Duponchel, Lép. de France*, i. p. 167. n. 56. figs. 1-4 (1832); but not of *Hübner*.

Var. Papilio (Hipparchia) Xenia, *Freyer, Neuere Beiträge*, vi. p. 133. n. 1030, pl. 566. figs. 3, 4 (1852).

Arge Hertha, var. *caucasica*, *Nordmann, Bull. Soc. Imp. Nat. Mosc.* p. 403, tab. 8. figs. 1-3 (1851).

Russia.

B.M.

This form is more distinct than *Cleanthe*; they are both evidently local forms of *Iapygia*. We have a single specimen of an *Arge* which seems intermediate between *Herta* and *Iapygia*; but it would be rash to venture, without clearer evidence, to consider these species mere variations of one type.

7. *Arge Ines.*

Papilio Ines, *Hoffmannsegg in Illig. Mag.* iii. p. 205 (1804).

Arge Ines, *Boisduval, Icon. Hist. Lép.* p. 145. n. 10, pl. 27. figs. 5, 6 (1832); also of *Lefebvre* and *Westwood*.

Satyrus Ines, *Duponchel, Lép. de France*, i. p. 162. n. 54, pl. 24. figs. 1, 2, (var.) 3, 4 (1832).

Papilio Thetis, *Hübner, Eur. Schmett.* i. pl. 43. figs. 196, 197 (1805).

Papilio Amphitrite, var., *Ochsenheimer, Schmett. v. Eur.* i. pt. 1. p. 252 (1807), ? var. pt. 2. pp. 236, 237 (1808).

Spain.

B.M.

8. *Arge Syllius.*

Papilio Syllius, *Herbst, Natursyst. Schmett.* viii. p. 15, tab. 182. figs. 8, 9 (1796); also of *Ochsenheimer*.

- Papilio* (Nymph. Gemm.) *Arge* (occitanica), *Esper*, *Schmett.* i. pt. 2. tab. 96. cont. 51. figs. 3, 4 (1777).
Papilio Psyche, *Hübner*, *Eur. Schmett.* i. pl. 44. figs. 198, 199; var. pl. 133. figs. 676, 677, pl. 137. figs. 696, 697 (1805).
Satyrus Psyche, *Godart*, *Enc. Méth.* ix. p. 508. n. 95 (1819); *Lép. de France*, ii. p. 138. n. 52, pl. 19. figs. 3, 4 (1822).
Arge Psyche, *Westwood* in *Gen. Diurn. Lepid.* p. 384. n. 8 (1851).
 South of France, Germany, ?? Port Natal. B.M.

Var. IXORA.

- Arge Ixora*, *Boisduval*, *Icon. Hist. Lép.* p. 140. n. 7, pl. 27. figs. 3, 4 (1832).
Var. ? Papilio Psyche, *Hübner*, *Eur. Schmett.* i. pl. 137. figs. 694, 695 (1805).
 South of France.

9. *Arge Amphitrite.*

- ♂. *Papilio Amphitrite*, *Hübner*, *Eur. Schmett.* i. pl. 43. figs. 196, 197 (1805).
Satyrus Amphitrite, *Godart*, *Enc. Méth.* ix. p. 508. n. 94 (1819).
Arge Amphitrite, *Hübner*, *Verz. bek. Schmett.* p. 60. n. 578 (1816); also of *Boisduval* (1832).
Papilio Arge, *Fuessly*, *Alt. Mag.* i. pt. 2. p. 202 (1775); also of *Bergsträsser*, *Schneider*, *Petagna*, and *Herbst*.
Papilio (Nymph. Gemm.) *Arge*, *Sulzer*, *Gesch. Insekt. Linn. Syst.* p. 144, pl. 16. figs. 8, 9 (1776); also of *Esper*.
Satyrus Arge, *Duponchel*, *Lép. de France*, i. p. 164. n. 55, pl. 24. figs. 5, 6 (1832).
Papilio Arge Sicala, *Borkhausen*, *Naturgesch. Schmett.* i. p. 107. n. 50 (1788).
 Naples. B.M.

As the name *Arge* has been adopted for the genus, it has been necessary to retain the synonymic name *Amphitrite*.

Var. PHERUSA.

- Papilio Pherusa*, *Dahl* (1823?).
Arge Pherusa, *Boisduval*, *Icon. Hist. Lép.* p. 141. n. 8, pl. 26. figs. 4-6 (1832).
Arge Pherusa, var. *Plesasure*, *Bellier de la Chavignerie*, *Ann. Ent. Soc. France*, p. 678 (1860).
 Sicily (Bellier), South of France ?

This insect must be considered a local form of *Amphitrite*; it is intermediate between it and the preceding species.

Genus 52. RHAPHICERA, *Butler* (1867).Typical Species *Rhaphicera Satricus*.1. *Rhaphicera Satricus*.♀. *Lasiommata Satricus*, *Hewitson & Westwood*, in *Gen. Diurn. Lepid.* p. 387. n. 14, pl. 64. fig. 4 (1851).*Rhaphicera Satricus*, *Butler*, *Ann. & Mag. Nat. Hist.* 3rd ser. xix. p. 164. n. 1, pl. 4. fig. 3 (1867).

Eastern Himalayas.

♀, B.M.

2. *Rhaphicera Moorei*.♀. *Rhaphicera Moorei*, *Butler*, *Ann. & Mag. Nat. Hist.* 3rd ser. xix. p. 164. n. 2, pl. 4. figs. 4, 4a (1867).

North-western Himalayas.

♀, Coll. Moore.

Genus 53. RAGADIA, *Westwood* (1851).Typical Species, *Ragadia Crisia*.1. *Ragadia Crisia*.*Euptychia Crisia*, *Hübner*, *Zuträge*, figs. 675, 676 (1806).*Neonympha* (*Ragadia*) *Crisia*, *Westwood* in *Gen. Diurn. Lepid.* p. 376. n. 26 (1851).*Hipparchia Makuta*, *Horsfield*, *Cat. Lep. Mus. East-Ind. Comp.* tab. 5. figs. 9, 9a (1829).*Ragadia Makuta*, *F. Moore*, *Cat. Lep. Mus. East-Ind. Comp.* i. p. 226. n. 473 (1857).

Java (Moore), Borneo, East Indies.

B.M.

2. *Ragadia Luzonia*.*Ragadia Luzonia*, *Felder*, *Wien. ent. Monats.* v. p. 304 (1861).

Luzon (Coll. Semper and Felder), Philippines.

B.M.

3. *Ragadia Melindena*.♀. *Ragadia Melindena*, *Felder*, *Lep. Nov. Ins. Philipp. in Lep. Diagn.* p. 45. n. 99.

Mindanao.

According to Felder, intermediate between (*Makuta*) *Crisia* and *Luzonia*; it can scarcely be more than a variety of the latter.4. *Ragadia Crisilda*.*Ragadia Crisilda*, *Hewitson*, *Exot. Butterf.* iii. p. 98, pl. 49. figs. 5, 6 (1862).

Sylhet.

Coll. Hewitson.

Genus 54. ACROPHTALMIA (*sic*), *Felder* (1861).Typical Species *Acrophtalmia Artemis*.1. *Acrophtalmia Artemis*.*Acrophtalmia Artemis*, *Felder*, *Wien. ent. Monatschr.* v. p. 305 (1861).

Luzon (Coll. Semper and Felder), Philippines.

B.M.

2. *Acrophtalmia Chione*.*Acrophtalmia Chione*, *Felder*, *Nov. Voy.* iii. tab. 68. figs. 12, 13 (published Sept. 1867).

Ternate.

B.M.

3. *Acrophtalmia Leuce*.*Acrophtalmia Leuce*, *Felder*, *Nov. Voy.* iii. p. 486. n. 828 (1867).Celebes (*Felder*).Genus 55. ARGYROPHENGA, *Doubleday* (1845).Typical Species *Argyrophenga Antipodum*.1. *Argyrophenga Antipodum*.*Argyrophenga Antipodum*, *E. Doubleday*, *Ann. & Mag. Nat. Hist.* xvi. p. 307 (1845); also of *Westwood*.

New Zealand.

B.M.

Genus 56. ARGYROPHORUS, *Blanchard* (1852).Typical Species *Argyrophorus argenteus*.1. *Argyrophorus argenteus*.*Argyrophorus argenteus*, *Blanchard* in *Gay's Fauna Chilena*, vii. p. 30, lam. 2. figs. 9-11 (1852) (*Chionobas* Arg. on plate).

Chili.

2. *Argyrophorus Williamsianus*, *White*, MS. Pl. IV. fig. 1.*Alæ supra fusæ, puncto subapicali indistincto nigro, ferrugineo cincto, punctoque ferrugineo subanali; ciliis albo variis.**Alæ anticæ subtus ochreo-ferrugineæ, margine externo fuscescente; macula elongata subapicali nigra; ciliis albis fusco variis: posticæ fusæ, venis apud basim albis, fascia media valde irregulari nigro-fusca a venis interrupta; maculis quinque discalibus nigris, secunda majore distincta, aliis indistinctis; ciliis albis.*Exp. alar. unc. $1\frac{1}{8}$.

Port Famine.

B.M.

Genus 57. *ÆNEIS*, *Hübner* (1816).Typical Species, *Æneis Norna* of *Esper*.1. *Æneis Jutta*.♀. *Papilio Jutta*, *Hübner*, *Eur. Schmett.* i. pl. 120. figs. 614, 615 (1805).*Æneis Jutta*, *Hübner*, *Verz. bek. Schmett.* p. 58. n. 560 (1816).*Chionobas Jutta*, *Boisduval*, *Icon. Hist. Lép.* p. 187. n. 3, pl. 38. figs. 1-4 (1832); also of *Herrich-Schäffer*.*Satyrus Jutta*, *Duponchel*, *Lép. de France*, i. p. 250. n. 91, pl. 40. figs. 3-5 (1832); also of *Zetterstedt* (1840).♀ *Var. Chionobas* *Balder*, *Herrich-Schäffer*, *Eur. Schmett.* i. tab. 79. figs. 385, 386 (1845).

Arctic America.

B.M.

Var. BALDERI.*Eumenis Balderi*, *Hübner*, *Zuträge Exot. Schmett.* figs. 981, 982 (1806).*Satyrus Balder*, *Boisduval* in *Guérin's Règne Animal*, p. 487, pl. 80. fig. 1 (1829-44); also of *Le Conte* (1833).*Chionobas Balder*, *Boisduval*, *Icon. Hist. Lép.* p. 189. n. 4, pl. 39. figs. 1-3 (1832); also (part.) *Herrich-Schäffer*.

Polar Regions, Hudson's Bay, Lapland (Westw.).

This form is not in the British-Museum Collection. The species included by Mr. Doubleday in his catalogue is the true *Jutta*. *Balderi* is allowed by Mr. Scudder and others to be nothing more than a variation from the true type of *Jutta*. It chiefly differs in being smaller, with more distinct fulvous bands and fewer ocelli. It may be a race, but can scarcely be separated as a species.

2. *Æneis Aello*.*Papilio* (Nymph. Phal.) *Aello*, *Esper*, *Schmett.* i. pt. 2. tab. 115. cont. 70. fig. 1 (1777).*Papilio Aello*, *Hübner*, *Eur. Schmett.* i. pl. 102. figs. 519-521 (1805); also of *Ochsenheimer* (1807).*Eumenis Aello*, *Hübner*, *Verz. bek. Schmett.* p. 58. sp. 554 (1816).*Satyrus Aello*, *Godart*, *Enc. Méth.* ix. p. 518. n. 15 (1819); also of *Zetterstedt* (1840), but not of *Herrich-Schäffer*.*Var. Chionobas Aello*, *Boisduval*, *Icon. Hist. Lép.* i. p. 183, pl. 36. figs. 1-3 (1832); also of *Duponchel* (*Satyrus Aello*).*Var. ? Papilio Norna*, *Hübner*, *Eur. Schmett.* i. pl. 31. figs. 141, 142, & 142* (1805).

Alps.

B.M

3. *Œneis Tarpeia*.

Papilio (Nymph. Gemm.) *Tarpeia*, *Esper*, *Schmett.* i. pt. 2. tab. 83. cont. 23. figs. 1, 2 (1777).

Papilio Tarpeia, *Ochsenheimer*, *Schmett. v. Eur.* i. p. 203. n. 18 (1807).

Satyrus Tarpeius, *Duponchel*, *Lép. de France*, i. p. 207. n. 71, pl. 31. figs. 6, 7 (1832); also of *Zetterstedt* (1840).

Papilio Tarpeja, *Borkhausen*, *Naturgesch. eur. Schmett.* i. p. 101. n. 41 (1788), ii. p. 209. n. 41 (1789); *Hübner*, *Eur. Schmett.* i. pl. 156. figs. 779-782 (1805).

Eumenis Tarpeja, *Hübner*, *Verz. bek. Schmett.* p. 58. n. 556 (1816).

Papilio (*Hipparchia*) *Tarpeja*, *Freyer*, *Neuere Beiträge*, v. p. 75. n. 753, tab. 427. figs. 3, 4 (1845).

Chionobas Tarpeja, *Herrich-Schäffer*, *Eur. Schmett.* i. tab. 13. figs. 61-64.

Papilio Celimene, *Cramer*, *Pap. Exot.* iv. p. 169, pl. 376. figs. E, F (1782).

Arctic America.

B.M.

Var. *NANNA*.

Chionobas Nanna, *Ménétriés*, *Bull. Acad. St. Pétersb.* tom. xvii. p. 216. n. 15 (1844); *Schrenck*, *Fauna Lepid. Sibér. orient. et l'Amour*, p. 38. n. 72, tab. 3. fig. 5 (1859).

River Amoor (*Schrenck*).

4. *Œneis Chryxus*.

Chionobas Chryxus, *Hewitson* in *Gen. Diurn. Lepid.* p. 383. n. 13, pl. 64. fig. 1 (1851).

Rocky Mountains.

B.M.

5. *Œneis Nevadensis*.

Chionobas Nevadensis, *Felder*, *Reise der Fregatte Novara*, iii. tab. 62. figs. 4, 5 (1867).

California, Sierra Nevada (*Felder*).

Closely allied to a new species from Vancouver's Island in the National Collection. It must come next to *Chryxus*.

6. *Œneis gigas*, sp. nov. Pl. II. fig. 2.

♂. Alæ supra fulvæ: anticæ elongatæ, marginibus costali et externo fuscis; fascia lata basali obliqua nigro-fusca; area basali rufo-fuscescente; ocellis duobus distinctis nigris albo pupillatis, superiore paulum majore: posticæ margine externo fusco undulato; ocello subanali nigro, albo pupillato; ciliis omnibus fuscis albo variis: corpus nigrum; antennis subferrugineis.

Anticæ subtus pallidiores, area basali minime fuscescente; costa et apice albicantibus et nigro lituratis; ocellis velut supra; margine externo fuscescente; ciliis albo variis: posticæ nivæ fusco lituratæ; fascia media velut in *Celæno* fere, irregulari, fuscescente, nigro-fusco limitata; nebula fusca basali, et altera marginali lata, punctos tres albos includente; ocello velut supra, sed minore; maculis ad venarum fines nigris; ciliis albo variis.

Exp. alar. unc. 2½.

Vancouver's Island.

B.M.

7. *Oeneis Urda*.

Hipparchia Urda, *Eversmann in Bull. Soc. Imp. Nat. Mosc.* xx. pt. 3. p. 69. n. 4, tab. 2. figs. 1-4 (1847).

Chionobas Urda, *Herrich-Schäffer, Eur. Schmett.* i. tab. 97. figs. 461-463.

Dauria.

8. *Oeneis Norna*.

♀. *Papilio* (Nymph. Gemm.) *Norna*, *Esper, Schmett.* i. pt. 2. p. 82, tab. 108. cont. 63. fig. 4 (1777).

Papilio Norna, *Thunberg, Diss. Acad. Ins. Su.* tab. 5. fig. 11 (1801); also of *Hübner* and *Ochsenheimer*.

Oeneis Norna, *Hübner, Verz. bek. Schmett.* p. 58. sp. 557 (1816).

Satyrus Norna, *Godart, Enc. Méth.* ix. p. 518. n. 116 (1819).

Var. *Papilio Hilda*, *Schneider, Magaz.* iv. p. 414. n. 3, v. p. 585 (1787-8).

Var. *Chionobas Norna*, *Boisduval, Icon. Hist. Léop.* p. 185. n. 2, pl. 36. figs 4-6 (1832).

Satyrus Norna, *Duponchel, Léop. de France*, i. p. 205. n. 70, pl. 31. figs. 4, 5 (1832).

Lapland.

♂ ♀, B.M.

Var. *CELÆNO*.

♂. *Papilio Celæno*, *Hübner, Eur. Schmett.* i. pl. 34. figs. 152, 153 (1805).

Lapland.

♂ ♀, B.M.

Var. *FULLA*.

Hipparchia (*Chionobas*) *Fulla*, *Eversmann in Bull. Soc. Imp. Nat. Mosc.* p. 614 (1851).

Chionobas Fulla, *Herrich-Schäffer, Eur. Schmett.* i. tab. 127. figs. 615, 616; also of *Westwood* (*Bore*, var. ?) (1852).

Eastern Siberia.

This form differs very little from *Celæno*.

9. *Œneis Uhleri*.

Chionobas Uhleri, *Reakirt, Proc. Ent. Soc. Philad.* vi. p. 143 (1866).

Rocky Mountains, Colorado Territory.

10. *Œneis Sculda*.

Hipparchia (*Chionobas*) *Sculda*, *Eversmann, Bull. Soc. Imp. Nat. Mosc.* p. 612 (1851).

Chionobas Sculda, *Herrich-Schäffer, Eur. Schmett.* i. tab. 127. figs. 613, 614.

Eastern Siberia.

Nearly allied to *Norna*, but with central band resembling that of *Crambis*.

11. *Œneis Crambis*.

Papilio (*Hipparchia*) *Crambis*, *Freyer, Neuere Beiträge*, v. p. 99. n. 783, tab. 440. figs. 3, 4 [not good] (1845).

Chionobas Crambis (*Ch. Bore*, var.), *Westwood in Gen. Diurn. Lepid.* p. 383. n. 11 (1851).

Chionobas Œno (*synonym*), *Scudder, Proc. Ent. Soc. Philad.* v. n. 5 (1865).

Lapland (Freyer), Arctic America.

♂ ♀, B.M.

12. *Œneis Œno*.

Chionobas Œno, *Boisduval, Icon. Hist. Lép.* p. 195. n. 7, pl. 39. figs. 4-6 (1832); also of *Herrich-Schäffer, Eur. Schmett.*

♀ var., *ibid.* i. tab. 27. figs. 123, 124 (1845).

Var. *Chionobas Also*, *Boisduval, Icon. Hist. Lép.* p. 197. n. 8, pl. 40. figs. 1, 2 (1832); also of *Herrich-Schäffer, Westwood, &c.* (B.M.). *Exitiosa*, *Harris* (*Boisd. cit.*).

Arctic America, Repulse Bay.

B.M.

13. *Œneis assimilis*, sp. nov. Pl. II. fig. 10.

Alæ supra forma et colore subhyalinae; *subtus coloribus Œnonis, fascia autem posticarum minus distincta et ad costam latiore, ob quam rem causam, multo magis regulari.*

Exp. alar. unc. 1 $\frac{7}{8}$.

Repulse Bay.

B.M.

This species is closely allied to *Œno*, but with longer wings and differently formed central band.

14. *Œneis semidea*.

Hipparchia semidea, *Say, Amer. Ent.* iii. pl. 50 (1828).

Chionobas semidea, *Edwards in Morr. Syn. Lepid. N. Amer.*
p. 351 (1860); also of *Scudder (Bost. Journ. vii.)*.

Cænonympha semidea, *Morris, Syn. Lep.* p. 80 (1860).

White Mountains.

B.M.

15. *Œneis subhyalina*.

Hipparchia subhyalina, *Curtis in Append. Ross's Narrative*
N. W. Passage, p. 68. n. 13 (1835).

Erebia subhyalina, *Westwood in Gen. Diurn. Lepid.* p. 380.
n. 56 (1851).

Arctic America.

B.M.

We do not possess the type of this species; but our specimens agree exactly with Mr. Curtis's description. They come near *Taygete*, but are of a more smoky colour.

16. *Œneis Taygete*.

Œneis Taygete, *Hübner, Samml. exot. Schmett.* iii. (1806–24).
Chionobas Taygete (part.), *Herrich-Schäffer, Eur. Schmett.* i.
tab. 24. figs. 114, 115 (1845); also of *Westwood and Ed-*
wards, ♀ (1862).

Chionobas Calais, *Scudder, Proc. Ent. Soc. Philad.* v. n. 3
(1865).

Labrador.

B.M.

17. *Œneis Bore*.

Papilio (Dan. Cand.) *Bore*, *Esper, Schmett.* i. pt. 2. tab. 100. cont.
55. fig. 1 (1777).

Œneis Bore, *Hübner, Verz. bek. Schmett.* p. 58. sp. 558
(1816).

Erebia Bore, *Dalman, Pap. Su.* p. 80. n. 7 (1824).

Chionobas Bore, *Boisduval, Icon. Hist. Lép.* p. 193. n. 6, pl. 37.
figs. 1–3 (1832).

Papilio Polixenes, *Fabricius, Ent. Syst.* iii. pt. 1. p. 152. n. 466
(1793).

Neonympha ? *Polixenes*, *Westwood in Gen. Diurn. Lepid.* p. 503.
n. 131 (1851).

Var. Papilio (Nymph. Gemm.) *Norna*, *Schneider, Mag.* iv.
p. 415. n. 4 (1787).

Papilio Norna, var. γ , *Thunberg, Diss. Acad.* p. 50 (1801).

Var. Satyrus Bore, *Duponchel, Lép. de France*, i. p. 209. n. 72,
pl. 32. figs. 1, 2 (1832).

Var. ? ♂. Papilio Fortunatus, *Fabricius, Ent. Syst.* iii. pt. 1.
p. 214. n. 670 (1793).

Satyrus Fortunatus, *Godart, Enc. Méth.* ix. p. 519. n. 117
(1819).

Lapland.

♂ ♀, B.M.

Var. Bootes.

Eumenis ? *Bootes*, *Hübner*, *Eur. Schmett.* i. figs. 1025–1028 (1805).

Chionobas Bootes, *Boisduval*, *Icon. Hist. Léop.* p. 191. n. 5, pl. 37. figs. 4–6 (1832); var. ♀, *Herrich-Schäffer*, tab. 80 (1845).

Satyrus Bootes, *Duponchel*, *Léop. de France*, i. p. 211. n. 73, pl. 32. figs. 3–5 (1832).

Papilio Bore, *Hübner*, *Eur. Schmett.* i. pl. 29. figs. 134–136 (1805); also of *Ochsenheimer*.

Chionobas Taygete (part.), *Herrich-Schäffer*, *Eur. Schmett.* i. tab. 24. figs. 112, 113 (1845).

Lapland, Labrador.

B.M.

This form is now generally allowed to be no more than a variety of *Bore*. It may be distinct; for the central band is not identical with that of *Bore*, being much less irregular and better defined. The upperside of the wings is of a brighter ochre-colour with darker lines, and the underside less streaked with minute stræ. However, it may be as well to consider it nothing more than a race.

18. *Ceneis pumilus*.

Chionobas pumilus, *Felder*, *Reise der Fregatte Novara*, iii. tab. 69. figs. 6, 7 (1867).

Eastern Himalayas.

Seems closely allied to *Bore* of Esper; but according to Felder it is distinct from all the known species.

Genus 58. ARGYNNINA, *Butler* (1867).**1. *Argynnina Hobartia*.**

Lasiommata Hobartia, *Westwood in Gen. Diurn. Lepid.* p. 387. n. 21 (1851).

Argynnina Hobartia, *Butler*, *Ann. & Mag. Nat. Hist.* 3rd ser. xix. p. 165. n. 1 (1867).

Lasiommata Lasus, *Hewitson*, *Trans. Ent. Soc. London*, 3rd ser. ii. p. 248. n. 9, pl. 16. fig. 5 (1865).

Van Diemen's Land.

B.M.

2. *Argynnina Lathoniella*.

Lasiommata ? (*Xenica*) *Lathoniella*, *Westwood in Gen. Diurn. Lepid.* p. 387. n. 25 (1851).

Argynnina Lathoniella, *Butler*, *Ann. & Mag. Nat. Hist.* 3rd ser. xix. p. 165. n. 2, see pl. 4. fig. 6 (1867).

Van Diemen's Land.

B.M.

3. *Argynnina Leprea*.

Lasiommata Leprea, *Hewitson in Trans. Ent. Soc. London*,
3rd ser. ii. p. 249. n. 10, pl. 16. figs. 6, 7 (1865).

Australia (Hewitson).

Genus 59. *GEITONEURA*, *Butler* (1867).

Typical Species *Geitoneura Klugii*.

1. *Geitoneura Klugii*.

♀. *Satyrus Klugii*, *Guérin, Voy. Coquille, Atlas, Ins.* pl. 17.
fig. 2 (1826).

♂ ♀. *Geitoneura Klugii*, *Butler, Ann. & Mag. Nat. Hist.* 3rd ser.
xix. p. 165. n. 1 (1867).

♂. *Satyrus Singa*, *Boisduval, Voy. de l'Astrolabe Entom.* pt. 1.
p. 144 (1832-33).

Lasiommata? (*Xenica*) *Singa*, *Westwood in Gen. Diurn. Lepid.*
p. 387. n. 23 (1851).

Lasiommata Philerope (part.), *Westwood, ibid.* n. 19 (1851).

King George's Sound, New Holland.

B.M.

2. *Geitoneura Achanta*.

Papilio Achanta, *Donovan, Ins. New Holland*, pl. 22. fig. 2
(1805).

Satyrus Achanta, *Godart, Enc. Méth.* ix. p. 500 (1819); *Bois-*
duval, Voy. de l'Astrolabe, Entom. pt. 1. p. 147 (1832-33).

Lasiommata? (*Xenica*) *Achanta*, *Westwood in Gen. Diurn.*
Lepid. p. 387. n. 22 (1851).

Xenica Achanta, *F. Moore, Cat. Lep. Mus. East-Ind. Comp.* i.
p. 228. n. 479 (1857).

Geitoneura Achanta, *Butler, Ann. & Mag. Nat. Hist.* 3rd ser.
xix. p. 165. n. 2 (1867).

Tisiphone Acanthe, *Hübner, Zuträge*, figs. 267, 268 (1806).

New Holland.

B.M.

3. *Geitoneura*? *Cordace*.

Tisiphone Cordace, *Hübner, Zuträge Exot. Schmett.* figs. 797,
798 (1806).

Lasiommata Cordace, *Westwood in Gen. Diurn. Lepid.* p. 387.
n. 13 (1851).

East India.

Genus 60. HYPOCYSTA, *Westwood* (1851)*.Typical Species *Hypocysta Irius*.1. *Hypocysta Hygea*.

Hypocista Hygea, *Hewitson, Exot. Butterf.* iii. p. 99, pl. 50.
figs. 2, 3 (1863).

Aru Islands.

B.M.

2. *Hypocysta Hecaerge*.

Hypocista Hecaerge, *Hewitson, Exot. Butterf.* iii. p. 99, pl. 50.
figs. 4, 5 (1863).

New Guinea.

B.M.

3. *Hypocysta Hæmonia*.

Hypocista Hæmonia, *Hewitson, Exot. Butterf.* iii. p. 99, pl. 50.
fig. 1 (1863).

Aru Islands.

B.M.

4. *Hypocysta Osyris*.

Satyrus Osyris, *Boisduval, Voy. de l'Astrolabe, Entom.* pt. 1.
p. 154 (1832-33).

Satyrus? *Osyris*, *Westwood in Gen. Diurn. Lepid.* p. 392. n. 48
(1851).

Hypocysta Osyris, *Felder in Wien. ent. Monatschr.* iv. n. 88,
Taf. 10. fig. 5 (1860).

Aru Islands.

B.M.

Var. Alis anticis magis fuscescentibus, posticis magis albican-
tibus.

Dorey (New Guinea).

B.M.

5. *Hypocysta Irius*.

Papilio Irius, *Fabricius, Syst. Ent.* p. 487 (1775-78); *Donovan,*
Ins. New Holland, pl. 28. fig. 1 (1805).

Satyrus Irius, *Godart, Enc. Méth.* ix. p. 548. n. 177 (1819);
also of *Boisduval* (1832-33).

Cænonympha Irius, *Westwood in Gen. Diurn. Lepid.* p. 398.
n. 27 (1851).

Neonympha Adiante, *Hübner, Zuträge Exot. Schmett.* figs. 545,
546 (1806).

New Holland.

B.M.

* This genus is only briefly characterized by Westwood in a remark on the variation of the genus *Cænonympha*, *Gen. Diurn. Lepid.* p. 397, lines 7-10 (1851).

6. *Hypocysta Antirius*. Pl. IV. fig. 9.

Alæ fulvæ, apice, margine externo et sæpe costa nigris: posticæ oculo minuto (rarius indistincto) nigro, argenteo pupillato; margine externo fuscescente.

Alæ subtus pallidiores, marginibus plerumque fusciscentibus: posticæ area basali angulariter obscuriore; ocellis duobus distinctis, uno apicali, altero anali, æqualibus, nigris, iride fulva, margine fusco, cinctura pupillaque argenteis; puncto interjecto et linea marginali argenteis.

Exp. alar. unc. $1\frac{1}{3}$ — $1\frac{1}{2}$.

Moreton Bay, Australia.

B.M.

Var. Linea media subtus dentata; marginibus supra nigerrimis.

North-west Australia.

B.M.

7. *Hypocysta Euphemia*.

Cænonympha (*Hypocysta*) *Euphemia*, Hewitson & Westwood in *Gen. Diurn. Lepid.* p. 398. n. 28, pl. 67. fig. 3 (1851).

New Holland.

B.M.

Genus 61. *ETEONA*, Westwood (1851).

Typical Species *Eteona Tisiphone*.

1. *Eteona Tisiphone*.

Panopæa *Tisiphone*, Hübner, *Zuträge Exot. Schmett.* figs. 911, 912 (1806).

Euterpe *Tisiphone*, Boisduval, *Sp. Gén. Lép.* pt. 1. p. 411. n. 8 (1836).

Eteona Tisiphone, E. Doubleday, *List Lep. Brit. Mus. App.* p. 21 (1848); also of Hewitson & Westwood in *Gen. Diurn. Lepid.* pl. 42. fig. 3 (1851).

Chili.

B.M.

2. *Eteona Eupolis*.

Eteona Eupolis, Hewitson in *Trans. Ent. Soc. London*, ser. 3. ii. p. 247. n. 6, pl. 16. fig. 2 (1865).

Rio Janeiro.

Coll. Hewitson and B.M.

Genus 62. *LYMANOPODA*, Westwood (1851).

Typical Species *Lymanopoda Samius*.

1. *Lymanopoda obsoleta*.

Lymanopoda (*Sarromia*) *obsoleta*, Westwood & Hewitson in *Gen. Diurn. Lepid.* p. 402. n. 3, pl. 67. fig. 5 (1851).

Bolivia, Bogota.

♂, B.M.

Var. Minor, ocellis subtus distinctis.

Venezuela.

♂ ♀, B.M.

The female differs a good deal from the male, more especially in the colouring of the underside, the front wings being ferruginous brown; the apex, costa, and a large, somewhat wedge-shaped sub-anal patch ochreous. There are also two circular transparent spots near the apex. The hind wings are pale ochreous brown, with the base and a transverse interno-discal subcentral band and one or two scarcely perceptible submarginal nebulæ dun-coloured.

2. *Lymanopoda lactea*.

Lymanopoda lactea, *Hewitson, Journal of Entomology*, p. 157. sp. 2, pl. 9. figs. 2, 3 (1862).

New Granada (Hewitson).

3. *Lymanopoda Samius*.

♂. *Lymanopoda Samius*, *Westwood & Hewitson Gen. Diurn. Lepid.* p. 402. n. 1, pl. 67. fig. 6 (1851).

Bogota (New Granada).

♂, B.M.

4. *Lymanopoda Ionius*.

♂. *Lymanopoda Ionius*, *Westwood in Gen. Diurn. Lepid.* p. 402. n. 2 (1851).

♂ as ♀. *Lymanopoda Samius*, *Hewitson & Westwood, op. cit.* pl. 67. fig. 7 (1851).

Bogota.

♂, B.M.

The types of *L. Samius* and *Ionius* have been carefully examined, and there seems no doubt that they are both males.

5. *Lymanopoda ferruginosa*, sp. nov. Pl. IV. fig. 3.

Alæ apice anticarum convexo; margine interno brevior; supra punctis duobus albidis, uno subapicali, altero discali: posticæ marginis medio magis productæ; hæ quoque subtus maculis omnibus magnis discalibus aurantiacis nec argenteis; macula autem discoidea fulvo-argentea: aliter velut in *L. Labda* (Hewits.).

Exp. alar. unc. $1\frac{2}{3}$.

Bolivia.

B.M.

This beautiful little species is closely allied to *Labda*, from which it chiefly differs in the rounded apex and short inner margin of the front wings, which show two white points on the upperside, also in all the larger silver spots on the underside of the hind wings being replaced by orange ones, the discoidal streak, however, being of a yellowish silver-colour.

6. *Lymanopoda Lebbæa*.

Lymanopoda Lebbæa, *Felder, Reise der Fregatte Novara*, iii.
p. 473. n. 810 (1867).

Bogota (Felder).

Dr. Felder says of this species that it stands near *L. Labda* of Hewitson, of which he only knows the figure, but that in the form of its wings it approaches *L. Ionius* of Westwood.

7. *Lymanopoda Lanassa*.

Lymanopoda Lanassa, *Felder, Reise der Fregatte Novara*, iii.
p. 474. n. 811 (1867).

Bogota (Felder).

Differs from *Leæna*, Hewits., in having longer wings and a narrower ochraceous band below.

8. *Lymanopoda apiculata*.

Lymanopoda apiculata, *Felder, Reise der Fregatte Novara*, iii.
p. 474. n. 812 (1867).

Bogota (Felder).

"Larger than *Leæna*, the wings longer within, the outer margin gently undulated below the apex."

9. *Lymanopoda perita*.

Pronophila perita, *Hewitson, Exot. Butterf.* iv. pl. incog. fig. 25
(1868).

Ecuador.

Coll. Hewitson.

It is difficult to say whether this is a *Lymanopoda*; but it certainly looks like one; an examination of the cell of the front wings would at once settle the question.

10. *Lymanopoda Labda*.

Lymanopoda Labda, *Hewitson, Journal of Entomology*, p. 157.
sp. 3, pl. 9. fig. 4 (1862).

New Granada.

B.M.

11. *Lymanopoda Leæna*.

Lymanopoda Leæna, *Hewitson, Journal of Entomology*, p. 156.
sp. 1, pl. 9. fig. 1 (1862).

New Granada (Quito), Amazon (Nauta).

B.M.

12. *Lymanopoda albocincta*.

Lymanopoda albocincta, Hewitson, *Journal of Entomology*,
p. 157. sp. 4, pl. 9. fig. 5 (1862).

Bogota.

B.M.

13. *Lymanopoda albomaculata*.

Lymanopoda albomaculata, Hewitson, *Journal of Entomology*,
p. 158. sp. 5, pl. 9. fig. 6 (1862).

Bolivia.

B.M.

Allied to *albocincta*, but differing considerably in the band of the underside.

14. *Lymanopoda venosa*, sp. nov. Pl. IV. fig. 5.

Alæ supra forma *albomaculata*, fuscae, margine apicali minime dilutiores.

Alæ subtus pallidiores, venis, et furca posticarum discoidea, nigris: anticæ maculis tribus decrescentibus subhastatis marginalibus mediis ferrugineis.

Exp. alar. unc. 2.

Bolivia.

B.M.

The form and size of *albomaculata*; coloured below something like an *Acraea*. The entire upperside and the underside of the front wings in our specimen, when magnified, are seen to be sprinkled over with gold scales.

15. *Lymanopoda Acraeida*, sp. nov. Pl. IV. fig. 6.

Alæ elongatæ, supra rufo-fuscae, fascia anticis submarginali rufo-ferruginea subarcuata a venis nigris interrupta, maculas sex sic formante; puncto in serie eadem costali albido: posticæ inter venas squamis nonnullis rufis indistinctissime sparsæ.

Alæ subtus pallidiores: anticæ cinerascens, costa discoidea ferruginea; fascia marginali intus bene limitata, extus fulvo diffusa, margine externo et apice olivaceo-fuscescentibus, a venis plicisque nigris interrupta: posticæ olivaceo-fuscae, basi diffuse rufescente, macula parva sub cellæ extimum et margine externo paulum rufescentibus, venis plicisque nigerrimis.

Exp. alar. unc. $2\frac{2}{3}$.

Bolivia.

B.M.

There can be little doubt that this species mimics some form of *Acraea*. It is longer in the wing than any of the known species of *Lymanopoda*.

Genus 63. *STEROMA*, Westwood (1851).Typical Species *Steroma Bega*.1. *Steroma Bega*.*Steroma Bega*, Hewitson & Westwood, *Gen. Diurn. Lepid.* p. 400.
n. 1, pl. 67. fig. 6 (1851).

Venezuela.

B.M.

Local form. ANDENSIS.*Steroma Andensis*, Felder, *Reise der Fregatte Novara*, iii. p. 475.
n. 814 (1867).

Bogota (Felder), Bolivia.

B.M.

2. *Steroma pronophila*.*Steroma pronophila*, Felder, *Reise der Fregatte Novara*, iii.
p. 475. n. 815 (1867).

Bogota.

B.M.

3. *Steroma superba*, sp. nov. Pl. V. fig. 6.

♂. Alæ supra piceo-nigræ, marginibus rufescentibus: anticæ macula permagna discali ovali sericea; areola basali alarum omnium brunneo rorata.

Alæ subtus pallidiores: anticæ cupreo-fuscæ, nitentes; area basali fuscescente, margine basali nigro variegato; apice ochraceo punctato: posticæ fuscæ, nigro irregulariter passim variegatæ; macula parva cellulari et altera costali ochreo-argenteis; punctis septem ochraceis quasi ocellis submarginalibus; ocello minutissimo anali, nigro, bipupillato; puncto longitudinali infraanali, squamisque nonnullis marginalibus sparsis violaceis.

Exp. alar. unc. 2½.

Bolivia.

B.M.

This is the largest species of *Steroma* yet described; it is of nearly the same shape and size as *Callerebia scanda*, Kollar; the wings are strongly produced, and the hind wings are waved, as in *Steroma pronophila*, Felder.Genus 64. *PEDALIODES*, Butler (1867)*.Typical Species *Pedaliodes Poësia*.1. *Pedaliodes Poësia*.*Pronophila Poësia*, Hewitson, *Trans. Ent. Soc. London*, 3rd ser. i.
p. 6. n. 10, pl. 3. fig. 19 (1861).

Bogota.

B.M.

* Ann. & Mag. Nat. Hist. xx. p. 267, fig. 1.

2. Pedaliodes Proerna.

Pronophila Proerna, Hewitson, *Trans. Ent. Soc. London*, 3rd ser. i. p. 9. n. 19, pl. 4. fig. 29 (1861).

Bogota.

B.M.

Var. Macula subanali posticarum subtus majore.

Bolivia.

B.M.

This chiefly differs from the preceding species in wanting the costal white spot on the underside of the hind wings. It is very doubtful if it is more than a variety of *Poesia*.

3. Pedaliodes granulata, sp. nov. Pl. IV. fig. 8.

? *Pronophila Perperna*, Felder, *Reise der Fregatte Novara*, iii. p. 468. n. 797 (1867).

Bogota (Felder).

? *id.* B.M.

"Larger than *P. Proerna*, Hewits., the margins of the wings more deeply sinuated." Dr. Felder seems entirely to have overlooked Mr. Hewitson's description of a species under the name of *Perperna*; indeed he has redescribed the latter under the name of *satyroides*.

4. Pedaliodes empusa.

Pronophila empusa, Felder, *Reise der Fregatte Novara*, iii. p. 468. n. 798 (1867).

Bogota (Felder).

Much smaller than *Proerna*, the underside of the hind wings resembling *Poesia*.

5. Pedaliodes Pisonia.

♀. *Pronophila Pisonia*, Hewitson, *Trans. Ent. Soc. London*, 3rd ser. i. p. 6, pl. 3. fig. 21, not 20 (1861).

Pronophila dejecta, H. W. Bates, *Ent. Month. Mag.* i. p. 179. n. 71 (1865).

Venezuela, (var.) Guatemala (Bates).

♂ ♀, B.M.

The insect described, and the variety of the same figured, as the male of this species seems to represent another form, as we have both sexes of *Pisonia* of equal size and perfectly similar in colour and pattern.

6. Pedaliodes Perperna.

♂ ♀. *Pronophila Perperna*, Hewitson, *Trans. Ent. Soc. London*, 3rd ser. i. p. 16 (1861).

Pronophila satyroides, Felder, *Reise der Fregatte Novara*, iii. p. 469. n. 801 (1867).

Venezuela.

♂, B.M.

We do not possess the female of this insect; the male differs from *Pisonia* in having an ocellus and a white spot on the underside of the front wings.

7. *Pedaliodes Phantias*.

Pronophila Phantias, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 6. n. 10, pl. 3. fig. 18 (1861).

Minas Geraes.

Coll. Hewitson.

8. *Pedaliodes Manis*.

♂ ♀. *Pronophila Manis*, *Felder, Reise der Fregatte Novara*, iii. p. 469. n. 799 (1867).

♂. *Pronophila Pisonia*, var., *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 6, pl. 3. fig. 20 (1861).

Bogota (Felder), Venezuela and (var.) Bogota.

B.M.

Our Bogota specimen differs from the Venezuela examples as follows:—

♂. Alæ margine externo minus undulato; alæ posticæ subtus minus rufescentes; alæ anticæ magis productæ.

Var. EREIBA.

♂. *Pronophila Ereiba*, *Felder, Reise der Fregatte Novara*, iii. p. 469. n. 800 (1867).

Bogota (Felder), Venezuela.

♂ ♀, B.M.

This differs very little from the preceding species, of which it is most likely only a variety.

Var. b. Alæ margine subintegro rotundato; fasciis subtus minus dentatis et confusis.

Bolivia.

♂, B.M.

Var. c. Alæ subtus fasciis obsoletis, margine subanali interno quadrate fulvescente.

Bolivia.

♂?, B.M.

Var. d. Macula subanali subtus distinctiore et lutescente, velut in *Phoenissa* formata.

Quito.

♀, B.M.

This last variety is much like the Upper-Amazon insect figured by Mr. Hewitson as a variety of *Paneis*.

9. *Pedaliodes Paneis*.

♂? *Pronophila Paneis*, *Hewitson, Trans. Ent. Soc. London*,

ser. 3. i. p. 8. n. 17, pl. 4. fig. 27 (1861), ♂ var. pl. 4. fig. 26;
var. ? *Exot. Butterf.* 4. pl. incog. fig. 27 (1868).

Nauta (Upper Amazon).

♂, B.M.

10. *Pedaliodes decorata*.

Pronophila decorata, *Felder, Reise der Fregatte Novara*, iii.
p. 470. n. 802, tab. 67. fig. 11 (1867).

Bogota.

B.M.

This and the following species differ slightly from the remaining species of *Pedaliodes*, but not sufficiently to warrant their separation as a genus. The colouring of the hind wings on the underside seems to indicate an affinity to the *Manis* group; but the plumbeous bands give them a very distinct appearance.

11. *Pedaliodes Porphyria*.

Pronophila Porphyria, *Felder, Reise der Fregatte Novara*, iii.
p. 470. n. 803 (1867).

Venezuela.

B.M.

The front wings in this species have a broader and more clumsy appearance than those of *decorata*.

12. *Pedaliodes Apuleja*.

Pronophila Apuleja, *Felder, Reise der Fregatte Novara*, iii. p. 471.
n. 804 (1867).

Venezuela.

id.? B.M.

The wings are shorter than in the two preceding species.

13. *Pedaliodes ocellifera*.

Pronophila ocellifera, *Felder, Reise der Fregatte Novara*, iii.
p. 471. n. 805 (1867).

Bogota (Felder).

Coll. Salvin.

14. *Pedaliodes encycla*.

Pronophila encycla, *Felder, Reise der Fregatte Novara*, iii.
p. 472. n. 806 (1867).

Venezuela.

B.M.

15. *Pedaliodes Phyllalia*.

Pronophila Phyllalia, *Hewitson, Exot. Butterf.* iv. pl. incog.
figs. 21, 22 (Jan. 1868).

Ecuador.

Coll. Hewitson.

16. **Pedaliodes Calisto.**

Pronophila Calisto, *Felder, Reise der Fregatte Novara*, iii. p. 472.
n. 807 (1867).

Bogota (Felder).

17. **Pedaliodes oculata.**

Pronophila oculata, *Felder, Reise der Fregatte Novara*, iii.
p. 472. n. 808 (1867).

Bogota.

B.M.

18. **Pedaliodes mycalesoides.**

Pronophila mycalesoides, *Felder, Reise der Fregatte Novara*, iii.
p. 473. n. 809 (1867).

Bogota (Felder).

19. **Pedaliodes Polusca.**

Pronophila Polusca, *Hewitson, Trans. Ent. Soc. London*, 3rd ser.
i. p. 10. n. 21, pl. 4. figs. 31, 32 (1861).

Bogota.

♂, B.M.

20. **Pedaliodes Pausia.**

Pronophila Pausia, *Hewitson, Trans. Ent. Soc. London*, 3rd ser.
i. p. 8. n. 16, pl. 4. fig. 25 (1861).

Bolivia.

B.M.

21. **Pedaliodes angularis**, sp. nov. Pl. IV. fig. 7.

Alæ supra fuscae, area basali anticarum obscuriore.

Alæ subtus coloribus partim Plotinæ, stria ochracea anticarum magis ad marginem approximante: costa posticarum rectissima, fascia media tenuiore magis angulata; plaga fulvo-ferruginea anali obsoleta; apice paulum albicante.

Exp. alar. unc. 1 $\frac{7}{8}$.

Peru.

B.M.

22. **Pedaliodes Piletha.**

Pronophila Piletha, *Hewitson, Trans. Ent. Soc. London*, 3rd ser.
i. p. 7. n. 14, pl. iii. fig. 23 (1861).

Venezuela.

♂, B.M.

23. **Pedaliodes Pedacia.**

Pronophila Pedacia, *Hewitson, Exot. Butterf.* iv. pl. incog. fig. 24
(1868).

Ecuador.

Coll. Hewitson.

24. *Pedaliodes Prytanis*.

Pronophila Prytanis, *Hewitson, Trans. Ent. Soc. London*, 3rd ser.
i. p. 7. n. 15, pl. iii. fig. 24 (1861).

Venezuela. ♂, B.M.

Var. *a*. *Fascia subanali posticarum subtus brevior et magis quadrata.*

Venezuela. ♀? B.M.

Var. *b*. *Fascia elongata triangulari, velut in Piletha.*

Venezuela. ♂, B.M.

This variety, as remarked by Mr. Hewitson, has the elongated triangular band of *Piletha*; it agrees, however, with *Prytanis* in form and size.

25. *Pedaliodes Phœnissa*.

Pronophila Phœnissa, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 9, pl. 4. fig. 30 (1861).

Bogota. ♂, B.M.

Differs from the preceding species in the more feebly scalloped margin, and in having brown fringes to the wings.

26. *Pedaliodes Physcoa*.

Pronophila Physcoa, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 5, pl. 3. fig. 17 (1861).

Bolivia. ♂, B.M.

27. *Pedaliodes Porina*.

Pronophila Porina, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 9. n. 18, pl. 4. fig. 28 (1861).

Bolivia. ♂ ♀, B.M.

28. *Pedaliodes Peucestas*.

Pronophila Peucestas, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 4. n. 7, pl. 2. figs. 13, 14 (1861).

(♂) Bogota, (♀) Quito. B.M.

29. *Pedaliodes Pallantis*.

Pronophila Pallantis, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 5. n. 8, pl. 2. figs. 15, 16 (1861).

Bogota. ♀, B.M.

30. *Pedaliodes Pylas*.

Pronophila Pylas, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 4. n. 6, pl. 2. figs. 11, 12 (1861).

Bogota. ♂, B.M.

31. **Pedaliodes Cocytia.**

Pronophila Cocytia, *Felder, Reise der Fregatte Novara*, iii. p. 468. n. 796 (1867).

Pronophila Phaesana, *Hewitson, Exot. Butterf.* iv. pl. incog. fig. 23 (1868).

Bogota (Felder).

Belongs to the *Parepa* group; the wings are less strongly dentated, the front wing somewhat waved below the apex, and the inner margin of the hind wing shorter than in all the other species. Its nearest ally is *P. Plotina*, Hew.

32. **Pedaliodes Plotina.**

Pronophila Plotina, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 4. n. 5, pl. 2. figs. 9, 10 (1861).

Venezuela.

♂, B.M.

33. **Pedaliodes Parepa.**

Pronophila Parepa, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 2. n. 1, pl. 1. figs. 1, 2 (1861).

Guayaquil.

♂, B.M.

34. **Pedaliodes Phila.**

Pronophila Phila, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 2. n. 2, pl. 1. figs. 3, 4 (1861).

Bolivia.

♂, B.M.

35. **Pedaliodes Phœnicusa.**

Pronophila Phœnicusa, *Hewitson, Exot. Butterf.* iv. pl. incog. figs. 18, 19 (1868).

Ecuador.

Coll. Hewitson.

36. **Pedaliodes Phæa.**

Pronophila Phæa, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 3. n. 3, pl. 1. figs. 5, 6 (1861).

Colombia.

♂, B.M.

Mr. Hewitson has figured a very distinct looking insect as a variety of this species (*Exot. Butterf.* iv. pl. incog. fig. 26, 1861).

37. **Pedaliodes ochrotænia.**

Pronophila ochrotænia, *Felder, Reise der Fregatte Novara*, iii. p. 467. n. 795 (1867).

Var. *β. fuscata*, *Felder, ibid.*

Bogota, New Granada.

B.M.

38. *Pedaliodes Philotera*.

Pronophila Philotera, *Hewitson, Exot. Butterf.* pl. incog. figs. 16, 17 (Jan. 1868).

Ecuador ?

Coll. Hewitson.

39. *Pedaliodes Peruda*.

Pronophila Peruda, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 3. n. 4, pl. 1. figs. 7, 8 (1861).

Nauta (Amazon).

♂, B.M.

Subgenus 1.

Alis posticis costa apud apicem convexe producta.

40. *Pedaliodes Panyasis*.

Pronophila Panyasis, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 7. n. 13, pl. 3. fig. 22 (1861).

Venezuela.

♂, B.M.

This insect, in the underside-markings, somewhat resembles some species of *Steroma*.

41. *Pedaliodes Napæa*.

Pronophila Napæa, *H. W. Bates, Ent. Month. Mag.* i. p. 179. n. 70 (1865).

Guatemala (central valleys).

Coll. Salvin.

42. *Pedaliodes Phanaræa*.

Pronophila Phanaræa, *Hewitson, Exot. Butterf.* iv pl. incog. fig. 15 (1868).

Ecuador.

Coll. Hewitson.

Genus 65. *GYROCHEILUS*, *Bulter* (1867)*.

Typical Species *Gyrocheilus Patrobas*.

1. *Gyrocheilus Patrobas*.

Pronophila Patrobas, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 16. n. 38, pl. 6. figs. 40, 41 (1861); *Exot. Butterf.* iv. pl. incog. fig. 20 (1868).

Mexico (Oajaca).

♂, B.M.

* This and the other genera of the *Pronophila* group were described, and their distinctive characters figured, in the 'Annals and Magazine of Natural History' for 1867.

Genus 66. OXEOSCHISTUS, *Butler* (1867).Typical Species *Oxeoschistus Puerta*.1. *Oxeoschistus Puerta*.

Pronophila Puerta, *Westwood* in *Gen. Diurn. Lepid.* p. 358.
n. 3 (1851); *Hewitson* in *Exot. Butterf.* ii. p. 28, pl. 41. fig.
12 (1859).

Venezuela.

♂ ♀, B.M.

2. *Oxeoschistus simplex*, sp. nov. Pl. IV. fig. 2.

Alæ anticæ costa subconvexa, margine externo fusco tenuiore;
posticæ latius ferrugineæ, maculis nigris majoribus; aliter velut in
O. Puerta.

Anticæ subtus obscuriores: posticæ fascia media latiore, ocellis
ferrugineo quadrate inclusis, fascia marginali haud denticulata:
aliter velut in *O. Puerta*.

Exp. alar. unc. $2\frac{1}{2}$.

Colombia.

♂, B.M.

3. *Oxeoschistus hilarus*.

Pronophila hilara, *Bates*, *Ent. Month. Mag.* i. p. 178, n. 60 (1865).

Guatemala (Pacific slope).

Coll. Salvin.

4. *Oxeoschistus Protogenia*.

♂ ♀. *Pronophila Protogenia*, *Hewitson*, *Trans. Ent. Soc. Lon-*
don, 3rd ser. i. p. 13. n. 28, pl. 6. fig. 38 (1861).

Bogota.

♂, B.M.

5. *Oxeoschistus pronax*.

Pronophila pronax, *Hewitson*, *Exot. Butterf.* ii. p. 82, pl. 41.
figs. 10, 11 (1859).

Nauta (Amazon).

♂, B.M.

6. *Oxeoschistus Propylea*.

Pronophila Propylea, *Hewitson*, *Exot. Butterf.* ii. p. 80, pl. 40.
figs. 5, 6 (1857).

New Granada.

Coll. Hewitson.

7. *Oxeoschistus Tauropolis*.

Pronophila Tauropolis, *Hewitson & Westwood* in *Gen. Diurn.*
Lepid. p. 358. n. 6, pl. 66. fig. 1 (1851).

Pronophila lætifera, *H. W. Bates*, *Ent. Month. Mag.* i. p. 164.
n. 68 (1865).

Mexico, Guatemala.

8. *Oxeoschistus Prochyta*.

Pronophila Prochyta, *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 13. n. 29, pl. 6. fig. 39 (1861).

Bolivia.

♂, B.M.

9. *Oxeoschistus Irmina*.

♂. *Pronophila Irmina*, *Hewitson & Westwood in Gen. Diurn. Lepid.* p. 358. n. 5, pl. 60. fig. 2 (1851); ♀, *Herrich-Schäffer, Lep. Exot.* figs. 85, 86.

Venezuela.

♂ ♀, B.M.

Genus 67. *LASIOPHILA*, *Felder* (1859).

Typical Species *Lasiophila Zapatoza*.

1. *Lasiophila Cirta*.

♂. *Lasiophila Cirta*, *Felder, Wien. ent. Monatsch.* iii. Taf. 6. fig. 1 (1859).

Pronophila Cirta (part.), *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 15. n. 37 (1861).

♂ as ♀. *Pronophila Præneste* (part.), *Hewitson, Exot. Butterf.* ii. p. 81, pl. 41. figs. 7, 9 (not 8) (1859).

Peru (*Felder*), New Granada (*Hewitson*), Nauta (Amazon).

♂, B.M.

2. *Lasiophila Præneste*.

♂. *Pronophila Præneste*, *Hewitson, Exot. Butterf.* ii. p. 81, pl. 41. fig. 8 (1859).

? ♂. *Lasiophila Circe*, *Felder, Wien. ent. Monatsch.* iii. n. 26 (1859).

Pronophila Cirta (part.), *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i. p. 15. n. 37 (1861).

Bogota.

♂, B.M.

3. *Lasiophila Zapatoza*.

♂. *Pronophila Zapatoza*, *Westwood in Gen. Diurn. Lepid.* p. 358. n. 4 (1851); ? *Herrich-Schäffer, Lep. Exot.* figs. 87, 88 (1850); *Hewitson, Exot. Butterf.* ii. p. 79, pl. 40. figs. 1, 2 (1857).

Venezuela.

♂ ♀, B.M.

The female of this species differs from the male in the brighter colouring of the upperside and the paler colouring of the underside. There do not appear to be any specimens in the collection that exactly agree with the description ("subtus serie punctorum nigrorum, antico albo pupillato"). We have no specimens from Bolivia.

4. *Lasiophila orbifera*, sp. nov. Pl. V. fig. 6.

♂. Alæ supra rufo-ferruginæ: anticæ area apicali nigra, maculis quinque (apicali elongata obliqua) rufo-ferrugineis discalibus serie undata positis: posticæ margine maculisque 5-6 submarginalibus rotundatis, apicalibus subconnectis, nigris.

♀. Major, maculis discalibus et (posticarum) postmacularibus flavidis.

Alæ subtus fere velut in *L. Zapatoza*, Westw.

Bolivia, Bogota.

♂ ♀, B.M.

Var. Alis paulum brevioribus.

Amazon (registered "Nautæ").

♂, B.M.

This species is allied to *Zapatoza*, of which Mr. Hewitson considers it to be a variety; but it is larger, and has the front wings black to beyond the termination of the discoidal cell. For some time it stood in the collection as Felder's *Circe*; but it does not quite agree with his description, and Mr. Hewitson considers the latter to represent his *Præneste*.

5. *Lasiophila Hewitsonia*, sp. nov. Pl. IV. fig. 4.

♂. Coloribus alarum anticarum et forma velut in *L. Cirta* (Felder), posticarum velut in *Præneste* (Hewitson): alæ subtus velut in *Phalæsia* (Hewitson).

Exp. alar. unc. 2½.

Bolivia.

♂, B.M.

Allied to the preceding species and to *Præneste*; the underside of the wings exactly as in *Phalæsia*. This species is considered by Mr. Hewitson to be a variety of *Præneste*; it, however, chiefly resembles the latter species in form; and, were the pattern of its coloration more distinctly outlined, nobody would hesitate about separating them. It is far more distinct than many of the species in the *Pedaliodes* group.

6. *Lasiophila Prosymna*.

♂. Pronophila Prosymna, Hewitson, Exot. Butterf. ii. p. 79, pl. 40, figs. 3, 4 (1857); as ♀, Trans. Ent. Soc. London, 3rd ser. i. p. 15. n. 35 (1861).

Bogota.

♂, B.M.

7. *Lasiophila Phalæsia*.

♂. Pronophila Phalæsia, Hewitson, Exot. Butterf. iv. pl. incog. figs. 13, 14 (Jan. 1868).

Var. Pronophila Prosymna (part.), Hewitson, Trans. Ent. Soc. London, 3rd ser. i. p. 15. n. 35, pl. 5. figs. 35, 36 (1861).

Ecuador (Hewitson), Bolivia, Quito.

♂, B.M.

Genus 68. *DÆDALMA*, *Hewitson* (1858).Typical Species *Dædalma Dinias*.1. *Dædalma Dinias*.

♂ ♀. *Dædalma Dinias*, *Hewitson*, *Exot. Butterf.* ii. p. 85, pl. 43.
figs. 1-3 (1858).

Bogota.

♂, B.M.

2. *Dædalma inconspicua*.

♂. *Dædalma inconspicua*, *Butler*, *Ent. Month. Mag.* iii. p. 77.
n. 2 (1866).

Quito.

♂, B.M.

3. *Dædalma Drusilla*.

♂. *Dædalma Drusilla*, *Hewitson*, *Exot. Butterf.* ii. p. 86, pl. 43.
fig. 7 (1858).

New Granada.

Coll. Hewitson.

4. *Dædalma Doræte*.

♂. *Dædalma Doræte*, *Hewitson*, *Exot. Butterf.* ii. p. 85, pl. 43.
figs. 4, 5 (1858).

New Granada.

Colls. Hewitson & Saunders.

5. *Dædalma Dorinda*.

Dædalma Dorinda, *Felder*, *Wien. ent. Monatschr.* vi. p. 28
(1862); *Nov. Voy.* iii. tab. 67. figs. 3, 4 (1867).

Bogota.

Coll. Hewitson.

6. *Dædalma Emilia*.

Dædalma Emilia, *Butler*, *Proc. Zool. Soc.* p. 40. n. 3, pl. 3. fig. 3
(1866).

Bogota.

B.M.

This may be only a brightly coloured variety of *Dorinda*; it seems, however, to differ slightly in the form of the hind wings.

Subgenus 1.

Costa posticarum subintegra, *caudis aliquando obsoletis*.

7. *Dædalma Phoronea*.

♂. *Pronophila Phoronea*, *Hewitson & Westwood*, *Gen. Diurn. Lepid.* p. 358. n. 7, pl. 60. fig. 1 (1861); also of *Hewitson* (1861).

Venezuela.

♂ ♀, B.M.

8. *Dædalma Phaselis*.

♂. *Pronophila Phaselis*, *Hewitson, Trans. Ent. Soc. London*, ser. 3. i. p. 14, pl. 6. fig. 37 (1861).

Venezuela.

♂, B.M.

Dædalma ? Drymæa (? *Pedaliodes*, near *angularis*).

Dædalma Drymæa, *Hewitson, Exot. Butterf.* ii. p. 86, pl. 43. fig. 6 (1858).

New Granada.

Coll. Hewitson.

Genus 69. *PRONOPHILA*, *Westwood* (1851).

Typical Species *Pronophila Thelebe*.

1. *Pronophila Thelebe*.

Pronophila Thelebe, *Hewitson & Westwood in Gen. Diurn. Lepid.* p. 358. n. 1, pl. 60. fig. 3 (1851); *Hewitson, Trans. Ent. Soc. London*, ser. 3. i. p. 11. n. 23 (1861).

Venezuela.

B.M.

2. *Pronophila Orcus*.

Satyrus Orcus, *Latreille in Humboldt & Bonpland's Obs. Zool.* ii. p. 72, pl. 35. figs. 1, 2 (1799–1804). *Pron. Orcus*, *Westw.* (1851).

Bogota.

B.M.

Var. Alis anticis supra velut in Thelebe albimaculatis.

Bolivia.

B.M.

3. *Pronophila Porsenna*.

Pronophila Porsenna, *Hewitson, Trans. Ent. Soc.* 3rd ser. i. p. 12. n. 25, pl. 5. fig. 34 (1861).

Quito, Bogota.

B.M.

4. *Pronophila obscura*, sp. nov. Pl. IV. fig. 10.

Alæ supra magnitudine et colore Thelebes, forma Orci.

Alæ subtus pallidiores: anticæ maculis elongatis discalibus nigris, maculas quatuor nigras cæruleo-albido pupillatas includentibus, serie directa digestas, et partim aurantiaco a lateribus marginatas: apice albicante: posticæ coloribus Orci, fasciarum forma et ocellorum positione magis iis Cordillerae simulantibus.

Exp. alar. unc. 3 $\frac{1}{8}$ –2 $\frac{7}{8}$.

Venezuela (two specimens).

B.M.

Most nearly allied to *Cordillera*, which it resembles in the number and position of its ocelli. The hind wings below are most like those of *P. Orcus*.

5. *Pronophila Cordillera*.

Pronophila Cordillera, *Westwood in Gen. Diurn. Lepid.* p. 358.
n. 2 (1851); *Hewitson, Trans. Ent. Soc. London*, 3rd ser. i.
p. 11. n. 24, pl. 5. fig. 33 (1861).

Bolivia.

B.M.

Genus 70. *CORADES*, *E. Doubleday* (1848).

Typical Species *Corades Enyo*.

1. *Corades Iduna*.

Corades Iduna, *Hewitson in Ann. & Mag. Nat. Hist.* 2nd ser. vi.
p. 437 (1850); *Hewitson & Westwood, Gen. Diurn. Lepid.*
p. 355. n. 2, pl. 63. fig. 1 (1851).

Bolivia.

B.M.

Local form. Macula discali anticarum minore nec geminata: alæ
posticæ longiores, colore flavo confuso maculam permagnam
formante.

Nauta (Amazon).

B.M.

2. *Corades Chelonis*.

Corades Chelonis, *Hewitson, Exot. Butterf.* iii. p. 71, pl. 36.
figs. 1, 2 (1863).

Bogota, Quito.

B.M.

3. *Corades Enyo*.

Corades Enyo, *Hewitson, Proc. Zool. Soc. London*, p. 117, pl. 4
(*Annulosa*) (1848).

Corades Auriga, *Herrich-Schäffer, Lep. Exot. sp. nov.* (Regens-
burg, 1856), ser. 11, Lief. 5. n. 82, 83 (1850-58).

Venezuela.

♂ ♀, B.M.

Local form. Alis posticis supra castaneo-fuscis.

Nauta (Amazon), Bogota, New Granada, Bolivia.

B.M.

4. *Corades Pannonia*.

Corades Pannonia, *Hewitson, Ann. & Mag. Nat. Hist.* 2nd ser.
vol. vi. p. 438, pl. 10. fig. 1, not 2 (1850); *Hewits. & West-*
wood, Gen. Diurn. Lepid. p. 355. n. 3 (1851).

Corades ichthya, *Herrich-Schäffer, Lep. Exot. sp. nov.* (Regens-
burg, 1856), ser. 11, Lief. 5. n. 84 (1850-58).

Venezuela.

B.M.

Local form. Alae subtus fascia media ochrea posticarum subobsoleta; punctis anticarum rufis obsoletis.

Bogota.

B.M.

5. *Corades Ulema*.

Corades Ulema, Hewitson in *Ann. & Mag. Nat. Hist.* 2nd ser. vi. p. 438 (1850); Hewitson & Westwood in *Gen. Diurn. Lepid.* p. 355. n. 4 (1851).

Bolivia.

B.M.

6. *Corades Chirone*.

Corades Chirone, Hewitson, *Exot. Butterf.* iii. p. 71, pl. 36. fig. 3 (1863).

Bogota, Quito.

♂, B.M.

7. *Corades argentata*, sp. nov.

Corades Pannonia ♀?, Hewitson in *Ann. & Mag. Nat. Hist.* 2nd ser. vi. p. 438, pl. 10. fig. 2 (1850).

Bolivia.

♂, B.M.

This species is undoubtedly distinct from *Pannonia*. It differs from it on the underside in the form of the central band of the hind wing, which, as Mr. Hewitson remarks, is "of silver," not of yellow, as in *Pannonia*. Our specimen is an undoubted male, so that the idea of its being the opposite sex of Mr. Hewitson's species at once falls to the ground. It is more nearly allied to *C. Cybele* than to *Pannonia*.

8. *Corades Cybele*.

Corades Cybele, Butler, *Proc. Zool. Soc. London*, p. 40. n. 2, pl. 3. fig. 2 (1866).

Bogota.

B.M.

9. *Corades Sareba*.

Corades Sareba, Hewitson, *Ann. & Mag. Nat. Hist.* 2nd ser. vi. p. 439, pl. 10. fig. 5 (1850); Hewitson & Westwood, *Gen. Diurn. Lepid.* p. 355. n. 6 (1851).

Bolivia.

B.M.

10. *Corades Medeba*.

Corades Medeba, Hewitson, *Ann. & Mag. Nat. Hist.* 2nd ser. vi. p. 439, pl. 10. fig. 4 (1850); Hewitson & Westwood, *Gen. Diurn. Lepid.* p. 355. n. 5 (1851).

Bolivia, Quito.

♂, B.M.

Local form. *Alæ subtus striga discoidali abbreviata.*

Bogota.

♂ ♀, B.M.

The female of this species has several reddish spots, and, in the front wings, a row of ill-defined ochreous spots on the upperside.

11. *Corades Cistene.*

Corades Cistene, *Hewitson, Exot. Butterf.* iii. p. 72, pl. 36. figs. 4, 5 (1863).

Bogota, Quito.

B.M.

Genus 71. *PTYCHANDRA*, *Felder* (1861).

Typical Species *Ptychandra Lorquinii*.

1. *Ptychandra Lorquinii.*

Ptychandra Lorquinii, *Felder, Wien. ent. Monatschr.* v. p. 304 (1861); *Nov. Voy.* iii. tab. 68. figs. 1-3 (1867).

Philippine Islands.

B.M.

This genus differs from every other genus of *Satyridæ* in the shortness of the discoidal cell of the front wings. In this particular it resembles the family *Eurytelidæ*.

2. *Ptychandra leucogyne.*

Ptychandra leucogyne, *Felder, Novara-Voyage*, iii. p. 498. n. 864 (1867).

Gilolo.

ADDITIONAL SPECIES.

1. *Satyrus Boöpis*.

Satyrus Boöpis, Behr, *Am. Calif. Acad. Nat. Scien.* (1864);
Edwards, Proc. Ent. Soc. Philad. p. 196. n. 4 (1866).

California.

Possibly a variety of *Satyrus* (*Minois*) *Nephele*, from which it chiefly differs in the absence of ocelli upon the under surface of the wings.

2. *Cænonympha infuscata*.

Cænonympha infuscata, M'Leay, in *Proc. Ent. Soc. New South Wales*, i. pt. 5. p. liii (1866).

Cape York.

The description of this species is very brief; but its locality will probably assist in its determination. All species of *Satyridae*, however, ought to be figured.

3. *Lethe? marginalis*.

Satyrus marginalis, Motschulsky, *Etudes Entomologiques*, ix. p. 29 (1860).

Japan.

May be nearly allied to *Lethe Diana*, Butler; but the description does not quite agree with it. It may possibly be a faulty description taken from a broken specimen. The work in which this species is described has not been recorded. *Satyrus bipunctatus*, Motschulsky, *Etudes Entom.* ix. p. 29 (1860)=Japanese form of *Satyrus* (*Minois*) *Dryas* of Scopoli.

4. *Aulocera Werang*.

Aulocera Werang, Lang, in *Ent. Month. Mag.* iv. p. 247 (1868).

Upper Kunáwur (Werang Pass), Cashmere (Goolmurg).

Most nearly allied to *A. Scylla*, Butler.

5. *Erites argentina*, sp. nov. Pl. V. fig. 8.

♀. Alæ supra fuscæ pallidæ, subhyalinæ: anticæ ocellis quinque, quinto maximo subanali, primo et quarto minimis, omnibus cæcis flavo cinctis: posticæ ocellis quatuor similibus sed magis distinctis, quarto paulum majore apud angulum analem; fasciis velut in *E. elegante*, sed magis distinctis.

Alæ subtus argenteo-glaucinae, area basali fusco liturata; fasciis

duabus mediis nigro limitatis fulvis, externa posticarum latiore, clariore et medio extus producta; margine externo albido lineis duabus ciliisque nigris: anticæ ocellis quinque nigerrimis pallide fulvo cinctis, nigro circumcinctis et argenteo decore pupillatis, magnitudine velut supra, distantibus: posticæ ocellis quinque nigerrimis, secundo et quinto paulum majoribus; primo et secundo velut in anticis, tertio et quinto flavo cinctis et nebula flavo-fusca circumcinctis, quarto minimo nebula flava circumcincta.

Exp. alar. unc. $2\frac{3}{8}$.

Labuan (Borneo).

♀, B.M.

E. argentina may be looked upon as nearly intermediate between *E. Madura* and *E. elegans*; it differs, however, from both in the form of the front wings, which in this species are much broader and more rounded at the apex. The large ocellus of the front wings seems to indicate its nearer relationship to *Madura*; but the general colouring of the underside is more like that of *elegans*. The form of the central bands, and the relative size of the ocelli, seem to preclude the possibility of *E. elegans* and *argentina* being opposite sexes of one species.

FOSSIL SPECIES.

(CYLLO) SEPULTA.

Cylo sepulta, Boisduval in *Ann. Ent. Soc. France*, p. 371, pl. 8 (1840); also of Westwood (1851).

Vanessa sepulta, Lefebvre in *Ann. Ent. Soc. France*, p. 71, pl. 3. figs. A-C (1851).

"Dans les marnes des environs d'Aix en Provence" (Boisduval).

This very interesting species was described, and admirably figured by Dr. Boisduval in the French 'Annales de la Société Entomologique' (1840); that gentleman considered it to be a Satyride allied to *Satyrus rohria*, *caumas*, &c.*, which it somewhat resembles in the form of the wings.

In the French 'Annales' (for 1851), M. Lefebvre published a note upon the species, in which he criticised Dr. Boisduval's paper, and stated that the fossil species, instead of being allied to *rohria*, was evidently a *Vanessa*—that the strong tail-like wing projection belonged to the front, and not to the hind wings, and represented the angular projection which occurs in all true *Vanessidae*, as an example of which he instanced *Vanessa (Junonia) Archesia* of Cramer. This remarkable note was, moreover, accompanied by figures of the species, representing the tail both upon the front and hind wings.

In the same volume of the 'Annales' Dr. Boisduval gives an excellent answer to M. Lefebvre's observations, in which he well remarks, "Nous n'avons jamais vu une seule espèce avec les ailes

* Species of *Lethe*.

supérieures anguleuses et appendiculées, et les ailes inférieures arrondies comme avec un compas ;” and certainly, did such an insect ever exist, its wings would be utterly useless as organs of flight, for they would invariably carry it downwards. In all insects which have small and rounded hind wings the costa of the front wings always far exceeds the inner margin in length and strength*, whereas in *M. Lefebvre’s* insect the reverse would be the case.

The true position of *C. sepulta* is undoubtedly in the family *Satyridae*; and, so far as can be judged from the beautiful figure in the ‘*Annales*’†, it is exactly intermediate in character between three nearly allied genera now existing, viz. *Neorina*, *Antirrhæa*, and *Anchiphlebia*, its more immediate allies being the commonest species in each of the above genera. Its characters are distributed between these three species as follows :—

	<i>Neorina</i> <i>Lewsi</i> , Boisd.	<i>Antirrhæa</i> <i>Philottetes</i> , Linn.	<i>Anchiphlebia</i> <i>Archæa</i> , Hübner.
Form of front wings	*		
Form of hind wings		*	
Tails of hind wings (intermediate in character between)	*	*	
Submarginal spots of front wings	*	*	*
Black disco-submarginal spots of hind wings		*	
Pale costal and discal banding of wings	*?		*
Limitation of dark dentated basal area of hind wings			*
Submarginal line of hind wings	*		

The venation appears to be nearly similar to that of *Anchiphlebia* (pl. 5. fig. 3). It is doubtful, however, whether the drawing of the veins has been sufficiently attended to to offer any reliable characters.

The species in this Catalogue, as they now stand, amount to upwards of 760, more than 400 of which are in the National Collection. Of these, 51 are described as new to science.

* As, for instance, in the *Sphingidæ*, *Heliconidæ*, &c.

† In this figure the neuration has not been very clearly defined, the veining of the hind wings not being continuous.

INDEX.

The genera are printed in Roman capitals, the species in Roman, the synonyms in italics, and the more distinct varieties or local forms are marked with an asterisk.

A.		Page		Page
Abd-el-Kader, <i>Pierret</i> (Satyrus)...	61		<i>Agrota</i> , <i>Butler</i> (Euptychia)	29
Abeona, <i>Donovan</i> (Papilio)	70		Aello, <i>Esper</i> (Pap. Nymph. Gemm.)	160
<i>Abeona</i> , <i>Butler</i> (Xenica)	70		Aello, <i>Boisduval</i> (Chionobas)	160
E. Doubleday (Hipparchia)	70		Godart (Satyrus)	160
Godart (Satyrus)	70		Hübner (Eumenis & Papilio)	160
Wallengren (Heteronympha)	70		<i>atherius</i> , <i>Esper</i> (Pap. Nymph. Gemm.)	75
Westwood (Lasiommata) ...	70		<i>Æthiops</i> , <i>Butler</i> (Mycalesis)	141
<i>Acanthe</i> , <i>Hübner</i> (Tisiphone).....	166		<i>Æthiops</i> , <i>Herbst</i> (Papilio)	81
<i>Achanta</i> , <i>Donovan</i> (Papilio)	166		<i>Æthiops minor</i> , <i>Villars</i> (Papilio)	73
<i>Achanta</i> , <i>Butler</i> (Geitoneura) ...	166		Afer, <i>Esper</i> (Pap. Nymph. Gemm.)	91
Godart (Satyrus)	166		Afer, <i>Butler</i> (Erebia)	91
Moore (Xenica)	166		Duponchel (Satyrus)	91
Westwood (Lasiommata) ...	166		Eversmann (Hipparchia) ...	91
Achine, <i>Scopoli</i> (Papilio)	122		Freyer (Pap. Hipparchia) ...	91
<i>Achine</i> , <i>Butler</i> (Pararge).....	122		Schneider (Papilio)	91
Acmenis, <i>Hübner</i> (Megisto)	24		affinis, <i>Butler</i> (Euptychia)	20
<i>Acmenis</i> , <i>Butler</i> (Euptychia)	24		Afra, <i>Boisduval</i> (Erebia)	91
Westwood (Neonympha) ...	24		Fabricius (Papilio)	91
<i>Acresida</i> , <i>Butler</i> (Lymanopoda)...	171		Godart (Satyrus)	91
ACROPTALMIA (sic), <i>Felder</i>	159		* <i>Aganippe</i> , <i>Felder</i> (Zethera)	109
Actæa, <i>Esper</i> (Pap. Nymph. Gemm.)	59		Agatha, <i>Butler</i> (Euptychia)	34
<i>Actæa</i> , <i>Borkhausen</i> (Papilio).....	59		Agave, <i>Borkhausen</i> (Papilio)	52
Butler (Hipparchia)	59		Esper (Pap. Nymph. Gemm.)	52
Fabricius (Papilio)	61		<i>Aglauros</i> , <i>Herbst</i> (Papilio).....	87
Godart (Satyrus)	59		Agnes, <i>Cramer</i> (Papilio).....	97
Lang (Papilio)	69		albinotata, <i>Butler</i> (Taygetia)	12
De Villers (Papilio)	57		albocincta, <i>Hewitson</i> (Lymanopoda)	171
<i>Actæus</i> , <i>Herbst</i> (Papilio)	59		albo-maculata, <i>Hewitson</i> (Lymanopoda)	171
<i>Adiantæ</i> , <i>Hübner</i> (Neonympha)...	167		Alcinö, <i>Felder</i> (Euptychia)	25
Adolphe, <i>Güérin</i> (Satyrus)	139		Alcyone, <i>Denis</i> (Papilio)	50
<i>Adolphe</i> , <i>Westwood</i> (Mycalesis)	139		Alcyone, <i>Boisduval</i> (Satyrus)	50
Adrasta, <i>Hübner</i> (Papilio)	124		Borkhausen (Papilio).....	73, 75
<i>Adrasta</i> , <i>Butler</i> (Amecera).....	124		Butler (Hipparchia)	50
<i>Adyte</i> , <i>Hübner</i> (Papilio).....	82		Fabricius (Papilio)	52
<i>Ægeria</i> , <i>Linnaeus</i> (Papilio).....	120		Godart (Satyrus)	52
<i>Ægeria</i> , <i>Butler</i> (Pararge)	120		Stewart (Papilio)	81
E. Doubleday (Lasiommata)	120		Alecto, <i>Hübner</i> (Papilio)	78
Esper (Pap. Nymph. Gemm.)	120		Alecto, <i>Boisduval</i> (Erebia)	77
Godart (Satyrus)	120		Freyer (Pap. Hipparchia) ...	77
Stephens (Hipparchia)	120		Hübner (Papilio)	77
Westwood (Lasiommata) ...	120			

	Page		Page
<i>Alexis</i> , Esper (Pap. Nymph. Gemm.)	83	<i>Annada</i> , F. Moore (Erebia)	96
Linnaeus (Papilio)	83	Butler (Callerebia)	96
<i>Allionia</i> , Cyrrilo (Papilio)	57	<i>anophthalma</i> , Felder (Taygetis)	14
Esper (Pap. Nymph. Gemm.)	57	<i>Anosia</i> , Godart (Satyrus)	71
<i>Allionii</i> , Hübner (Papilio)	57	<i>Anthe</i> , Godart (Satyrus)	55
<i>Allionius</i> , Herbst (Papilio)	57	Ochsenheimer (Papilio)	55
<i>Alope</i> , Fabricius (Papilio)	62	<i>Anthelea</i> , Hübner (Papilio)	51
<i>Alope</i> , E. Doubleday (Enodia?)	62	<i>Anthelea</i> , Boisduval (Satyrus)	52
Godart (Satyrus)	62	E. Doubleday (Hipparchia)	51
Reakirt (Enodia)	62	Freyer (Pap. Hipparchia)	51, 52
Westwood (Satyrus Enodia)	62	Herrich-Schäffer (Satyrus)	51
<i>Also</i> , Boisduval (Chionobas)	163	<i>Antipodum</i> , E. Doubleday (Argyrophenga)	159
<i>amabilis</i> , Boisduval (Cyllo)	5	<i>Antirius</i> , Butler (Hypocysta)	168
<i>amabilis</i> , Butler (Hipo)	5	<i>ANTIRRHÆA</i> , Westwood	107
<i>Amarillis</i> , Ernst (Papilio)	69	<i>Antonoë</i> , Cramer (Papilio)	30
Herbst (Papilio)	43	<i>Antonoë</i> , Butler (Euptychia)	30
<i>Amaryllis</i> , Cramer (Papilio)	43	Herbst (Papilio)	53
Geoffroy (Papilio)	68	Westwood (Euptychia)	31
<i>Amaryllis</i> , Butler (Epinephele)	68	<i>Anysis</i> , Hewitson (Debis)	117
Eversmann (Hipparchia)	43	<i>Anysis</i> , Butler (Lethe)	117
Freyer (Pap. Hipparchia)	43	<i>Aphnius</i> , Godart (Satyrus)	152
Godart (Satyrus)	69	<i>Aphnius</i> , Hewitson (Ypthima)	152
Herrich-Schäffer (Satyrus)	43	Westwood (Ypthima)	152
Westwood (Cænonympha)	43	<i>apiculata</i> , Felder (Lymanopoda)	170
<i>Ambasara</i> , Moore (Melanitis)	4	<i>Apuleja</i> , Felder (Pronophila)	175
<i>ambigua</i> , Butler (Euptychia)	21	<i>Apuleja</i> , Butler (Pedaliodes)	175
<i>Ambiorix</i> , Wallengren (Neosatyrys)	96	<i>Arachne</i> , Fabricius (Papilio)	80
<i>AMECERA</i> , Butler	123	<i>Arachne</i> , Butler (Erebia)	80
<i>AMPHIDECTA</i> , Butler	7	Denis (Papilio)	56
<i>Amphithea</i> , Menétrés (Ypthima)	148	Esper (Pap. Nymph. Gemm.)	56
<i>Amphithea</i> , Hewitson (Ypthima)	149	Godart (Satyrus)	80
<i>Amphitrite</i> , Hübner (Papilio)	157	Hübner (Papilio)	81
<i>Amphitrite</i> , Godart (Satyrus)	157	<i>Aramis</i> , Hewitson (Mycalesis)	129
Hübner (Arge)	157	<i>Aranea</i> , Fabricius (Papilio)	33
Ochsenheimer (Papilio)	156	Godart (Satyrus)	33
<i>Amyntas</i> , Poda (Papilio)	40	<i>Arcadia</i> , Cramer (Papilio)	118
<i>Amyntas</i> , Butler (Cænonympha)	40	<i>Arcadia</i> , Butler (Lethe)	118
Scopoli (Papilio)	41	Hübner (Temenis)	118
<i>Anacardii</i> , Fabricius (Papilio)	102	Westwood (Debis)	118
<i>ANADEBIS</i> , Butler	110	<i>Arcania</i> , Linnaeus (Papilio)	41
<i>Anapita</i> , Moore (Mycalesis)	146	<i>Arcania</i> , Eversmann (Hipparchia)	41
<i>Anaxias</i> , Hewitson (Mycalesis)	141	Hübner (Papilio)	41
<i>Anaxogorus</i> , Asamuss (Cænonympha)	41	Hufnagel (Pap. Nymph. Gemm.)	41
<i>ANCHIPHLEBIA</i> , Butler	106	<i>arcanioides</i> , Pierret (Satyrus)	41
Andensis, Felder (Steroma)	172	<i>arcanioides</i> , Butler (Cænonympha)	41
<i>Andromacha</i> , Hübner (Oreas)	114	<i>Arcanius</i> , E. Doubleday (Cænonympha)	41
<i>Andromeda</i> , Cramer (Papilio)	11	Esper (Pap. Nymph. Gemm.)	41
Fabricius (Papilio)	101	Godart (Satyrus)	41
<i>Andromeda</i> , Butler (Callitæra)	101	Linnaeus (Papilio)	41
E. Doubleday (Satyrus)	11	Stephens (Hipparchia)	41
Godart (Satyrus)	101	<i>Arcensia</i> , Cramer (Papilio)	2
Hübner (Cithærias)	101	<i>Arcesilaë</i> , Hübner (Cærois)	1
Westwood (Hætera)	101	<i>Arcesilaus</i> , Cramer (Papilio)	1
Westwood (Taygetis)	11	<i>Archæa</i> , Hübner (Antirrhæa)	106
<i>angularis</i> , Butler (Euptychia)	19	<i>Archæa</i> , Butler (Anchiphlebia)	106
Butler (Pedaliodes)	176	Westwood (Hætera)	106
<i>angulosa</i> , Butler (Mycalesis)	130		

	Page		Page
<i>Archebates, Ménétriés</i> (Satyrus)...	97	<i>Aristeus</i> , Westwood (Satyrus) ...	56
<i>Archebates</i> , Butler (Calisto)	97	<i>armilla</i> , Butler (Euptychia)	22
Westwood (Euptychia)	98	<i>armillata</i> , Butler (Taygetis)	8
<i>Archemor</i> , Godart (Satyrus)	100	<i>Artemis</i> , Felder (Acrophtalmia)...	159
* <i>arctoides</i> , Hewitson (Ypthima)	151	<i>ascanioides</i> , Westwood (Cæno-	
<i>Arctous</i> , Fabricius (Papilio)	151	nympha)	41
<i>Arctous</i> , Donovan (Papilio)	151	<i>Ascanius</i> , Westwood (Cæno-	
Godart (Satyrus)	151	nympha)	41
Hewitson (Ypthima)	151	<i>Asochis</i> , Hewitson (Mycælesis) ...	131
Westwood (Ypthima)	151	<i>Asophis</i> , Hewitson (Mycælesis) ...	144
<i>arcuata</i> , Butler (Lethe)	114, 190	<i>assimilis</i> , Butler (Cneis)	163
<i>areolata</i> , Butler (Euptychia)	37	<i>Asterope</i> , Klug (Hipparchia)	148
<i>areolatus</i> , Abbot (Papilio)	37	<i>Asterope</i> , Hewitson (Ypthima)	148
<i>areolatus</i> , Godart (Satyrus)	37	Westwood (Ypthima)	148
<i>Arete</i> , Fabricius (Papilio)	73	<i>Astyoche</i> , Erichson (Hætera)	104
<i>Arete</i> , Bellier (Erebia)	75	<i>Astyoche</i> , Butler (Pierella)	104
Butler (Oreina)	73	Hewitson (Hætera)	104
Cramer (Papilio)	114	<i>Aswa</i> , Moore (Melanitis)	5
Müller (Papilio)	64	<i>Atalanta</i> , Butler (Euptychia)	23
Westwood (Erebia)	74	<i>Athene</i> , Borkhausen (Papilio)	62
<i>Arethusa</i> , Denis (Papilio)	53	* <i>atratus</i> , Esper (Pap. Nymph.	
<i>Arethusa</i> , E. Doubleday (Hipparchia)	53	Gemm.)	91
Esper (Pap. Nymph. Gemm.)	53	<i>Atrax</i> , Felder (Cyllo)	6
Freyer (Pap. Hipparchia) ...	54	<i>Atrax</i> , Butler (Hippo)	6
Godart (Satyrus)	53	<i>Atropos</i> , Hübner (Papilio)	156
Hübner (Papilio)	54	<i>AULOCERA</i> , Butler	49
<i>Arethusus</i> , Herbst (Papilio)	53	<i>auricruda</i> , Butler (Mycælesis) ...	131
<i>Aretus</i> , Herbst (Papilio)	74	<i>Auriga</i> , Herrich-Schäffer (Co-	
<i>Argante</i> , Cramer (Papilio)	21	rades)	185
<i>Argante</i> , Hübner (Megisto)	21	<i>Aurora</i> , Felder (Hætera)	101
Westwood (Euptychia)	21	<i>Aurora</i> , Butler (Callitera)	101
<i>ARGE</i> , Hübner	153	<i>austera</i> , Butler (Euptychia)	17
<i>Arge</i> , Duponchel (Satyrus)	157	<i>Autonoë</i> , Borkhausen (Papilio)	53
Fabricius (Papilio)	155	E. Doubleday (Hipparchia)	53
Fuessly (Papilio)	157	Esper (Pap. Nymph. Gemm.)	53
Godart (Satyrus)	155	Freyer (Pap. Hipparchia) ...	53
Sulzer (Pap. Nymph. Gemm.)	157	Godart (Satyrus)	53
<i>Arge nemausiaca</i> , Esper (Pap. Nymph. Gemm.)	154	<i>Avatara</i> , Moore (Satyrus)	50
<i>occitanica</i> , Esper (Pap. Nymph. Gemm.)	157	<i>Avatara</i> , Butler (Aulocera)	50
<i>Russica</i> , Esper (Pap. Nymph. Gemm.)	155	<i>Ayaya</i> , Butler (Euptychia)	34
<i>Sicula</i> , Borkhausen (Papilio)	157		
<i>argentata</i> , Butler (Corades)	186		
<i>argenteus</i> , Blanchard (Argyrophorus)	159		
<i>argenteus</i> , Swainson (Satyrus) ...	36		
<i>argentina</i> , Butler (Erites)	188		
<i>Argulus</i> , Godart (Satyrus)	21		
<i>Argus</i> , Butler (Ypthima)	150		
<i>ARGYNNINA</i> , Butler	165		
<i>ARGYROPHENGA</i> , E. Doubleday	159		
<i>ARGYROPHORUS</i> , Blanchard	159		
<i>Argyrospila</i> , Butler (Euptychia)	18		
<i>Ariane</i> , Boisduval (Satyrus)	63		
<i>Ariane</i> , Reakirt (Enodia)	63		
* <i>Aristeus</i> , Bonelli (Pap. Satyrus)	56		
Freyer (Pap. Hipparchia) ...	56		

B.

<i>Baladeva</i> , Moore (Zophoëssa)	109
<i>Balder</i> , Boisduval (Satyrus)	160
Herrich-Schäffer (Chionobas)	160
* <i>Balderi</i> , Hübner (Eumenis)	160
<i>Baldiva</i> , Moore (Lasiommata) ...	127
<i>Baldiva</i> , Butler (Amecera)	127
* <i>Baldus</i> , Fabricius (Papilio)	150
Godart (Satyrus)	150
Hewitson (Ypthima)	150
Westwood (Ypthima)	150
* <i>Banksia</i> , Fabricius (Papilio) ...	3
Butler (Hipparchioides)	100
Butler (Melanitis)	3
E. Doubleday (Lasiommata)	100
Godart (Satyrus)	3
Moore (Melanitis)	3
Westwood (Cyllo)	3

	Page		Page
<i>Cassius</i> , Godart (Satyrus)	93, 94	<i>Clarissa</i> , Hübner (Euptychia) ...	16
Westwood (Erebia)	94	<i>Claudina</i> , Borkhausen (Papilio)...	74
<i>Cassus</i> , <i>Linnaeus</i> (Papilio)	72	* <i>Cleanthe</i> , Boisduval (Arge)	156
<i>Cassus</i> , Godart (Satyrus)	72	Duponchel (Satyrus)	156
Trimen (Leptoneura)	72	<i>Cleo</i> , Hübner (Papilio)	86
Westwood (Erebia)	72	<i>Cleopatra</i> , <i>Felder</i> (Taygetis)	11
<i>Casta</i> , Bates (Antirrhæa)	108	<i>Clerimene</i> , Hübner (Neonympha) ..	37
* <i>Castor</i> , <i>Esper</i> (Pap. Nymph. Gemm.)	87	* <i>Clerimon</i> , <i>Fabricius</i> (Papilio) ...	133
<i>Caumas</i> , Godart (Satyrus)	118	Godart (Satyrus)	133
<i>Celia</i> , <i>Cramer</i> (Papilio)	8	Westwood (Lasiommata) ...	133
<i>Celia</i> , Hübner (Taygetis)	8	<i>Climene</i> , <i>Esper</i> (Pap. Nymph. Gemm.)	121
<i>Celimene</i> , <i>Cramer</i> (Papilio)	161	<i>Climene</i> , Butler (Lasiommata) ...	121
<i>Celmis</i> , Godart (Satyrus)	19	Butler (Pararge)	121
<i>Celmis</i> , Westwood (Euptychia) ...	19	Eversmann (Hipparchia) ...	121
* <i>Celano</i> , Hübner (Papilio)	162	Godart (Satyrus)	121
<i>Cepheus</i> , Butler (Mycæsis)	134	Hübner (Papilio)	121
* <i>Ceres</i> , Butler (Canonympha) ...	47	<i>Clite</i> , Hübner (Papilio)	43
<i>Ceto</i> , Hübner (Erebia)	91	<i>Clorimena</i> , <i>Stoll</i> (Papilio)	36
<i>Ceto</i> , Freyer (Pap. Hipparchia) ...	91	<i>Clorimena</i> , Butler (Euptychia) ...	36
Godart (Satyrus)	91	<i>Clorimene</i> , Herrich-Schäffer (Neonympha)	37
Westwood (Erebia)	91	<i>Clotho</i> , Boisduval (Arge)	155
<i>Ceylonica</i> , <i>Hewitson</i> (Ypthima) ..	152	Duponchel (Satyrus)	156
<i>Ceylonica</i> , Butler (Ypthima)	152	Hübner (Papilio)	155
<i>Chandica</i> , <i>Moore</i> (Debis)	119	<i>Cluena</i> , <i>Drury</i> (Pap. Nymph.) ...	32
<i>Chandica</i> , Butler (Lethe)	119	<i>Cluena</i> , Westwood (Euptychia) ...	32
<i>Cheena</i> , <i>Moore</i> (Epinephele)	65	<i>Clueria</i> , E. Doubleday (Euptychia) ..	32
<i>Chelonis</i> , <i>Hewitson</i> (Corades)	185	Fabricius (Papilio)	32
<i>Chelys</i> , <i>Fabricius</i> (Papilio)	9	Godart (Satyrus)	32
<i>Chelys</i> , Godart (Satyrus)	9	<i>Clymene</i> , <i>Fabricius</i> (Papilio)	121
Westwood (Taygetis)	9	Herrich-Schäffer (Lasiommata)	121
<i>Chenu</i> , <i>Guérin</i> (Satyrus)	151	Westwood (Satyrus)	121
<i>Chenu</i> , Butler (Ypthima)	151	<i>Clymenus</i> , E. Doubleday (Lasiommata)	121
<i>Chilensis</i> , E. Doubleday (Erebia) ..	58	<i>Clyte</i> , Hübner (Dira)	71
<i>Chiliensis</i> , <i>Guérin</i> (Satyrus)	58	<i>Clytus</i> , <i>Linnaeus</i> (Papilio)	71
<i>Chiliensis</i> , Butler (Hipparchia) ...	58	<i>Clytus</i> , E. Doubleday (Erebia) ..	71, 72
Westwood (Erebia)	58	Esper (Pap. Nymph. Gemm.) ..	71
<i>Chione</i> , <i>Felder</i> (Acrophthalmia) ..	159	Godart (Satyrus)	71
<i>Chirone</i> , <i>Hewitson</i> (Corades)	186	Wallengren (Leptoneura) ...	71
<i>Chlorimene</i> , Hübner (Euptychia) ..	33	<i>Coctei</i> , <i>Guérin</i> (Satyrus)	68
<i>Chloris</i> , <i>Cramer</i> (Papilio)	33	<i>Coctei</i> , Butler (Epinephele)	68
<i>Chloris</i> , Westwood (Euptychia) ...	33	Westwood (Erebia)	68
<i>Chorinæus</i> , <i>Fabricius</i> (Papilio) ...	1	<i>Cocytia</i> , <i>Felder</i> (Pronophila)	178
<i>Chorinæus</i> , Duncan (Arpidea) ...	1	<i>Cocytia</i> , Butler (Pedaliodes)	178
Westwood (Cœrois)	1	<i>coelestis</i> , Butler (Euptychia)	29
<i>Chrysogone</i> , <i>Hewitson</i> (Taygetis) ..	10	<i>CÆLITES</i> , Westwood	111
<i>Chryxus</i> , <i>Hewitson</i> (Chionobas) ...	161	Constantia, <i>Cramer</i> (Papilio)	5
<i>Chryxus</i> , Butler (Eneis)	161	Constantia, Boisduval (Cyllo) ...	5
<i>cineros</i> , Butler (Mycæsis)	138	Godart (Satyrus)	5
<i>Circe</i> , E. Doubleday (Hipparchia) ...	50	Hübner (Hipo)	5
Fabricius (Papilio)	50	<i>CORADES</i> , E. Doubleday	185
Felder (Lasiophila)	181	<i>Cordace</i> , Hübner (Tisiphone)	166
Godart (Satyrus)	50	<i>Cordace</i> , Butler (Geitoneura ?) ...	166
<i>Cirta</i> , <i>Felder</i> (Lasiophila)	181	Westwood (Lasiommata) ...	166
<i>Cirta</i> , <i>Hewitson</i> (Pronophila) ...	181	<i>Cordillera</i> , Westwood (Pronophila) ..	185
<i>Cissa</i> , Hübner (Cithærias)	101	<i>Cordula</i> , <i>Fabricius</i> (Papilio)	60
<i>Cissia</i> , <i>Cramer</i> (Papilio)	14		
E. Doubleday (Euptychia) ...	14		
<i>Cistene</i> , <i>Hewitson</i> (Corades)	187		
<i>Clarissa</i> , <i>Cramer</i> (Papilio) ...	16, 190		

	Page		Page
<i>Cordula</i> , E. Doubleday (Hipparchia)	60	<i>Davus</i> , Godart (Satyrus)	46
Godart (Satyrus)	60	Ochsenheimer (Papilio)	47
<i>Corinna</i> , Hübner (Papilio)	43	Stephens (Hipparchia)	46
<i>Corinna</i> , Boisduval (Satyrus) ...	43	<i>decorata</i> , Felder (Pronophila) ...	175
E. Doubleday (Cænonympha) ...	43	<i>decorata</i> , Butler (Pedaliodes) ...	175
Freyer (Pap. Hipparchia) ...	43	<i>Deianira</i> , Hewitson (Mycælesis) ...	142
Godart (Satyrus)	43	<i>Deidamia</i> , Eversmann (Hipparchia)	123
Ochsenheimer (Papilio)	44	<i>Deidamia</i> , Butler (Pararge)	123
<i>Corinus</i> , Westwood (Papilio)	44	Westwood (Lasiommata) ...	123
<i>Corycia</i> , Hewitson (Cænura)	147	<i>Dejanira</i> , Esper (Pap. Nymph. Gemm.)	122
<i>cosmophila</i> , Hübner (Neonympha) ...	36	Eversmann (Hipparchia) ...	123
<i>cosmophila</i> , Butler (Euptychia) ...	36	Godart (Satyrus)	123
<i>COSMOSATYRUS</i> , Felder	59	Hübner (Enodia)	122
<i>Crambis</i> , Freyer (Pap. Hipparchia)	163	Linnæus (Papilio)	122
<i>Crambis</i> , Butler (Cneis)	163	Westwood (Lasiommata) ...	123
Westwood (Chionobas)	163	<i>dejecta</i> , Bates (Pronophila) ...	173
<i>Cramer</i> , Butler (Cyllo)	5	<i>Demophile</i> , Freyer (Pap. Hipparchia)	47
<i>Cramer</i> , Butler (Hippio)	5	<i>dendrophilus</i> , Trimen (Satyrus) ...	113
<i>Crantor</i> , Fabricius (Papilio)	18	<i>dendrophilus</i> , Butler (Neope) ...	113
<i>Crantor</i> , Godart (Satyrus)	18	Trimen (Debis)	113
Westwood (Euptychia)	19	<i>Dexamenus</i> , Hewitson (Mycælesis) ...	142
<i>Criahna</i> , Westwood (Cyllo)	111	<i>Diana</i> , Butler (Debis)	117
<i>Criahna</i> , Butler (Neorina)	111	<i>Diana</i> , Butler (Lethe)	117
<i>Crisia</i> , Hübner (Euptychia)	158	* <i>diaphana</i> , Lucas (Hætera)	102
<i>Crisia</i> , Westwood (Neon. Ragadia)	158	<i>Dinarbas</i> , Hewitson (Debis)	117
<i>Crisilda</i> , Hewitson (Ragadia)	158	<i>Dinarbas</i> , Butler (Lethe)	117
<i>Cyamites</i> , Boisduval (Satyrus) ...	142	<i>Dindymene</i> , Cramer (Papilio)	104
<i>Cyamites</i> , Butler (Mycælesis)	142	Godart (Satyrus)	104
Westwood (Cænonympha)	142	Westwood (Hætera)	104
<i>Cybele</i> , Butler (Corades)	186	<i>Dinias</i> , Hewitson (Dædalma)	183
<i>Cyclopius</i> , Herrich-Schäffer (Erebia)	92	<i>Diniche</i> , Hewitson (Mycælesis) ...	144
<i>Cyclopius</i> , Eversmann (Hipparchia)	92	<i>Dinon</i> , Hewitson (Mycælesis) ...	142
Westwood (Satyrus)	92	<i>Dioxippe</i> , Hübner (Papilio)	88
<i>CYLOGENES</i> , Butler	6	<i>Dioxippe</i> , Duponchel (Satyrus) ...	88
<i>Cymela</i> , Cramer (Papilio)	17	<i>Disa</i> , Thunberg (Papilio)	88
<i>Cymelia</i> , Hübner (Megisto)	17	<i>Disa</i> , Freyer (Pap. Hipparchia) ...	88
<i>Cyrillus</i> , Herbst (Papilio)	61	Westwood (Erebia)	88
		Zetterstedt (Satyrus)	88
D.		<i>discoidalis</i> , Kirby (Hipparchia) ...	92
<i>Dædale</i> , Bergsträsser (Papilio) ...	54	<i>discoidalis</i> , Westwood (Erebia) ...	93
<i>DÆDALMA</i> , Hewitson	183	<i>divergens</i> , Butler (Euptychia) ...	17
* <i>Daidis</i> , Hewitson (Mycælesis) ...	145	<i>Dohnii</i> , Zeller (Triphysa)	49
<i>Dalmata</i> , Godart (Satyrus)	91	<i>Dora</i> , Hewitson (Mycælesis)	142
<i>Damaris</i> , G. R. Gray (Orinoma) ...	110	<i>Doræte</i> , Hewitson (Dædalma) ...	183
<i>Danaë</i> , Hufnagel (Papilio)	56	<i>Dorilis</i> , Borkhausen (Papilio) ...	44
<i>D'Arceti</i> , Lefebvre (Satyrus)	154	<i>Dorinda</i> , Felder (Dædalma) ...	183
<i>Darena</i> , Felder (Debis)	115	<i>Dorion</i> , Hübner (Papilio)	44
<i>Darena</i> , Butler (Lethe)	115	* <i>Doris</i> , Cramer (Papilio)	28, 139
<i>Daretis</i> , Hewitson (Debis)	115	Westwood (Euptychia)	28
<i>Daretis</i> , Butler (Lethe)	115	<i>Dorothea</i> , Cramer (Papilio)	132
<i>Davendra</i> , Moore (Epinephele) ...	65	<i>Dorothea</i> , Hübner (Neonympha) ...	132
<i>Davus</i> , Fabricius (Papilio)	46	<i>Dorus</i> , Esper (Pap. pleb.-rural) ...	44
<i>Davus</i> , Dalman (Erebia)	47	<i>Dorus</i> , E. Doubleday (Cænonympha)	44
E. Doubleday (Cænonympha) ...	46	Godart (Satyrus)	44
		Herbst (Papilio)	44
		<i>Dorycus</i> , Boisduval (Satyrus)	141

	Page
<i>Doryceus</i> , Hewitson (<i>Mycalesis</i>)	141
Westwood (<i>Cænonympha</i>)	141
<i>Doxes</i> , Godart (<i>Satyrus</i>)	35
<i>Doxes</i> , Westwood (<i>Euptychia</i>)	35, 36
<i>Dracontis</i> , Hübner (<i>Pieris</i>)	105
<i>Dracontis</i> , Butler (<i>Pierella</i>)	105
Westwood (<i>Hætera</i>)	105
* <i>Dromus</i> , Fabricius (<i>Papilio</i>)	85
Godart (<i>Satyrus</i>)	86
Westwood (<i>Erebia</i>)	86
<i>Drusia</i> , Cramer (<i>Papilio</i>)	133
<i>Drusilla</i> , Hewitson (<i>Dædalma</i>)	183
<i>Dryas</i> , Felder (<i>Epinephele</i>)	67
Scopoli (<i>Papilio</i>)	61
<i>Dryas</i> , Butler (<i>Satyrus</i>)	61
Esper (<i>Pap. Nymph. Gemm.</i>)	61
<i>Drymæa</i> , Hewitson (<i>Dædalma</i>)	184
<i>Drypetis</i> , Hewitson (<i>Debis</i>)	114
<i>Drypetis</i> , Butler (<i>Lethe</i>)	114
<i>Duboulayi</i> , Butler (<i>Hipparchio-</i> <i>ides</i>)	100
<i>Duboulayi</i> , Butler (<i>Heteronympha</i>)	100
<i>Duponchelii</i> , Guérin (<i>Satyrus</i>)	141
* <i>Duryodana</i> , Felder (<i>Cylo</i>)	2
<i>Duryodana</i> , Butler (<i>Melanitis</i>)	2
<i>Dynsate</i> , Hewitson (<i>Debis</i>)	115
<i>Dynsate</i> , Butler (<i>Lethe</i>)	115
<i>Dyrta</i> , Felder (<i>Debis</i>)	115
<i>Dyrta</i> , Butler (<i>Lethe</i>)	115

E.

<i>Ebusa</i> , Cramer (<i>Papilio</i>)	32
<i>Ebusa</i> , Hübner (<i>Euptychia</i>)	29, 32
<i>Echo</i> , Cramer (<i>Papilio</i>)	10
<i>Echo</i> , Westwood (<i>Taygetis</i>)	11
* <i>Edda</i> , Ménétériés (<i>Erebia</i>)	89
<i>Egea</i> , Borkhausen (<i>Papilio</i>)	73
<i>Egeria</i> , Hübner (<i>Pararge</i>)	120
<i>Electra</i> , Butler (<i>Euptychia</i>)	19
<i>elegans</i> , Butler (<i>Erites</i>)	147
<i>Eliasis</i> , Hewitson (<i>Mycalesis</i>)	132
<i>ELINA</i> , Blanchard	64
<i>Elionas</i> , Hewitson (<i>Mycalesis</i>)	131
<i>Embla</i> , Thunberg (<i>Papilio</i>)	88
<i>Embla</i> , Boisduval (<i>Erebia</i>)	89
Dalman (<i>Erebia</i>)	88
Duponchel (<i>Satyrus</i>)	89
Freyer (<i>Pap. Hipparchia</i>)	88
Ochsenheimer (<i>Papilio</i>)	89
Zetterstedt (<i>Satyrus</i>)	88
<i>embolima</i> , Butler (<i>Debis</i>)	115
<i>embolima</i> , Butler (<i>Lethe</i>)	115
<i>Emilia</i> , Butler (<i>Dædalma</i>)	183
<i>empusa</i> , Felder (<i>Pronophila</i>)	173
<i>empusa</i> , Butler (<i>Pedaliodes</i>)	173
<i>encycla</i> , Felder (<i>Pronophila</i>)	175
<i>encycla</i> , Butler (<i>Pedaliodes</i>)	175
<i>Enyo</i> , Butler (<i>Euptychia</i>)	27
Hewitson (<i>Corades</i>)	185

<i>Euus</i> , Butler (<i>Euptychia</i>)	24
<i>Epimenides</i> , Ménétériés (<i>Lasiom-</i> <i>mata</i>)	122
<i>Epimenides</i> , Butler (<i>Pararge</i>)	122
<i>Epiminthia</i> , Westwood (<i>Celites</i>)	111
<i>EPINEPHELE</i> , Hübner	64
<i>Epinephele</i> , Felder (<i>Neonympha</i>)	24
<i>Epinephele</i> , Butler (<i>Euptychia</i>)	24
<i>Epiphron</i> , Knoch (<i>Papilio</i>)	72
<i>Epiphron</i> , Godart (<i>Satyrus</i>)	72
Herrich-Schäffer (<i>Erebia</i>)	72
<i>Episodes</i> , Butler (<i>Erebia</i>)	80
<i>Epistigne</i> , Duponchel (<i>Satyrus</i>)	84
<i>Epistygne</i> , Hübner (<i>Papilio</i>)	84
<i>Epistygne</i> , Boisduval (<i>Erebia</i>)	84
<i>EREBIA</i> , Dalman	72
<i>erebioides</i> , Felder (<i>Idioneura</i>)	99
* <i>Ereiba</i> , Felder (<i>Pronophila</i>)	174
<i>Erichsonia</i> , Felder (<i>Cylo</i>)	4
<i>Erichsonia</i> , Butler (<i>Melanitis</i>)	4
<i>Erigone</i> , Butler (<i>Euptychia</i>)	18
<i>Erina</i> , Fabricius (<i>Papilio</i>)	76
<i>Eriphile</i> , Westwood (<i>Erebia</i>)	75
<i>Eriphule</i> , Butler (<i>Euptychia</i>)	19
* <i>Eriphyle</i> , Freyer (<i>Pap. Hip-</i> <i>parchia</i>)	75
<i>ERITES</i> , Westwood	146
<i>Ero</i> , Bremer (<i>Erebia</i>)	88
<i>erubescens</i> , Butler (<i>Taygetis</i>)	10
<i>erycina</i> , Butler (<i>Euptychia</i>)	36
<i>Erynia</i> , Esper (<i>Pap. Nymph.</i> <i>Gemm.</i>)	77, 85
* <i>Erythia</i> , Hübner (<i>Papilio</i>)	54
<i>Esmeralda</i> , E. Doubleday (<i>Hætera</i>)	101
<i>Esmeralda</i> , Butler (<i>Callitera</i>)	101
<i>ETREONA</i> , Westwood	168
<i>Ethus</i> , Fabricius (<i>Papilio</i>)	88
<i>Eudora</i> , Esper (<i>Pap. Nymph.</i> <i>Gemm.</i>)	64
<i>Eudora</i> , Butler (<i>Epinephele</i>)	64
Fabricius (<i>Papilio</i>)	64
Godart (<i>Satyrus</i>)	64
<i>Eudorus</i> , Herbst (<i>Papilio</i>)	64
<i>Eugenia</i> , Felder (<i>Neonympha</i>)	26
* <i>Eumenis</i> , Freyer (<i>Pap. Hip-</i> <i>parchia</i>)	90
* <i>Eumonia</i> , Ménétériés (<i>Erebia</i>)	83
<i>Euphemia</i> , Hewitson (<i>Cæno-</i> <i>nympha</i>)	168
<i>Euphemia</i> , Butler (<i>Hypocysta</i>)	168
<i>Eupolia</i> , Hewitson (<i>Eteona</i>)	168
<i>EUPTYCHIA</i> , Hübner	14, 32
<i>Euptychidia</i> , Butler (<i>Taygetis</i>)	13
<i>euptychioides</i> , Felder (<i>Celites</i>)	112
<i>Europa</i> , Fabricius (<i>Papilio</i>)	114
<i>Europa</i> , Hübner (<i>Oreas & Lethe</i>)	114
Westwood (<i>Debis</i>)	144
<i>Euryale</i> , Esper (<i>Pap. Nymph.</i> <i>Gemm.</i>)	82
<i>Euryale</i> , Boisduval (<i>Erebia</i>)	82

	Page		Page
<i>Euryale</i> , Freyer (Pap. Hipparchia)	82	<i>fumata</i> , Butler (Euptychia)	38
Godart (Satyrus)	82	<i>funebria</i> , Godart (Satyrus)	128
Herrich-Schäffer (Erebia) 74, 82		<i>funebria</i> , Westwood (Mycalesis)	128
Hübner (Papilio)	82	<i>furina</i> , Hewitson (Euptychia) ...	35
<i>Eurydia</i> , Linnæus (Papilio)	123	<i>fuscum</i> , Felder (Dasyomma)	144
<i>Eurythris</i> , Godart (Satyrus)	17	<i>fuscum</i> , Butler (Mycalesis)	144
<i>Eurytris</i> , Fabricius (Papilio)	17		
Westwood (Neonympha)	17	G.	
<i>Eurytus</i> , Fabricius (Papilio)	17	<i>galactinus</i> , Boisduval (Satyrus) ...	48
<i>Eurytus</i> , Butler (Euptychia)	17	<i>galactinus</i> , But'er (Cænonympha) ..	48
<i>Eusirus</i> , Hopffer (Mycalesis)	128	<i>Galatea</i> , Ochsenheimer (Papilio)	153
<i>Evadne</i> , Cramer (Papilio)	128	Schrank (Maniola)	153
<i>Evadne</i> , Hübner (Mycalesis)	128	<i>Galathea</i> , Linnæus (Papilio)	153
<i>Evenus</i> , Hopffer (Mycalesis)	129	<i>Galathea</i> , Boisduval (Arge)	153
<i>Evenus</i> , Trimen (Mycalesis)	128	Esper (Pap. Nymph. Gemm.)	153
<i>Eversmanni</i> , Eversmann (Hip-		Godart (Satyrus)	153
parchia)	127	Stephens (Hipparchia)	153
<i>Eversmanni</i> , Butler (Amecera) ...	127	<i>Galazera</i> , Esper (Pap. Nymph.	
Westwood (Lasiommatata) ...	127	Gemm.)	153
* <i>Evias</i> , Lefebvre (Erebia)	83	Galesus, Godart (Satyrus)	22
<i>excavata</i> , Butler (Taygetis)	8	<i>Galesus</i> , Butler (Euptychia)	22
		Westwood (Neonympha)	22
F.		<i>Gamaliba</i> , Butler (Mycalesis)	139
<i>Fagi</i> , Scopoli (Papilio)	50	<i>Gambius</i> , E. Doubleday (Myc-	
<i>Fagi</i> , Butler (Hipparchia)	50	lesis)	129
<i>fallax</i> , Felder (Neonympha)	23	<i>Gaschkevitchii</i> , Felder (Lasiom-	
<i>fallax</i> , Butler (Euptychia)	23	matata)	113
<i>fasciata</i> , Butler (Erebia)	92	GEITONEURA, Butler	166
Hewitson (Yphthima)	151	<i>Gelanor</i> , Godart (Satyrus)	100
<i>fasciata</i> , Butler (Yphthima)	151	<i>gemma</i> , Hübner (Neonympha) ...	37
<i>fatua</i> , Freyer (Pap. Hipparchia) ..	57	<i>gemma</i> , Boisduval (Satyrus)	37
<i>fatua</i> , Herrich-Schäffer (Satyrus) ..	57	Butler (Euptychia)	37
Hübner (Taygetis)	11	<i>gemma</i> , Butler (Euptychia) ...	35
<i>Fauna</i> , E. Doubleday (Hipparchia) ..	56	<i>George</i> , Freyer (Pap. Hipparchia) ..	85
Esper (Pap. Nymph. Gemm.)	56	Gera, Hewitson (Euptychia)	35
Godart (Satyrus)	56	<i>Gerdrutius</i> , Fabricius (Papilio) ...	4
Sulzer (Papilio)	56	<i>Gerdrutius</i> , Butler (Melanitis) ...	4
FAUNULA, Felder	95	Westwood (Cyllo?)	5
Felderi, Butler (Mycalesis)	144	<i>Germainii</i> , Felder (Tetraphlebia) ..	95
<i>ferruginosa</i> , Butler (Lymanopoda)		Geryon, Felder (Antirrhæa)	108
.....	169	<i>Gesion</i> , Esper (Pap. Nymph.	
<i>Ferula</i> , Fabricius (Papilio)	60	Gemm.)	88
<i>Fidia</i> , Linnæus (Papilio)	57	Quensel (Papilio)	88
<i>Fidia</i> , E. Doubleday (Hipparchia) ..	58	<i>Geticus</i> , Esper (Papilio)	39
Esper (Pap. Nymph. Gemm.)	57, 58	<i>Getulia</i> , Felder (Mycalesis)	141
Godart (Satyrus)	58	<i>Geyeri</i> , Herrich-Schäffer (Satyrus) ..	53
Herbst (Papilio)	57	<i>Geyeri</i> , Butler (Hipparchia)	53
<i>Flora</i> , Philippi (Satyrus)	58	<i>gigas</i> , Butler (Euptychia)	30
<i>Flora</i> , Butler (Hipparchia?)	58	Butler (Eneis)	161
<i>Fortunatus</i> , Fabricius (Papilio) ...	164	<i>Girondius</i> , Godart (Satyrus)	106
Godart (Satyrus)	164	<i>glacialis</i> , Esper (Pap. Nymph.	
* <i>Francisca</i> , Cramer (Papilio)	132	Gemm.)	77
<i>Franciscus</i> , Herbst (Papilio)	132	<i>glacialis</i> , Butler (Oreina)	77
<i>fraterna</i> , Butler (Mycalesis)	145	<i>glaucina</i> , Bates (Euptychia)	28
* <i>Fulla</i> , Eversmann (Hip. Chiono-		<i>Glaucopis</i> , Felder (Epinephele) ...	61
bas)	162	<i>Glaucopis</i> , Butler (Satyrus)	61
<i>Fulla</i> , Herrich-Schäffer (Chiono-		<i>Glycerion</i> , Borkhausen (Papilio) ..	40
bas)	162	GNOPHODES, Westwood	6
<i>fulvescens</i> , Felder (Cyllo)	3	<i>Gnophodes</i> , Butler (Melanitis) ...	5
		<i>Goalpara</i> , Moore (Zophocæsa)	109

	Page
Goante, <i>Esper</i> (Pap. Nymph. Gemm.)	84
Goante, Boisduval (Erebia)	84
Godart (Satyrus)	84
Herrich-Schäffer (Erebia) ...	85
Ochsenheimer (Papilio)	84
Gokala, <i>Moore</i> (Melanitis)	4
Gopa, <i>Felder</i> (Mycalesis)	140
Gorge, <i>Esper</i> (Pap. Nymph. Gemm.)	85
Gorge, Boisduval (Erebia)	85
Freyer (Pap. Hipparchia) ...	85
Godart (Satyrus)	78
Hübner (Papilio)	85
Treitschke (Hipparchia)	84
Gorgone, Boisduval (Erebia)	84
Gorgone, Herrich-Schäffer (Erebia)	84, 85
Gorgophone, <i>Bellier</i> (Erebia)	85
Goschkevitchii, <i>Ménétriés</i> (Lasiommata)	113
Goschkevitchii, <i>Butler</i> (Neope) ...	113
Gotama, <i>F. Moore</i> (Mycalesis) ...	134
granulata, <i>Butler</i> (Pedaliodes) ...	173
Griela, <i>Fabricius</i> (Papilio)	88
Godart (Satyrus)	88
Grimon, <i>Godart</i> (Satyrus)	21
Grimon, <i>Butler</i> (Euptychia)	21
E. Doubleday (Neonympha)	21
Griphe, <i>Felder</i> (Neonympha)	27
GYROCHEILUS, <i>Butler</i>	179

H.

Hæmonia, <i>Hewitson</i> (Hypocista) ..	167
HÆTERA, <i>Fabricius</i>	102
*Halimede, <i>Schrenck</i> (Arge)	154
Hanibal, <i>Herbst</i> (Papilio)	44
Harmonia, <i>Butler</i> (Euptychia) ...	24
Harpalyce, <i>Butler</i> (Hetera)	101
Harpalyce, <i>Butler</i> (Callitara)	101
Harpypia, <i>Felder</i> (Neonympha) ...	41
Hebe, <i>Trimen</i> (Ypthima)	147
Hebe, <i>Hewitson</i> (Cænura)	147
Hecæerge, <i>Hewitson</i> (Hypocista) ..	167
Heidenreichii, <i>Lederer</i> (Satyrus) ..	54
Heidenreichii, <i>Butler</i> (Hipparchia) ..	54
Hela, <i>Felder</i> (Antirrhæa)	106
Hela, <i>Butler</i> (Anchiphlebia)	106
Helena, <i>Westwood</i> (Cyllo)	3
Helicta, <i>Hübner</i> (Neonympha) ...	37
Hübner (Oreas)	37
Helle, <i>Cramer</i> (Papilio)	15
Helvina, <i>Hewitson</i> (Hætera)	105
helvina, <i>Butler</i> (Pierella)	105
Hercyna, <i>Hübner</i> (Tisiphone) ...	71
Heri, <i>F. Moore</i> (Mycalesis)	134
Hermes, <i>Fabricius</i> (Papilio)	23
Hermes, <i>Butler</i> (Euptychia)	23
E. Doubleday (Neonympha)	23

Hermes, <i>Godart</i> (Satyrus)	23, 30
Herbst (Papilio)	30
Hermessa, <i>Hübner</i> (Euptychia) ...	30
Hermione, Boisduval (Satyrus) ...	51
Godart (Satyrus)	50
Herbst (Papilio)	51
Linnæus (Papilio)	50
Hermione major, <i>Esper</i> (Pap. Nymph. Gemm.)	50
Hermione minor, <i>Esper</i> (Pap. Nymph. Gemm.)	51
Hero, <i>Linnæus</i> (Papilio)	40
Hero, <i>De Geer</i> (Papilio)	46
E. Doubleday (Cænonympha)	40
Esper (Pap. pleb. rural.)	40
Godart (Satyrus)	40
Schrank (Maniola)	40
Stephens (Hipparchia)	40
Herophile, <i>Hübner</i> (Calisto)	97
Herse, <i>Cramer</i> (Papilio)	33
Herse, <i>Borkhausen</i> (Papilio)	85
Denis (Papilio)	69
Hübner (Euptychia)	33
Herseis, <i>Godart</i> (Satyrus)	33
Westwood (Euptychia)	33
*Herta, <i>Herbst</i> (Papilio)	155
Boisduval (Arge)	155
Hertha caucasica, <i>Nordmann</i> (Arge)	156
Hesione, <i>Cramer</i> (Papilio)	138, vi
Hesione, <i>Godart</i> (Satyrus)	138
Hübner (Mycalesis)	138
Wallengren (Orsotriæna) ...	139
hestioides, <i>Felder</i> (Zethera)	110
HETERONYMPHA, <i>Wallengren</i>	99
Hewitsonia, <i>Butler</i> (Lasiophila) ...	182
Hewitsonii, <i>Butler</i> (Euptychia) ...	34
Doumet (Idiomorphus)	127
Lederer (Erebia)	84
hiemalis, <i>Butler</i> (Euptychia)	35
Hiera, <i>Hübner</i> (Papilio)	126
Hiera, Boisduval (Satyrus)	126
Butler (Amecera)	127
Westwood (Lasiommata) ...	127
hilara, <i>Bates</i> (Pronophila)	180
Felder (Neonympha)	27
hilara, <i>Butler</i> (Euptychia)	27
Butler (Oxeoschistus)	180
Hilda, <i>Hewitson</i> (Neorina)	111
Hilda, <i>Schneider</i> (Papilio)	162
Himachala, <i>Moore</i> (Theope)	110
Himachala, <i>Butler</i> (Anadobis)	111
Moore (Ethope)	110
HIPIO, <i>Hübner</i>	5
HIPPARCHIA, <i>Fabricius</i>	50
Hippia, <i>Cramer</i> (Papilio)	93
Hippia, E. Doubleday (Erebia?) ...	93
Wallengren (Pseudonympha) ..	93
Westwood (Erebia)	93
Hippius, <i>Herbst</i> (Papilio)	93

	Page		Page
<i>Hippodice</i> , Hübner (Papilio)	60	<i>Iocius</i> , Hewitson (Idiomorphus)...	127
<i>Hippolite</i> , Freyer (Pap. Hipparchia)	52	<i>Ichthya</i> , Herrich-Schäffer (Co-	
Herbst (Papilio)	52	rades)	185
<i>Hippolyte</i> , Esper (Pap. Nymph. Gemm.)	52	<i>Ida</i> , Esper (Pap. Nymph. Gemm.)	69
<i>Hippolyte</i> , Boisduval (Satyrus)	52	<i>Ida</i> , Borkhausen (Papilio)	69
E. Doubleday (Hipparchia)	52	Butler (Epinephele)	69
Zetterstedt (Papilio)	52	Godart (Satyrus)	69
<i>Hippolytus</i> , Herbst (Papilio)	60	IDIOMORPHUS, Doumet	127
<i>Hippomedusa</i> , Boisduval (Erebia)	90	IDIONEURA, Felder	99
<i>Hispania</i> , Butler (Erebia)	86	<i>Iduna</i> , Hewitson (Corades).....	185
<i>Hobartia</i> , Westwood (Lasiommata)	165	<i>Igoleta</i> , Felder (Mycalasis)	133
<i>Hobartia</i> , Butler (Argynnis).....	165	<i>incerta</i> , Hewitson (Amechania) ...	110
Homconympha, Felder	95	<i>incerta</i> , Butler (Zethers).....	110
<i>Hortensia</i> , Hewitson (Hætera) ...	106	<i>inconspicua</i> , Butler (Dædalma)...	183
<i>Hortensia</i> , Butler (Pierella)	106	<i>Ines</i> , Hoffmannsegg (Papilio).....	156
<i>Hortona</i> , Hewitson (Hætera)	106	<i>Ines</i> , Boisduval (Arge)	156
<i>Hortona</i> , Butler (Pierella)	106	Duponchel (Satyrus)	156
<i>Hübneri</i> , Butler (Euptychia)	22	<i>infuscata</i> , M. Leay (Cænonympha)	188
Felder (Satyrus)	52	<i>Inica</i> , Hewitson (Ypthima)	151
<i>Hübneri</i> , Butler (Hipparchia) ...	52	<i>Inica</i> , Butler (Ypthima)	151
<i>humilia</i> , Felder (Stygnus)	96	<i>injusta</i> , Wallengren (Mycalasis)	129
<i>humilis</i> , Butler (Cœlites)	112	<i>inornata</i> , Edwards (Cænonympha)	47
<i>Hyagriva</i> , Moore (Ypthima)	152	Felder (Taygetis).....	13
<i>Hyceta</i> , Hewitson (Hætera)	103	<i>insignis</i> , Butler (Euptychia)	38
<i>Hyceta</i> , Butler (Pierella).....	103	<i>Iolaus</i> , Bonelli (Pap. Satyrus).....	51
<i>Hygea</i> , Hewitson (Hypocista).....	167	<i>Ionius</i> , Westwood (Lymanopoda)	169
<i>Hylata</i> , Ménétriés (Hipparchia)...	155	<i>Iphia</i> , Eversmann (Hipp. Cænonympha)	48
<i>Hylata</i> , Butler (Arge)	155	<i>Iphigenus</i> , Herbst (Papilio)	39
<i>Hymenæa</i> , Felder (Hætera)	102	<i>Iphia</i> , Borkhausen (Papilio).....	47
<i>Hypæsia</i> , Hewitson (Hætera)	103	Butler (Cænonympha)	47
<i>Hyperanthe</i> , Hübner (Enodia)	63	Denis (Papilio)	40
<i>Hyperanthus</i> , Fuessly (Papilio) ...	63	E. Doubleday (Cænonympha)	40
Godart (Satyrus)	63	Eversmann (Hipparchia) ...	40
Stephens (Hipparchia)	63	Freyer (Pap. Hipparchia) ...	41
<i>Hyperantus</i> , Linnaeus (Papilio)...	63	Godart (Satyrus)	40
<i>Hyperantus</i> , Butler (Satyrus).....	63	Stephens (Hipparchia)	47
<i>hyperbioides</i> , Wallengren (Pseudonympha)	94	<i>Iris</i> , Felder (Neonympha)	34
<i>Hyperbius</i> , Linnaeus (Papilio) ...	93	<i>Iris</i> , Butler (Euptychia)	34
<i>Hyperbius</i> , Cramer (Papilio)	94	<i>Irius</i> , Fabricius (Papilio)	167
Godart (Satyrus).....	93	<i>Irius</i> , Butler (Hypocysta)	167
Wallengren (Pseudonympha)	93	Godart (Satyrus).....	167
Westwood (Erebia).....	93	Westwood (Cænonympha).....	167
HYPOCYSTA, Westwood.....	167	<i>Irmina</i> , Hewitson (Pronophila)...	181
<i>Hyrania</i> , Kollar (Satyrus)	116	<i>Irmina</i> , Butler (Oxeoschistus) ...	181
<i>Hyrania</i> , Butler (Lethe)	116	<i>Isabella</i> , Butler (Debis)	119
Westwood (Debis)	116	<i>Isana</i> , Kollar (Satyrus)	116
<i>Hysius</i> , Godart (Satyrus)	97	Westwood (Debis)	116
<i>Hysius</i> , Westwood (Calisto)	97	<i>Isis</i> , Thunberg (Papilio)	47
I.		<i>Isis</i> , Westwood (Cænonympha)...	47
<i>Ianthe</i> , Hübner (Papilio)	73, 76	Zetterstedt (Satyrus)	47
Schrank (Maniola)	54	<i>Ismene</i> , Cramer (Papilio)	2
Westwood (Erebia).....	76	Butler (Melanitis)	2
<i>Ianthe major</i> , Esper (Papilio).....	54	<i>Ita</i> , Felder (Mycalasis)	145
<i>Iapygia</i> , Esper (Pap. Nymph. Gemm.)	155	<i>Italus</i> , Hewitson (Idiomorphus)...	127
<i>Iapygia</i> , Butler (Arge)	155	<i>Itonia</i> , Hewitson (Ypthima)	150
Cyrillus (Papilio)	155	<i>Itonia</i> , Butler (Ypthima)	150
		<i>Itonis</i> , Hewitson (Euptychia).....	36
		<i>Itys</i> , Felder (Mycalasis)	145
		<i>Ixora</i> , Boisduval (Arge)	167

J.	Page
Janardana, Moore (Mycalesis) ...	136
Janira, Linnæus (Papilio)	66
Janira, Esper (Pap. Nymph. Gemm.)	66
Freyer (Pap. Hipparchia) ...	66
Godart (Satyrus)	66
Stephens (Hipparchia)	66
janirioides, Blanchard (Satyrus) ...	68
*Janirula, Esper (Pap. Nymph. Gemm.)	66
Janirus, Herbst (Papilio)	66
Japonica, Butler (Neope)	113
Joanna, Butler (Enodia)	70
Joanna, Butler (Xenica)	70
Jopas, Hewitson (Mycalesis)	136
Jovita, Felder (Neonympha)	25
Jovita, Butler (Euptychia)	25
Junia, Cramer (Papilio)	30
Junia, Hübner (Euptychia)	30
Junonia, Butler (Euptychia)	35
Butler (Mycalesis)	146
Jurtina, Hübner (Papilio)	64
Linnæus (Papilio)	66
*Justina, Cramer (Papilio)	135
Hübner (Mycalesis)	135
Justinella, Butler (Mycalesis)	135
Jutta, Hübner (Papilio)	160
Jutta, Boisduval (Chionobas) ...	160
Duponchel (Satyrus)	160
Hübner (Ceneis)	160

K.

Kalinda, Moore (Erebia)	78
Kansa, Moore (Debis)	119
Kansa, Butler (Lethe)	119
Kefersteinii, Herrich-Schäffer (Erebia)	74
Klugii, Guérin (Satyrus)	166
Klugii, Butler (Geitoneura)	166

L.

Labda, Hewitson (Lymanopoda) ...	170
Laccine, Felder (Neonympha) ...	27
*Laches, Fabricius (Papilio)	12
Godart (Satyrus)	12
Lachesis, Hübner (Papilio)	154
Lachesis, Boisduval (Arge)	154
Godart (Satyrus)	154
lactea, Hewitson (Lymanopoda) ...	169
latifica, Bates (Pronophila)	180
*Laidion, Borkhausen (Papilio) ...	47
*Lalassis, Hewitson (Mycalesis) ...	137
*Lamia, Sulzer (Pap. Nymph. Gemm.)	104
Felder (Hætera)	104
Lanassa, Felder (Lymanopoda) ...	170
Lappona, Esper (Pap. Nymph. Gemm.)	87

	Page
*Lara, Donovan (Papilio)	150
Larissa, Hübner (Papilio)	155
Larissa, Boisduval (Arge)	155
Freyer (Pap. Hipparchia) ...	155
laroides, Westwood (Xpthima) ...	150
Larus, Felder (Taygetis)	9
Larymna, Hewitson (Hætera)	104
LASIOPHILA, Felder	181
Lasus, Hewitson (Lasiommata) ...	165
Lathoniella, Westwood (Las. Xenica)	165
Lathoniella, Butler (Argynnina) ...	165
Latia, Butler (Euptychia)	36
Latiaria, Hewitson (Debis)	117
Latiaria, Butler (Lethe)	117
Latona, Felder (Hætera)	103
Latona, Butler (Pierella)	103
Lea, Cramer (Papilio)	29
Lea, Godart (Satyrus)	29, 30
Westwood (Euptychia)	29
Leena, Hewitson (Lymanopoda) ...	170
Leander, Esper (Pap. pleb. rural.) ...	42
Leander, E. Doubleday (Cænonympha)	43
Eversmann (Hipparchia) ...	43
Fabricius (Papilio)	42
Freyer (Pap. Hipparchia) ...	42
Godart (Satyrus)	42
Lebba, Hewitson (Lymanopoda) ...	170
Leda, Linnæus (Papilio)	1
Leda, Cramer (Papilio)	2
Fabricius (Melanitis)	2
Godart (Satyrus)	2
Hübner (Oreas and Hipio) ...	1, 2
Westwood (Cyllo)	2
Lefebvrei, Guérin (Satyrus)	64
*Lefebvrei, Boisduval (Erebia) ...	79
Butler (Elina)	64
Duponchel (Satyrus)	79
Herrich-Schäffer (Erebia) ...	78, 79
Lena, Linnæus (Papilio)	105
Lena, Butler (Pierella)	105
Cramer (Papilio)	105
Godart (Satyrus)	105
Hübner (Oreas and Pieris) ...	105
Westwood (Hætera)	105
Leprea, Hewitson (Lasiommata) ...	166
Leprea, Butler (Argynnina)	166
leptoneuroides, Felder (Cosmossatyrus)	59
Lethe, Butler (Euptychia)	18
LETHE, Hübner	114
Leuce, Felder (Acrophthalmia) ...	159
Leucoglene, Felder (Faunula)	95
Leucogyne, Felder (Ptychandra) ...	187
*leucomelas, Esper (Pap. Nymph. Gemm.)	153
Boisduval (Arge)	154
Borkhausen (Papilio)	154
Freyer (Pap. Hipparchia) ...	154

	Page		Page
<i>Läbye, Linnaeus</i> (Papilio)	30	<i>Madura, Westwood</i> (Sat. Erites) ...	147
<i>Libye, Butler</i> (Euptychia)	30	<i>Mapius, Godart</i> (Satyrus)	28
<i>libyoidea, Butler</i> (Euptychia) ...	31	<i>Mapius, Westwood</i> (Euptychia) ...	28
<i>Ligea, Linnaeus</i> (Papilio)	82	<i>Mara, Linnaeus</i> (Papilio)	125
<i>Ligea, Dalman</i> (Erebia)	82	<i>Mara, Borkhausen</i> (Papilio)	124
Esper (Papilio)	90	Butler (Amecera)	125
Freyer (Pap. Hipparchia) ...	82	*Esper (Pap. Nymph. Gemm.)	125
Godart (Satyrus)	82	Godart (Satyrus)	125
Stephens (Hipparchia)	82	Hübner (Papilio)	125
Westwood (Oreina)	83	Schränk (Maniola)	125
<i>Limonias, Philippi</i> (Satyrus)	67	Westwood (Lasiommata) ...	125
<i>Limonias, Butler</i> (Epinephele) ...	67	<i>meroides, Felder</i> (Lasiommata) ...	126
<i>Lindigii, Felder</i> (Antirrhæa)	108	<i>Mæmula, Felder</i> (Lasiommata) ...	126
<i>Liriope, Cyrellus</i> (Papilio)	154	<i>Marula, Butler</i> (Amecera)	126
<i>Lisandra, Cramer</i> (Papilio)	149	<i>Magus, Fabricius</i> (Papilio)	152
<i>Lisandra, Butler</i> (Ypthima)	149	<i>Maianæa, Hewitson</i> (Mycalesis) ...	131
<i>Lisidice, Hübner</i> (Euptychia)	28	<i>Makuta, Horsfield</i> (Hipparchia) ...	158
<i>liturata, Butler</i> (Euptychia)	22	Moore (Ragadia)	158
<i>Lizetta, Cramer</i> (Papilio)	44	<i>Malsara, Hewitson</i> (Mycalesis) ...	138
<i>Lorquini, Felder</i> (Psychandra) ...	187	<i>*Malsara, Moore</i> (Mycalesis) ...	136
<i>Loryma, Hewitson</i> (Ypthima) ...	148	<i>Malsarida, Butler</i> (Mycalesis) ...	134
<i>Lowii, Hewitson</i> (Cyllo)	111	<i>Mamerta, Cramer</i> (Papilio)	136
<i>Lowii, Butler</i> (Neorina)	111	<i>Mamerta, Butler</i> (Mycalesis)	136
<i>Lupita, Reakirt</i> (Neonympha) ...	39	Moore (Mycalesis)	140
<i>Lupita, Butler</i> (Euptychia)	39	<i>Mamurra, Herrich-Schäffer</i> (Sa-	
<i>lusitanica, Petiver</i> (Papilio)	58	tyrus)	53
<i>Luzonia, Felder</i> (Ragadia)	158	<i>Manasses, Felder</i> (Neonympha) ...	26
<i>Lybie, Sulzer</i> (Papilio)	30	<i>Manasses, Butler</i> (Euptychia) ...	26
<i>Lycæon, Müller</i> (Papilio)	64	<i>*Mancinus, Hewitson</i> (Erebia) ...	89
<i>Lydia, Cramer</i> (Papilio)	15	<i>Mandane, Kirby</i> (Cæonympha) .	41
<i>Lydia, Hübner</i> (Euptychia)	15	<i>mandata, Moore</i> (Mycalesis)	139
<i>Lydius, Godart</i> (Satyrus)	15	<i>mandosa, Butler</i> (Mycalesis) ...	139
<i>*Lylus, Esper</i> (Pap. pleb. rural.)	45	<i>*Manipa, Boisduval</i> (Satyrus) ...	145
Freyer (Pap. Hipparchia) ...	45	Felder (Mycalesis)	145
Godart (Satyrus)	45	Westwood (Cæonympha) ...	145
Herrich-Schäffer (Satyrus) ...	46	<i>Manis, Felder</i> (Pronophila)	174
Westwood (Cæonympha) ...	45	<i>Manis, Butler</i> (Pedaliodes)	174
LYMANOPTERA, Westwood	168	<i>Manthara, Felder</i> (Debis)	119
<i>Lysandra, Moore</i> (Ypthima)	149	<i>Manthara, Butler</i> (Lethe)	119
<i>Lysidice, Cramer</i> (Papilio)	28	<i>Manto, Denis</i> (Papilio)	86
Hübner (Oreas)	28	<i>Manto, Dalman</i> (Erebia)	87
Westwood (Euptychia)	28	*Esper (Pap. Nymph. Gemm.)	76
<i>*Lyssa, Hübner</i> (Papilio)	124	Freyer (Pap. Hipparchia) ...	86
Boisduval (Satyrus)	124	Godart (Satyrus)	87
<i>Lyssianassa, Boisduval</i> (Arge) ...	156	Herbst (Papilio)	76
Dahl (Papilio)	156	Schränk (Maniola)	40
M.		<i>mantoides, Butler</i> (Erebia)	87
<i>Maackii, Bremer</i> (Lasiommata) ...	122	<i>marginalis, Butler</i> (Cæonympha) ...	42
<i>Maackii, Butler</i> (Pararge)	122	Motschulsky (Satyrus)	188
<i>Maccabeus, Herbst</i> (Papilio)	76	<i>marginalis, Butler</i> (Lethe?)	188
<i>Machabeus, Westwood</i> (Erebia) ...	76	<i>Marmoræ, Hübner</i> (Papilio)	51
<i>Macleanania, Bates</i> (Hætera) ...	103	<i>Marmoræ, Butler</i> (Hipparchia) ...	51
<i>macrophthalmus, Eversmann</i> (Sa-		<i>marmorata, Butler</i> (Euptychia) ...	21
tyrus)	58	<i>Marpeesa, Hewitson</i> (Debis)	13
<i>Maderakal, Guérin</i> (Satyrus)	126	<i>Marpeesa, Hewitson</i> (Taygetis) ...	13
<i>Maderakal, Butler</i> (Amecera)	126	<i>Martius, Fabricius</i> (Papilio)	140
<i>Madjioosa, Butler</i> (Mycalesis) ...	137	<i>Martius, Godart</i> (Satyrus)	140
<i>Madura, Horsfield</i> (Hipparchia) ...	146	Westwood (Mycalesis)	140
<i>Madura, Moore</i> (Erites)	147	<i>Maurisius, Esper</i> (Pap. Nymph.	
		Gemm.)	76

	Page		Page
<i>Maurus, Esper</i> (Papilio).....	79	<i>Ménétréii</i> , Butler (Pararge).....	122
<i>Maurus</i> , Butler (Erebia).....	79	* <i>Moone</i> , Esper (Pap. Nymph.	
<i>Medea, Denis</i> (Papilio).....	81	Gemm.).....	120
<i>Medea</i> , Eversmann (Hipparchia).....	83	Butler (Lasiommata).....	120
Freyer (Pap. Hipparchia) ...	81	Cramer (Papilio).....	120
Hübner (Papilio).....	81	Wagner (Hipparchia).....	120
Schrank (Maniola).....	90	meridionalis, <i>Felder</i> (Melanargia).....	154
<i>Medeba, Hewitson</i> (Corades).....	186	meridionalis, Butler (Arge).....	154
<i>Medus</i> , Fabricius (Papilio).....	139, 190	<i>Mermeria</i> , Cramer (Papilio).....	7
<i>Medusa, Denis</i> (Papilio).....	90	<i>Mermeria</i> , Godart (Satyrus).....	7
<i>Medusa</i> , Borkhausen (Papilio) ...	81	Hübner (Taygetis).....	7
Dalman (Erebia).....	81	<i>Merope, Fabricius</i> (Papilio).....	99
Eversmann (Hipparchia) ...	90	<i>Merope</i> , Butler (Hipparchioides).....	100
Freyer (Pap. Hipparchia).....	90, 91	E. Doubleday (Lasiommata).....	100
Godart (Satyrus).....	90	Godart (Satyrus).....	100
Westwood (Erebia).....	90	Wallengren (Heteronympha).....	100
<i>Megara, Linnaeus</i> (Papilio).....	123	<i>Messene, Hewitson</i> (Mycalesis) ...	142
<i>Megara</i> , Butler (Amecera).....	124	<i>Mestra, Hewitson</i> (Mycalesis).....	137
Esper (Pap. Nymph.) ...	123, 124	<i>Metagera, Butler</i> (Euptychia) ...	35
Godart (Satyrus).....	123, 124	<i>Methora, Hewitson</i> (Ypthima) ...	149
Hübner (Dira).....	123	<i>Miltiades, Fabricius</i> (Papilio).....	108
Stephens (Hipparchia).....	123	<i>Miltiades</i> , Butler (Charaxes?) ...	108
Strom (Pararge).....	124	Godart (Satyrus).....	108
Westwood (Lasiommata) ...	123	Westwood (Antirrhæa?).....	108
<i>Megamede, Hewitson</i> (Mycalesis).....	136	Westwood (Nymphalis?) ...	108
<i>Megisto, Hübner</i> (section).....	14	<i>Milyas, Hewitson</i> (Mycalesis).....	129
<i>Mehadeva, Boisduval</i> (Satyrus) ...	143	mima, <i>Butler</i> (Euptychia).....	38
<i>Mehadeva</i> , Butler (Mycalesis).....	143	<i>Minerva</i> , Godart (Satyrus).....	118
Westwood (Cænonympha).....	143	<i>Mineus, Linnaeus</i> (Papilio).....	135
<i>Mekara, Moore</i> (Debia).....	119	<i>Mineus</i> , Godart (Satyrus).....	135
<i>Mekara</i> , Butler (Lethe).....	119	Westwood (Mycalesis).....	134, 135
<i>Melampus, Fuessly</i> (Papilio).....	75	<i>MINOIS, Hübner</i> (section).....	61
<i>Melampus</i> , Boisduval (Erebia) ...	75	<i>Minois, Hewitson</i> (Mycalesis).....	144
Butler (Oreina).....	75	<i>Mintha, Hübner</i> (Dira).....	71
Esper (Pap. Nymph. Gemm.).....	73	<i>Mintha</i> , Trimen (Leptoneura) ...	71
Freyer (Pap. Hipparchia).....	75	Westwood (Erebia).....	71
Godart (Satyrus).....	75	* <i>Miriam</i> , Fabricius (Papilio).....	132
Herbst (Papilio).....	73	mirifica, <i>Butler</i> (Lasiommata) ...	100
Newman (Erebia).....	73	mirifica, <i>Butler</i> (Heteronympha).....	100
<i>melancholica</i> , Herrich-Schäffer		Butler (Hipparchioides).....	101
(Erebia).....	84	<i>Miris</i> , Fabricius (Papilio).....	40
<i>MELANITIS, Fabricius</i>	1	<i>Mnasicles, Hewitson</i> (Mycalesis).....	141
* <i>Melas</i> , Herbst (Papilio).....	79	<i>Mnemon</i> , Haworth (Papilio).....	73
Boisduval (Erebia).....	79	* <i>Mnestra</i> , Esper (Papilio).....	75
Duponchel (Satyrus).....	80	Boisduval (Erebia).....	75
Freyer (Pap. Hipparchia) ...	79	Duponchel (Satyrus).....	75
Godart (Satyrus).....	79	Freyer (Pap. Hipparchia) ...	75
Herrich-Schäffer (Erebia) ...	79	Godart (Satyrus).....	75
<i>Melindena, Felder</i> (Ragadia).....	158	Herrich-Schäffer (Erebia) ...	75
<i>Melusina, Herrich-Schäffer</i> (Ere-		Hübner (Papilio).....	75
bia).....	83	modesta, <i>Butler</i> (Euptychia).....	21
<i>Melusina</i> , Fabricius (Papilio).....	132	* <i>mærens</i> , Westwood (Erebia).....	77
Godart (Satyrus).....	132	<i>Molina, Hübner</i> (Euptychia).....	27
Westwood (Mycalesis?).....	132	<i>Mollina, Hübner</i> (Euptychia) ...	27
<i>Menalcas</i> , Poda (Papilio).....	45	<i>Mollina</i> , E. Doubleday (Euptychia).....	27
<i>Menander, Drury</i> (Papilio).....	101	Westwood (Neonympha) ...	27
<i>Menander</i> , Butler (Callitæra).....	101	<i>monachus, Blanchard</i> (Satyrus).....	66
<i>Menava, Moore</i> (Lasiommata) ...	126	<i>monachus</i> , Butler (Epinephele) ...	66
<i>Menava</i> , Butler (Amecera).....	126	<i>Montrolii</i> , Feisthamel (Satyrus).....	64
<i>Ménétréii</i> , Bremer (Satyrus) ...	122	Gay (Elina).....	64

	Page		Page
<i>Montrolis</i> , Westwood (<i>Lasiommata</i>)	64	<i>Nelamus</i> , Boisduval (<i>Erebia</i>)	73
<i>Moorei</i> , Butler (Neope)	113	* <i>Neleus</i> , Freyer (Pap. <i>Hipparchia</i>)	86
Butler (<i>Rhaphicera</i>)	158	* <i>Nelo</i> , Hübner (Papilio)	79
<i>Moorei</i> , Felder (<i>Mycalesis</i>)	140	<i>Neomiris</i> , Godart (<i>Satyrus</i>)	51
<i>Morio</i> , Giorna (Papilio)	77	<i>Neomyris</i> , Boisduval (<i>Satyrus</i>) ..	51
<i>Morna</i> , Fabricius (Papilio)	107, vi	E. Doubleday (<i>Hipparchia</i>)	51
<i>Morna</i> , Butler (<i>Antirrhæa</i>)	107	NEONYMPHA, Hübner (section)	35
Godart (<i>Satyrus</i>)	107	Neonympha, Felder (<i>Taygetis</i>) ..	13
Westwood (<i>Hætera</i>)	107	NEOPE, Butler	112
<i>Morpæa</i> , Butler (<i>Gnophodes</i>) ..	7	Neoridas, Boisduval (<i>Erebia</i>)	81
* <i>Motschulskii</i> , Bremer (<i>Satyrus</i>) ..	148	Neoridas, Borkhausen (Papilio) ..	81
<i>Motschulskii</i> , Hewitson (<i>Yphthima</i>)	148	Duponchel (<i>Satyrus</i>)	81
<i>Mucia</i> , Hewitson (<i>Mycalesis</i>)	142	Treitschke (<i>Hipparchia</i>)	81
<i>Muirheadii</i> , Felder (<i>Lasiommata</i>) ..	113	NEORINA, Westwood	111
<i>Muirheadii</i> , Butler (Neope)	113	NEOSATYRUS, Wallengren	96
<i>Musa</i> , Felder (<i>Zethera</i>)	110	Nephele, Kirby (<i>Hipparchia</i>)	63
<i>Musarion</i> , Borkhausen (Papilio) ..	46	Nephele, Butler (<i>Satyrus</i>)	63
<i>mycalesoides</i> , Felder (<i>Pronophila</i>) ..	176	Hufnagel (Papilio)	45
<i>mycalesoides</i> , Butler (<i>Pedaliodes</i>) ..	176	Westwood (<i>Erebia</i>)	63
<i>Mycaena</i> , Cramer (Papilio)	2	Nereis, Drury (Papilio)	103
<i>Myncea</i> , Cramer (Papilio)	15	Nereis, Butler (<i>Pierella</i>)	103
<i>Myncea</i> , Godart (<i>Satyrus</i>)	15	Godart (<i>Satyrus</i>)	103
Hübner (<i>Euptychia</i>)	15	Hübner (<i>Cithærias</i>)	103
		Westwood (<i>Hætera</i>)	103
N.		* <i>Nerine</i> , Boisduval (<i>Erebia</i>)	78
<i>Nada</i> , Moore (Debis)	117	Duponchel (<i>Erebia</i>)	85
<i>Naidion</i> , Borkhausen (Papilio) ..	42	Freyer (Pap. <i>Hipparchia</i>) ..	78
<i>Nala</i> , Felder (<i>Mycalesis</i>)	140	Herrich-Schäffer (<i>Erebia</i>) ..	78
* <i>Nanna</i> , Ménétries (<i>Chionobas</i>) ..	161	Nevadensis, Felder (<i>Chionobas</i>) ..	161
<i>Napæa</i> , Bates (<i>Pronophila</i>)	179	Nevadensis, Butler (<i>Eneis</i>)	161
<i>Napæa</i> , Butler (<i>Pedaliodes</i>)	179	Nicetas, Hewitson (Debis)	116
<i>Narasingha</i> , Moore (<i>Yphthima</i>) ..	152	Nicetas, Butler (<i>Lethe</i>)	116
<i>Narcissus</i> , Fabricius (Papilio) ..	146	Nicotia, Hewitson (<i>Mycalesis</i>) ..	139
<i>Narcissus</i> , E. Doubleday (<i>Cænonympha</i>)	146	Nirmala, Moore (<i>Erebia</i>)	96
Godart (<i>Satyrus</i>)	146	Nirmala, Butler (<i>Callerebia</i>)	96
<i>Narcysus</i> , Westwood (<i>Mycalesis</i>) ..	146	Norax, Bonelli (Papilio)	44
<i>Nareda</i> , Kollar (<i>Satyrus</i>)	148	Norna, Esper (Pap. Nymph. Gemm.) ..	162
<i>Nareda</i> , Hewitson (<i>Yphthima</i>) ..	148	Norna, Boisduval (<i>Chionobas</i>) ..	162
<i>Narica</i> , Hübner (Papilio)	68	Duponchel (<i>Satyrus</i>)	162
<i>Narica</i> , Boisduval (<i>Satyrus</i>)	68	Godart (<i>Satyrus</i>)	162
Butler (<i>Epinephele</i>)	68	Hübner (<i>Eneis</i>)	162
Freyer (Pap. <i>Hipparchia</i>) ..	68	Hübner (Papilio)	160
<i>Natalii</i> , Boisduval (<i>Satyrus</i>)	94	Thunberg (Papilio)	162, 164
<i>Natalii</i> , Butler (<i>Pseudonympha</i>) ..	94	Nortia, Hewitson (<i>Euptychia</i>)	35
Westwood (<i>Erebia</i>)	94	Nossis, Hewitson (<i>Euptychia</i>)	26
<i>Nautilus</i> , Butler (<i>Mycalesis</i>)	136	Nothia, Hewitson (<i>Colletes</i>)	111
<i>nebulosa</i> , Butler (<i>Euptychia</i>)	26	Numeria, Felder (<i>Neonympha</i>) ..	25
Felder (<i>Mycalesis</i>)	129	Numeria, Butler (<i>Euptychia</i>)	25
<i>Necys</i> , Godart (<i>Satyrus</i>)	31	Numilia, Felder (<i>Neonympha</i>) ..	25
<i>Necys</i> , Westwood (<i>Euptychia</i>) ..	31	Numilia, Butler (<i>Euptychia</i>)	25
<i>Neelgherriensis</i> , Hewitson (Debis) ..	115	Nymphæa, Butler (<i>Taygetis</i>)	9
<i>Neelgherriensis</i> , Guérin (<i>Satyrus</i>) ..	115	nymphosa, Butler (<i>Taygetis</i>)	10
<i>Neelgherriensis</i> , Westwood (Debis) ..	115		
* <i>Negra</i> , Felder (<i>Hætera</i>)	103	O.	
<i>Negrito</i> , Felder (Debis)	119	obscura, Butler (<i>Euptychia</i>)	31
<i>Negrito</i> , Butler (<i>Lethe</i>)	119	Butler (<i>Pronophila</i>)	184
<i>Neilgherriensis</i> , Moore (Debis) ..	115	* <i>obsoleta</i> , Felder (<i>Cyllo</i>)	3
		Butler (<i>Melanitis</i>)	3
		obsoleta, Westwood (<i>Sarromia</i>) ..	168

	Page
<i>ocellifera</i> , <i>Felder</i> (Pronophila) ...	175
<i>ocellifera</i> , <i>Butler</i> (Pedaliodes) ...	175
<i>ochracea</i> , <i>Butler</i> (Euptychia)	21
<i>Edwards</i> (Cænonympha) ...	48
<i>ochrotænia</i> , <i>Felder</i> (Pronophila) ...	178
<i>ochrotænia</i> , <i>Butler</i> (Pedaliodes) ...	178
<i>Ocirrhoë</i> , <i>Fabricius</i> (Papilio)	14
<i>Ocirrhoë</i> , <i>Westwood</i> (Euptychia) ...	14
<i>Ocnus</i> , <i>Butler</i> (Euptychia)	19
<i>Eversmann</i> (Hipparchia) ...	87
<i>Ocnus</i> , <i>Herrich-Schäffer</i> (Erebia) ...	87
<i>ocreata</i> , <i>non descr.</i> (Pierella)	105
<i>oculata</i> , <i>Felder</i> (Pronophila)	176
<i>oculata</i> , <i>Butler</i> (Pedaliodes)	176
<i>Ocyptete</i> , <i>Fabricius</i> (Papilio)	15
<i>Ocyptete</i> , <i>Godart</i> (Satyrus)	15
<i>Hübner</i> (Euptychia)	15
<i>Ocyrrhoë</i> , <i>Godart</i> (Satyrus)	14
<i>Hübner</i> (Euptychia & <i>Oreas</i>) ...	14
<i>Odipe</i> , <i>Hübner</i> (Cænonympha) ...	39
<i>Odippe</i> , <i>Borkhausen</i> (Papilio) ...	39
<i>Odippus</i> , <i>Fabricius</i> (Papilio)	39
<i>Odipus</i> , <i>E. Doubleday</i> (Cænonympha)	39
<i>Eversmann</i> (Hipparchia) ...	39
<i>Godart</i> (Satyrus)	39
<i>Ochsenheimer</i> (Papilio)	39
<i>Oene</i> , <i>Esper</i> (Pap. Nymph. Gemm.) ...	89
<i>Oene</i> , <i>Boisduval</i> (Erebia)	90
<i>Duponchel</i> (Satyrus)	90
<i>Freyer</i> (Pap. Hipparchia) ...	90
<i>Godart</i> (Satyrus)	89
<i>Hübner</i> (Papilio)	89
<i>Westwood</i> (Erebia)	89
<i>Oeneis</i> , <i>Hübner</i>	160
<i>Oeno</i> , <i>Boisduval</i> (Chionobas)	163
<i>Oeno</i> , <i>Butler</i> (Oeneis)	163
<i>Scudder</i> (Chionobas)	163
<i>Enomais</i> , <i>Hübner</i> (Minois and <i>Oreas</i>)	100
<i>Onatas</i> , <i>Hewitson</i> (Mycalesis) ...	140
<i>orbifera</i> , <i>Butler</i> (Lasiophila)	182
<i>Orcus</i> , <i>Latreille</i> (Satyrus)	184
<i>Orcus</i> , <i>Westwood</i> (Pronophila) ...	184
<i>ORESSINOMA</i> , <i>Westwood</i>	153
<i>ORINOMA</i> , <i>Westwood</i>	110
<i>Orinoma</i> , <i>E. Doubleday</i> (Satyrus) ...	110
<i>Oroatis</i> , <i>Hewitson</i> (Mycalesis) ...	138
<i>Orseis</i> , <i>Hewitson</i> (Mycalesis)	136
<i>Ostrea</i> , <i>Westwood</i> (Mycalesis) ...	138
<i>Osyris</i> , <i>Boisduval</i> (Satyrus)	167
<i>Osyris</i> , <i>Felder</i> (Hypocysta)	167
<i>Westwood</i> (Satyrus)	167
<i>Otrea</i> , <i>Cramer</i> (Papilio)	132
<i>Otrea</i> , <i>E. Doubleday</i> (Mycalesis) ...	138
<i>Hübner</i> (Mycalesis)	138
<i>Moore</i> (Mycalesis)	132
<i>Ottomana</i> , <i>Herrich-Schäffer</i> (Erebia)	86
<i>Oxechistis</i> , <i>Butler</i>	180

P.

	Page
<i>Pacarus</i> , <i>Godart</i> (Satyrus)	38
<i>Pacarus</i> , <i>Butler</i> (Euptychia)	38
<i>Westwood</i> (Neonympha) ...	38
<i>Padma</i> , <i>Kollar</i> (Satyrus)	49
<i>Padma</i> , <i>Butler</i> (Aulocera)	49
* <i>Pæas</i> , <i>Esper</i> (Pap. Nymph. Gemm.) ...	61
<i>Hübner</i> (Papilio)	61
<i>Pæon</i> , <i>Godart</i> (Satyrus)	20
<i>Pæon</i> , <i>Westwood</i> (Euptychia) ...	20
<i>Pagyris</i> , <i>Godart</i> (Satyrus)	36
<i>Pagyris</i> , <i>Westwood</i> (Euptychia) ...	36
<i>Palemon</i> , <i>Ernst</i> (Papilio)	44
<i>Pales</i> , <i>Philippi</i> (Epinephele)	67
<i>Palladia</i> , <i>Butler</i> (Euptychia)	16
<i>Pallantis</i> , <i>Hewitson</i> (Pronophila) ...	177
<i>Pallantis</i> , <i>Butler</i> (Pedaliodes) ...	177
<i>pallescens</i> , <i>Butler</i> (Epinephele) ...	65
<i>Pamphila</i> , <i>Hübner</i> (Papilio)	46
<i>Pamphile</i> , <i>Godart</i> (Satyrus)	45
<i>Hübner</i> (Cænonympha)	45
<i>pamphiloides</i> , <i>Reakirt</i> (Cænonympha)	44
<i>Pamphilus</i> , <i>Linnaeus</i> (Papilio) ...	45
<i>Pamphilus</i> , <i>Boisduval</i> (Satyrus) ...	46
<i>E. Doubleday</i> (Cænonympha) ...	45, 46
<i>Esper</i> (Pap. pleb. rural.)	45
<i>Godart</i> (Satyrus)	45
<i>Schrank</i> (Maniola)	45
<i>Stephens</i> (Hipparchia)	45
<i>Panda</i> , <i>Boisduval</i> (Satyrus)	94
<i>Panda</i> , <i>Butler</i> (Pseudonympha?) ...	94
<i>Westwood</i> (Erebia)	94
<i>Pandocus</i> , <i>Moore</i> (Ypthima) ...	149
<i>Pandrosus</i> , <i>Herbst</i> (Papilio)	87
<i>Paneis</i> , <i>Hewitson</i> (Pronophila) ...	174
<i>Paneis</i> , <i>Butler</i> (Pedaliodes)	174
<i>Pannonia</i> , <i>Hewitson</i> (Corades), 185, 186	
<i>Panyasis</i> , <i>Hewitson</i> (Pronophila) ...	179
<i>Panyasis</i> , <i>Butler</i> (Pedaliodes) ...	179
<i>PARAMECERA</i> , <i>Butler</i>	98
<i>Paramegara</i> , <i>Hübner</i> (Papilio) ...	124
<i>Paramegara</i> , <i>Butler</i> (Amecera) ...	124
<i>PARARGE</i> , <i>Hübner</i>	120
<i>Parepa</i> , <i>Hewitson</i> (Pronophila) ...	178
<i>Parepa</i> , <i>Butler</i> (Pedaliodes)	178
<i>Parisatis</i> , <i>Kollar</i> (Satyrus)	58
<i>Parisatis</i> , <i>Butler</i> (Hipparchia) ...	58
<i>Parmenio</i> , <i>Boeber</i> (Papilio)	92
<i>Parmenio</i> , <i>Herrich-Schäffer</i> (Erebia) ...	92
<i>Parmeno</i> , <i>Hewitson</i> (Gnophodes) ...	6
<i>Parneus</i> , <i>Chenu</i> (Gnophodes) ...	6
<i>Parnassia</i> , <i>Felder</i> (Zethera)	109
<i>Pasiphaë</i> , <i>Esper</i> (Pap. Nymph. Gemm.)	69
<i>Pasiphaë</i> , <i>Boisduval</i> (Satyrus) ...	69
<i>Borkhausen</i> (Papilio)	69
<i>Butler</i> (Epinephele)	69
<i>Patnia</i> , <i>Moore</i> (Mycalesis)	146
<i>Patrobas</i> , <i>Hewitson</i> (Pronophila) ...	179

	Page		Page
<i>Patrobas</i> , Butler (<i>Gyrocheilus</i>) ...	179	<i>Pharte</i> , Boisduval (<i>Erebia</i>)	74
<i>Pausia</i> , Hewitson (<i>Pronophila</i>) ...	176	Butler (<i>Oreina</i>)	74
<i>Pausia</i> , Butler (<i>Pedaliodes</i>)	176	Duponchel (<i>Satyrus</i>)	74
<i>Pedacia</i> , Hewitson (<i>Pronophila</i>) ...	176	Freyer (Pap. <i>Hipparchia</i>) ...	74
<i>Pedacia</i> , Butler (<i>Pedaliodes</i>)	176	Godart (<i>Satyrus</i>)	74
<i>PEDALIODES</i> , Butler	172	<i>Phaselis</i> , Hewitson (<i>Pronophila</i>)	184
<i>Pegala</i> , Fabricius (<i>Papilio</i>)	62	<i>Phaselis</i> , Butler (<i>Dædalma</i>)	184
<i>Pegala</i> , Godart (<i>Satyrus</i>)	62	* <i>Phedima</i> , Cramer (<i>Papilio</i>)	3
Westwood (<i>Enodia</i>)	62	Butler (<i>Melanitis</i>)	3
<i>pellucida</i> , Butler (<i>Hetera</i>)	101	<i>Phægea</i> , Borkhausen (<i>Papilio</i>) ...	91
<i>Pelopea</i> , Klug (<i>Hipparchia</i>)	52	* <i>Pherusa</i> , Dahl (<i>Papilio</i>)	157
<i>Pelopea</i> , Westwood (<i>Satyrus</i>) ...	52	Boisduval (<i>Arge</i>)	157
<i>Peloria</i> , Felder (<i>Neonympha</i>)	26	Phidon, Hewitson (<i>Mycalesis</i>) ...	141
<i>Peloria</i> , Butler (<i>Euptychia</i>)	26	Phila, Hewitson (<i>Pronophila</i>)	178
<i>Penelea</i> , Cramer (<i>Papilio</i>)	13	Phila, Butler (<i>Pedaliodes</i>)	178
<i>Penelea</i> , Butler (<i>Taygetis</i>)	13	<i>Philaidia</i> , Borkhausen (<i>Papilio</i>)	43
Godart (<i>Satyrus</i>)	13	<i>Philaretos</i> , Felder (<i>Antirrhæa</i>) ...	108
<i>Penelope</i> , Fabricius (<i>Papilio</i>) ...	16, vi	<i>Philea</i> , E. Doubleday (<i>Cæno-</i>	
Godart (<i>Satyrus</i>)	16	nympha)	42
<i>Perdiccas</i> , Hewitson (<i>Mycalesis</i>) ...	138	Freyer (Pap. <i>Hipparchia</i>) ...	42
<i>perfusca</i> , Butler (<i>Euptychia</i>) ...	18	Hübner (<i>Papilio</i>)	42
<i>Peribæa</i> , Fabricius (<i>Papilio</i>)	38	<i>Philerope</i> , Boisduval (<i>Satyrus</i>) ...	100
<i>Peribæa</i> , Butler (<i>Euptychia</i>)	38	<i>Philerope</i> , Butler (<i>Heteronympha</i>)	100
Godart (<i>Satyrus</i>)	38	Butler (<i>Hipparchioides</i>)	100
Westwood (<i>Neonympha</i>) ...	39	Westwood (<i>Lasiommata</i>) 100, 166	
<i>Periphas</i> , Godart (<i>Satyrus</i>)	18	<i>Phileus</i> , Godart (<i>Satyrus</i>)	42
<i>Periphas</i> , Butler (<i>Euptychia</i>)	18	Philippa, Butler (<i>Euptychia</i>)	30
<i>perita</i> , Hewitson (<i>Lymanopoda</i>) ...	170	<i>Philis</i> , Cramer (<i>Papilio</i>)	101
<i>Perperna</i> , Hewitson (<i>Pronophila</i>)	173	<i>Philoctetes</i> , Clerck (<i>Papilio</i>)	107
<i>Perperna</i> , Butler (<i>Pedaliodes</i>) ...	173	<i>Philoctetes</i> , E. Doubleday (<i>Antir-</i>	
Felder (<i>Pronophila</i>)	173	rhæa)	107
<i>Persephone</i> , Hübner (<i>Papilio</i>) ...	55	Godart (<i>Satyrus</i>)	107
<i>Persephone</i> , Butler (<i>Hipparchia</i>)	55	Westwood (<i>Hætera</i>)	107
Esper (Pap. Nymph. Gemm.)	78	<i>Philoctetis</i> , Hübner (<i>Pieris</i>)	107
* <i>Perseus</i> , Fabricius (<i>Papilio</i>)	133	<i>Philomela</i> , Hübner (<i>Ypthima</i>) ...	152
<i>Peruda</i> , Hewitson (<i>Pronophila</i>) ...	179	* <i>Philomela</i> , Esper (Pap. Nymph.	
<i>Peruda</i> , Butler (<i>Pedaliodes</i>)	179	Gemm.)	82
<i>petrosus</i> , De Prunner (<i>Papilio</i>) ...	77	Hübner (<i>Papilio</i>)	82
<i>Peucestas</i> , Hewitson (<i>Pronophila</i>)	177	<i>Philopæmen</i> , Felder (<i>Antirrhæa</i>)	108
<i>Peucestas</i> , Butler (<i>Pedaliodes</i>) ...	177	<i>Philotera</i> , Hewitson (<i>Pronophila</i>)	179
<i>Phæa</i> , Hewitson (<i>Pronophila</i>) ...	178	<i>Philotera</i> , Butler (<i>Pedaliodes</i>) ...	179
<i>Phæa</i> , Butler (<i>Pedaliodes</i>)	178	<i>Philoxena</i> , Hübner (<i>Cæno-</i>	
<i>Phædra</i> , Esper (Pap. Nymph.		nympha)	46
Gemm.)	69	* <i>Philoxenus</i> , Esper (Pap. pleb.	
Godart (<i>Satyrus</i>)	62	rural.)	46
Linnaeus (<i>Papilio</i>)	61	Butler (<i>Cænonympha</i>)	46
Schrenck (<i>Maniola</i>)	62	<i>Phineus</i> , Butler (<i>Euptychia</i>)	26
Stephens (<i>Hipparchia</i>)	62	<i>Phocion</i> , Fabricius (<i>Papilio</i>)	37
<i>Phæasana</i> , Hewitson (<i>Pronophila</i>)	178	<i>Phocion</i> , Godart (<i>Satyrus</i>)	37
<i>Phalæsia</i> , Hewitson (<i>Pronophila</i>)	182	Westwood (<i>Neonympha</i>) ...	37
<i>Phalæsia</i> , Butler (<i>Lasiophila</i>) ...	182	<i>Phœnicusa</i> , Hewitson (<i>Pronophila</i>)	178
<i>Phanaræa</i> , Hewitson (<i>Pronophila</i>)	179	<i>Phœnicusa</i> , Butler (<i>Pedaliodes</i>) ...	178
<i>Phanaræa</i> , Butler (<i>Pedaliodes</i>) ...	179	<i>Phœnissa</i> , Hewitson (<i>Pronophila</i>)	177
<i>Phanias</i> , Hewitson (<i>Pronophila</i>)	174	<i>Phœnissa</i> , Butler (<i>Pedaliodes</i>) ...	177
<i>Phanias</i> , Butler (<i>Pedaliodes</i>)	174	<i>Phorcys</i> , Freyer (Pap. Hip-	
<i>Pharella</i> , Butler (<i>Euptychia</i>)	24	parchia)	91
<i>Phares</i> , Godart (<i>Satyrus</i>)	24	<i>Phoronea</i> , Hewitson (<i>Pronophila</i>)	183
<i>Phares</i> , Butler (<i>Euptychia</i>) ...	24	<i>Phoronea</i> , Butler (<i>Dædalma</i>)	183
Westwood (<i>Neonympha</i>) ...	24	<i>Phronius</i> , Godart (<i>Satyrus</i>)	20
<i>Pharte</i> , Esper (<i>Papilio</i>)	74	<i>Phronius</i> , Butler (<i>Euptychia</i>) ...	20

	Page
<i>Phronius</i> , Westwood (Neonympha) ..	20
<i>Phryne</i> , <i>Pallas</i> (Pap. Nymph. Gemm.) ..	48
<i>Phryne</i> , Borkhausen (Papilio) ..	48
<i>E. Doubledayi</i> (Cænonympha) ..	48
<i>Esper</i> (Pap. pleb. rural.) ..	48
<i>Herrich-Schäffer</i> (Satyrus) ..	49
<i>Phryneus</i> , <i>Fabricius</i> (Papilio) ..	49
<i>Godart</i> (Satyrus) ..	49
<i>Phyllalia</i> , <i>Hewitson</i> (Pronophila) ..	175
<i>Phyllalia</i> , <i>Butler</i> (Pedaliodes) ..	175
<i>Physcoa</i> , <i>Hewitson</i> (Pronophila) ..	177
<i>Physcoa</i> , <i>Butler</i> (Pedaliodes) ..	177
<i>picea</i> , <i>Butler</i> (Euptychia) ..	28
<i>Piera</i> , <i>Linnaeus</i> (Papilio) ..	102
<i>Piera</i> , <i>Boisduval</i> (Hætera) ..	102
<i>Cramer</i> (Papilio) ..	102
<i>Godart</i> (Satyrus) ..	102
<i>Hübner</i> (Oreas & Cithærias) ..	102
<i>Westwood</i> (Hætera) ..	102
<i>PIERRELLA</i> , <i>Westwood</i> ..	103
<i>Pieria</i> , <i>Butler</i> (Euptychia) ..	17
<i>Pierra</i> , <i>Sulzer</i> (Papilio) ..	102
<i>pignerator</i> , <i>Butler</i> (Amphidecta) ..	7
<i>pilata</i> , <i>Butler</i> (Euptychia) ..	29
<i>Piletha</i> , <i>Hewitson</i> (Pronophila) ..	176
<i>Piletha</i> , <i>Butler</i> (Pedaliodes) ..	176
<i>pilosella</i> , <i>Fabricius</i> (Papilio) ..	69
<i>Pimpla</i> , <i>Felder</i> (Neonympha) ..	20
<i>Felder</i> (Satyrus) ..	60
<i>Pimpla</i> , <i>Butler</i> (Euptychia) ..	20
<i>Pimplea</i> , <i>Erichson</i> (Cynthia) ..	109
<i>Pimplea</i> , <i>Felder</i> (Zethera) ..	109
<i>Westwood</i> (Diadema ?) ..	109
<i>pirata</i> , <i>Esper</i> (Pap. Nymph. Gemm.) ..	54
<i>Hübner</i> (Papilio) ..	54
<i>Pirene</i> , <i>Esper</i> (Pap. Nymph. Gemm.) ..	78
<i>Pirene</i> , <i>Butler</i> (Erebia) ..	78
<i>Hübner</i> (Papilio) ..	84
<i>Pisidice</i> , <i>Klug</i> (Hipparchia) ..	57
<i>Pisidice</i> , <i>Westwood</i> (Satyrus) ..	57
<i>Pisonia</i> , <i>Hewitson</i> (Pronophila) ..	173, 174
<i>*Pitho</i> , <i>Hübner</i> (Papilio) ..	80
<i>Plesaire</i> , <i>Bellier</i> (Arge) ..	157
<i>Plotina</i> , <i>Hewitson</i> (Pronophila) ..	178
<i>Plotina</i> , <i>Butler</i> (Pedaliodes) ..	178
<i>plumbeola</i> , <i>Butler</i> (Tetraphlebia ?) ..	95
<i>Pluto</i> , <i>Esper</i> (Pap. Nymph. Gemm.) ..	77
<i>*Podarce</i> , <i>Herrich-Schäffer</i> (Satyrus) ..	60
<i>Poesia</i> , <i>Hewitson</i> (Pronophila) ..	172
<i>Poesia</i> , <i>Butler</i> (Pedaliodes) ..	172
<i>Poliozona</i> , <i>Felder</i> (Epinephele) ..	67
<i>Polixenes</i> , <i>Fabricius</i> (Papilio) ..	164
<i>Westwood</i> (Neonympha) ..	164

	Page
<i>*Pollux</i> , <i>Esper</i> (Pap. Nymph. Gemm.) ..	87
<i>Poltya</i> , <i>Prittwitz</i> (Neonympha) ..	24
<i>Poltya</i> , <i>Butler</i> (Euptychia) ..	24
<i>Polusca</i> , <i>Hewitson</i> (Pronophila) ..	176
<i>Polusca</i> , <i>Butler</i> (Pedaliodes) ..	176
<i>Polydama</i> , <i>Haworth</i> (Papilio) ..	46
<i>Polydecta</i> , <i>Cramer</i> (Papilio) ..	135
<i>Polydecta</i> , <i>Godart</i> (Satyrus) ..	135
<i>Westwood</i> (Mycælesis) ..	132, 135
<i>*Polymeda</i> , <i>Jermyns</i> (Papilio) ..	46
<i>Butler</i> (Cænonympha) ..	46
<i>Scooli</i> (Papilio) ..	63
<i>Polyphemus</i> , <i>Butler</i> (Euptychia) ..	32
<i>Pompilia</i> , <i>Felder</i> (Neonympha) ..	25
<i>Pompilia</i> , <i>Butler</i> (Euptychia) ..	25
<i>*Pontica</i> , <i>Freyer</i> (Pap. Hipparchia) ..	51
<i>Westwood</i> (Satyrus) ..	51
<i>Porina</i> , <i>Hewitson</i> (Pronophila) ..	177
<i>Porina</i> , <i>Butler</i> (Pedaliodes) ..	177
<i>Porphyria</i> , <i>Felder</i> (Pronophila) ..	175
<i>Porphyria</i> , <i>Butler</i> (Pedaliodes) ..	175
<i>Porsenna</i> , <i>Hewitson</i> (Pronophila) ..	184
<i>Portlandia</i> , <i>Fabricius</i> (Papilio) ..	114
<i>Portlandia</i> , <i>Boisduval</i> (Satyrus) ..	114
<i>Westwood</i> (Debis) ..	164
<i>Præneste</i> , <i>Hewitson</i> (Pronophila) ..	181
<i>Præneste</i> , <i>Butler</i> (Lasiophila) ..	181
<i>*Priauri</i> , <i>Pierret</i> (Satyrus) ..	55
<i>*Procida</i> , <i>Herbst</i> (Papilio) ..	153
<i>Boisduval</i> (Arge) ..	153
<i>Freyer</i> (Pap. Hipparchia) ..	153
<i>Prochyta</i> , <i>Hewitson</i> (Pronophila) ..	181
<i>Prochyta</i> , <i>Butler</i> (Oxoeschistus) ..	181
<i>Proerna</i> , <i>Hewitson</i> (Pronophila) ..	173
<i>Proerna</i> , <i>Butler</i> (Pedaliodes) ..	173
<i>pronax</i> , <i>Hewitson</i> (Pronophila) ..	180
<i>pronax</i> , <i>Butler</i> (Oxoeschistus) ..	180
<i>Pronoe</i> , <i>Borkhausen</i> (Papilio) ..	81
<i>Esper</i> (Pap. Nymph. Gemm.) ..	81
<i>Freyer</i> (Pap. Hipparchia) ..	80
<i>Ochsenheimer</i> (Papilio) ..	80
<i>PRONOPHILA</i> , <i>Westwood</i> ..	184
<i>Pronophila</i> , <i>Butler</i> (Euptychia) ..	26
<i>Felder</i> (Steroma) ..	172
<i>Propylea</i> , <i>Hewitson</i> (Pronophila) ..	180
<i>Propylea</i> , <i>Butler</i> (Oxoeschistus) ..	180
<i>Proserpina</i> , <i>Denis</i> (Papilio) ..	50
<i>Proserpina</i> , <i>Butler</i> (Hipparchia) ..	50
<i>Cyrellus</i> (Papilio) ..	61
<i>Esper</i> (Pap. Nymph. Gemm.) ..	50
<i>Westwood</i> (Satyrus) ..	50
<i>Prosymna</i> , <i>Hewitson</i> (Pronophila) ..	182
<i>Prosymna</i> , <i>Butler</i> (Lasiophila) ..	182
<i>Protogenia</i> , <i>Hewitson</i> (Pronophila) ..	180
<i>Protogenia</i> , <i>Butler</i> (Oxoeschistus) ..	180
<i>Prytanis</i> , <i>Hewitson</i> (Pronophila) ..	177
<i>Prytanis</i> , <i>Butler</i> (Pedaliodes) ..	177
<i>PSEUDONYMPHA</i> , <i>Wallengren</i> ..	93

	Page		Page
* <i>Psodea</i> , Hübner (Papilio).....	90	<i>Rhamnusia</i> , Westwood (Satyrus)...	65
Freyer (Pap. Hipparchia) ...	90	<i>RHAPHICERA</i> , Butler.....	158
Godart (Satyrus).....	90	<i>Rhea</i> , Fabricius (Papilio).....	104
Herrich-Schäffer (Erebia) ...	90	<i>Rhea</i> , Butler (Pierella)	104
<i>Psyche</i> , Godart (Satyrus)	157	Hübner (Pieris).....	104
Hübner (Papilio).....	157	<i>Ridingsii</i> , Edwards (Satyrus).....	59
Westwood (Arge)	157	<i>Ridingsii</i> , Butler (Hipparchia?) ..	59
PTYCHANDRA, Felder	187	<i>Rinda</i> , Ménétries (Cænonympha) ..	43
Puerta, Westwood (Pronophila)...	180	<i>Rohria</i> , Fabricius (Papilio)	116
Puerta, Butler (Oxeoschistus) ...	180	<i>Rohria</i> , Butler (Lethe)	116
Pulaha, Moore (Lasiommata?) ...	112	Godart (Satyrus).....	116
<i>Pulaha</i> , Butler (Neope) 112, 113		Westwood (Debis).....	116
Moore (Enope)	112	<i>Rossii</i> , Curtis (Hipparchia)	89
pulchella, Felder (Epinephele) ...	65	<i>Rossii</i> , Westwood (Erebia).....	89
pulchra, Felder (Epinephele).....	65	<i>Roxandra</i> , Nordmann (Pararge) ..	122
pumilus, Felder (Chionobas).....	165	<i>Roxane</i> , Felder (Epinephele).....	69
<i>pumilus</i> , Butler (Ceneis).....	165	<i>Roxelana</i> , Cramer (Papilio)	121
Purana, Felder (Debis)	118	<i>Roxelana</i> , Butler (Pararge)	121
<i>Purana</i> , Butler (Lethe)	118	E. Doubleday (Lasiommata) ..	121
pusilla, Felder (Homœonympha) ..	95	Freyer (Pap. Hipparchia)	121
<i>Pylarge</i> , Hübner (Papilio).....	40	Godart (Satyrus).....	121
<i>Pylas</i> , Hewitson (Pronophila) ...	177	Hübner (Dira).....	121
<i>Pylas</i> , Butler (Pedaliodes).....	177	* <i>Runeka</i> , Moore (Mycalesis).....	139
<i>Pyraemon</i> , Butler (Euptychia)....	37	<i>rustica</i> , Butler (Euptychia)	32
* <i>Pyrenaica</i> , Herrich-Schäffer			
(Erebia)	75		S.
<i>Pyrene</i> , Freyer (Papilio).....	79	<i>Saadi</i> , Kollar (Satyrus)	48
Westwood (Erebia).....	78	<i>Saadi</i> , Butler (Cænonympha) ...	48
<i>Pyrrha</i> , Denis (Papilio)	76	<i>Sabacus</i> , Trimen (Erebia)	93, 94
<i>Pyrrha</i> , Butler (Oreina).....	76	<i>Sabacus</i> , Butler (Pseudonympha) ..	93
Freyer (Pap. Hipparchia) ...	76	Fabricius (Papilio).....	40
Godart (Satyrus).....	76	<i>Sabina</i> , Felder (Neonympha).....	30
Hübner (Papilio).....	76	<i>Sabina</i> , Butler (Euptychia).....	30
Westwood (Erebia).....	76, 77	<i>Saitza</i> , Hewitson (Mycalesis).....	128
		<i>Saga</i> , Butler (Mycalesis).....	130
Q.		<i>Saitia</i> , Hewitson (Zipaetis)	98
<i>Quantius</i> , Godart (Satyrus)	31	<i>Sakra</i> , Moore (Ypthima)	149
<i>Quantius</i> , Butler (Euptychia) ...	31	<i>Salome</i> , Fabricius (Papilio)	70
Westwood (Neonympha) ...	31	<i>Salvini</i> , Butler (Euptychia)	37
		* <i>Samba</i> , Moore (Mycalesis)	137
R.		<i>Samio</i> , Hewitson (Debis).....	118
* <i>Rasaces</i> , Hewitson (Mycalesis) ..	132	<i>Samio</i> , Butler (Lethe).....	118
RAGADIA, Westwood.....	158	<i>Samius</i> , Westwood (Lymanopoda) ..	169
<i>Rebecca</i> , Fabricius (Papilio)	9	<i>Sanaos</i> , Hewitson (Mycalesis).....	131
Godart (Satyrus).....	10	<i>Sanatana</i> , Moore (Mycalesis)	138
Westwood (Taygetis).....	10	<i>Saraswati</i> , Kollar (Satyrus).....	49
<i>Remulia</i> , Cramer (Papilio).....	143	<i>Saraswati</i> , Butler (Aulocera).....	49
<i>Remulia</i> , Westwood (Mycalesis) ..	143	Moore (Satyrus)	49
<i>renata</i> , Cramer (Papilio).....	20	<i>Sareba</i> , Hewitson (Corades)	186
<i>renata</i> , Butler (Euptychia).....	20	<i>Satricus</i> , Hewitson (Lasiommata) ..	158
Herrich-Schäffer (Neonympha)	20	<i>Satricus</i> , Butler (Rhaphiocera) ...	158
Westwood (Mycalesis?).....	20	<i>Saturnus</i> , Butler (Euptychia).....	27
<i>Rhacotis</i> , Hewitson (Mycalesis?) ..	130	<i>satyrina</i> , Bates (Taygetis)	13
<i>Rhamnusia</i> , Freyer (Pap. Hipparchia)	65	<i>Satyrion</i> , Esper (Pap. Dan. fest.) ..	42
<i>Rhamnusia</i> , Butler (Epinephele) ..	65	<i>Satyrion</i> , Freyer (Pap. Hipparchia)	42
Herrich-Schäffer (Erebia) ...	65	Herrich-Schäffer (Satyrus)....	42
		Ochsenheimer (Papilio)	42
		<i>SATYRUS</i> , Westwood (subgenus) ...	59
		<i>Saundersii</i> , Butler (Euptychia) ...	38

	Page
<i>Scæa</i> , Hübner (Papilio)	84
<i>scanda</i> , Moore (Debis)	117
<i>Kollar</i> (Erebia)	96
<i>scanda</i> , Butler (Callerebia)	96
(Lethe)	117
<i>Schakra</i> , Kollar (Satyrus)	126
<i>Schakra</i> , Butler (Amecera)	126
Westwood (Lasiommata) ...	126
<i>Schrenckii</i> , Ménétries (Prono- phila)	122
<i>Schrenckii</i> , Bremer (Nemophila)	122
Butler (Pararge)	122
<i>Schrenckii</i> , Ménétries (Prono- phila)	122
<i>Sciathis</i> , Hewitson (Mycælesis) ...	130
<i>Scipio</i> , Boisduval (Erebia)	78
Duponchel (Satyrus)	78
<i>Sculda</i> , Eversmann (Hipparchia)	163
<i>Sculda</i> , Herrich-Schäffer (Chio- nobas)	163
<i>Scoylax</i> , Hewitson (Zipaetis)	98
<i>Scylla</i> , Butler (Aulocera)	60
* <i>Sedakovii</i> , Eversmann (Hip- parchia)	83
Herrich-Schäffer (Erebia) ...	81
Westwood (Erebia)	83
* <i>Segonax</i> , Hewitson (Debis)	113
<i>Semele</i> , Linnaeus (Papilio)	55
<i>Semele</i> , Boisduval (Satyrus)	56
Esper (Pap. Nymph. Gemm.)	55
Godart (Satyrus)	56
Herrich-Schäffer (Satyrus) ...	56
Hübner (Papilio)	56
Stephens (Hipparchia)	56
<i>semidea</i> , Say (Hipparchia)	163
<i>semidea</i> , Butler (Cneis)	163
Edwards (Chionobas)	164
Morris (Cænonympha)	164
<i>Sempera</i> , Felder (Ypthima)	151
<i>Semperi</i> , Butler (Mycælesis)	137
<i>sepulta</i> , Boisduval (Cyllo)	189
<i>sepulta</i> , Lefebvre (Vanessa)	189
<i>sericeella</i> , Bates (Euptychia)	32
<i>Servatius</i> , Godart (Satyrus)	128
<i>Servilia</i> , Wallengren (Neomensas)	66
<i>Servilia</i> , Butler (Epinephele) ...	66
<i>Sesara</i> , Hewitson (Xois)	147
<i>Shakra</i> , Butler (Amecera)	126
<i>Shiva</i> , Boisduval (Satyrus)	143
<i>Shiva</i> , Butler (Mycælesis)	143
Westwood (Cænonympha)	143
<i>Sicelis</i> , Hewitson (Debis)	118
<i>Sicelis</i> , Butler (Lethe)	118
<i>Sidonis</i> , Hewitson (Debis)	116
<i>Sidonis</i> , Butler (Lethe)	116
<i>silvestris</i> , Edwards (Satyrus)	63
<i>similis</i> , Butler (Euptychia)	17
<i>simplex</i> , Butler (Oxoecisthus) ...	180
<i>Singa</i> , Boisduval (Satyrus)	166
Westwood (Lasiommata) ...	166

	Page
<i>Sinorix</i> , Hewitson (Debis)	118
<i>Sinorix</i> , Butler (Lethe)	118
<i>Sirius</i> , Fabricius (Papilio)	144
<i>Sirius</i> , Butler (Mycælesis)	144
Godart (Satyrus)	145
Westwood (Cænonympha)	145
<i>Sita</i> , Felder (Neorina)	111
* <i>Solandra</i> , Fabricius (Papilio) ...	3
Boisduval (Cyllo)	3
Butler (Melanitis)	3
<i>Sophia</i> , Acerbi (Papilio)	88
<i>Sosybius</i> , Fabricius (Papilio)	22
<i>Sosybius</i> , Butler (Euptychia) ...	22
Godart (Satyrus)	22
Westwood (Neonympha)	22
<i>Statilinus</i> , Hufnagel (Papilio) ...	56
<i>Statilinus</i> , Freyer (Pap. Hippar- chia)	56
Herrich-Schäffer (Satyrus) ...	56
<i>Stellera</i> , Eschscholtz (Hippar- chia)	150
<i>Stellera</i> , Butler (Ypthima)	150
Hewitson (Ypthima)	151
STEROMA, Westwood	172
<i>Sthenele</i> , Boisduval (Satyrus) ...	63
<i>Sthennyo</i> , Grashin (Erebia)	87
<i>Stheno</i> , Hübner (Papilio)	89
<i>Stirius</i> , Godart (Satyrus)	92
<i>straminea</i> , Butler (Euptychia) ...	19
<i>Stubbendorffii</i> , Ménétries (Ere- bia)	74
* <i>Stygne</i> , Godart (Satyrus)	78
Fischer (Hipparchia)	81
Freyer (Pap. Hipparchia) ...	84
Herrich-Schäffer (Erebia) ...	78
Hübner (Papilio)	84
STYGNUS, Felder	96
<i>Styz</i> , Freyer (Pap. Hipparchia)	80
<i>subhyalina</i> , Curtis (Hipparchia)	164
<i>subhyalina</i> , Butler (Cneis)	164
Westwood (Erebia)	164
<i>Sudra</i> , Felder (Mycælesis)	140
<i>Sunbecca</i> , Eversmann (Hippar- chia)	48
<i>Sunbecca</i> , Herrich-Schäffer (Sa- tyrus)	48
Westwood (Cænonympha) ...	48
<i>superba</i> , Butler (Steroma)	172
<i>Sura</i> , Hewitson (Zophoessa)	108
<i>Suradeva</i> , Moore (Melanitis)	6
<i>Suradeva</i> , Butler (Cyllogenes) ...	6
<i>Suyudana</i> , Moore (Melanitis) ...	4
<i>Swaha</i> , Kollar (Satyrus)	49
<i>Syllius</i> , Herbst (Papilio)	156
<i>Sylvia</i> , Bates (Taygetis)	11
<i>Sylvina</i> , Felder (Neonympha) ...	25
<i>Sylvina</i> , Butler (Euptychia) ...	25
<i>Synclimene</i> , Hübner (Epinephele)	121
<i>Syrcis</i> , Hewitson (Debis)	118
<i>Syrcis</i> , Butler (Lethe)	118

	Page		Page
T.			
<i>Tabitha</i> , Fabricius (Papilio)	133	<i>Tircis</i> , Herrich-Schäffer (Phryne)	49
<i>Tagala</i> , Felder (Mycalesis)	143	Zeller (Triphysa)	49
* <i>Taitensis</i> , Felder (Cyllo)	2	<i>TISIPHONE</i> , Westwood	71
Butler (Melanitis)	2	<i>Tisiphone</i> , Hübner (Panopea) ...	168
<i>Tamatava</i> , Boisduval (Satyrus) ...	94	<i>Tisiphone</i> , Boisduval (Euterpe) ...	168
<i>Tamatava</i> , Butler (Pararge)	94	E. Doubleday (Eteona) ..	168
E. Doubleday (Erebia?)	94	Fabricius (Papilio)	71
Westwood (Erebia)	94	* <i>Tissiphone</i> , Esper (Pap. Nymph.	
<i>Tarpeia</i> , Esper (Pap. Nymph.		Gemm.)	77
Gemm.)	161	* <i>Titea</i> , Klug (Hipparchia)	154
<i>Tarpeia</i> , Ochsenheimer (Papilio) ..	161	Boisduval (Arge)	154
<i>Tarpeius</i> , Duponchel (Satyrus) ...	161	<i>Tithonius</i> , Linnaeus (Papilio)	69
<i>Tarpeja</i> , Borkhausen (Papilio) ...	161	<i>Tithonus</i> , Stephens (Hipparchia) ..	49
Freyer (Pap. Hipparchia) ...	161	Westwood (Satyrus)	69
Herrich-Schäffer (Chionobas) ..	161	<i>Tolumnia</i> , Cramer (Papilio)	34
Hübner (Eumenis)	161	<i>Tolumnia</i> , Godart (Satyrus) ...	33, 34
<i>Tauropolis</i> , Hewitson (Pronophila) ..	180	Hübner (Euptychia)	34
<i>Tauropolis</i> , Butler (Pedaliodes) ...	180	tricolor, Hewitson (Euptychia) ...	34
<i>Taygete</i> , Hübner (Cneis)	164	<i>Trimenii</i> , Butler (Pseudonympha) ..	94
<i>Taygete</i> , Herrich-Schäffer (Chiono-		<i>TRIPHYSA</i> , Zeller	48
bas)	164, 165	<i>tristis</i> , Bremer (Erebia)	92
<i>Taygetina</i> , Butler (Anchiphlebia) ..	107	* <i>tristis</i> , Felder (Cyllo)	2
<i>TAYGETIS</i> , Hübner	7	Butler (Melanitis)	2
<i>Teneates</i> , Ménestries (Hipparchia) ..	154	Guérin (Argynnis & Satyrus) ..	68
<i>tenebrosus</i> , Blanchard (Faunus) ..	8	Herrich-Schäffer (Erebia) ...	77
<i>terminus</i> , Fabricius (Papilio)	143	Westwood (Erebia)	68
<i>terminus</i> , Godart (Satyrus)	144	<i>Tullia</i> , Hübner (Papilio)	46
Westwood (Mycalesis?)	144	Müller (Pap. Nymph.)	46
<i>terrestris</i> , Butler (Euptychia) ...	16	<i>Tyndarellus</i> , Herbst (Papilio) ...	85
<i>TETRAPHLEBIA</i> , Felder	95	<i>Tyndarus</i> , Esper (Pap. Nymph.	
<i>Thamyra</i> , Cramer (Papilio)	12	Gemm.)	85
<i>Thamyra</i> , Hübner (Oreas & Tay-		<i>Tyndarus</i> , Freyer (Papilio)	86
getis)	12	Ochsenheimer (Papilio)	86
<i>Theano</i> , Eversmann (Hipp. Ere-		<i>Typhia</i> , Hewitson (Oressinoma) ..	153
bia)	74	<i>Typhon</i> , Haworth (Papilio)	46
Tauscher (Papilio)	74		
<i>Thelebe</i> , Hewitson (Pronophila) ..	184	U.	
<i>Thelephassa</i> , Hübner (Eumenis) ..	51	<i>Uhleri</i> , Reakirt (Chionobas)	163
<i>Thelephassa</i> , Butler (Hipparchia) ..	51	<i>Uhleri</i> , Butler (Cneis)	163
Herrich-Schäffer (Satyrus) ...	52	<i>Ulema</i> , Hewitson (Corades)	186
Klug (Hipparchia)	52	<i>undulata</i> , Butler (Euptychia)	23
<i>Themis</i> , Butler (Euptychia)	16	<i>Urania</i> , Butler (Euptychia)	29
<i>Thetis</i> , Hübner (Papilio)	156	<i>Urda</i> , Eversmann (Hipparchia) ..	162
<i>Thyrsis</i> , Freyer (Pap. Hipparch-		<i>Urda</i> , Butler (Cneis)	162
chia)	44	Herrich-Schäffer (Chionobas) ..	162
<i>Thyrsis</i> , Herrich-Schäffer (Saty-		<i>usitata</i> , Butler (Euptychia)	17
rus)	44		
Westwood (Canonympha) ...	44	V.	
<i>Tigellus</i> , Boisduval (Satyrus) ...	125	<i>Valdivia</i> , Felder (Epinephela) ...	67
Bonelli (Papilio)	125	<i>Valentina</i> , Cramer (Papilio)	13
Butler (Amecera)	125	<i>Valentina</i> , Butler (Taygetis)	13
Freyer (Pap. Hipparchia) ...	125	Godart (Satyrus)	13
Westwood (Lasioommata) ...	125	<i>Vamana</i> , Moore (Melanitis)	3
<i>Tiphon</i> , Schneider (Papilio)	40	vanessoides, Blanchard (Elina) ...	64
Schränk (Maniola)	46	<i>Varaha</i> , Moore (Melanitis)	4
* <i>Tircis</i> , Godart (Satyrus)	120	<i>variabilis</i> , Butler (Euptychia) ...	19
Butler (Lasioommata)	120	<i>vastata</i> , Butler (Euptychia)	31
Cramer (Papilio)	49	<i>velutina</i> , Felder (Cyllo)	6

	Page
<i>vehutina</i> , Butler (Hipio)	6
<i>venosa</i> , Butler (Lymanopoda) ...	171
<i>Verma</i> , Kollar (Satyrus).....	116
<i>Verma</i> , Butler (Lethe)	116
Westwood (Debis)	116
<i>Vesagus</i> , Hewitson (Erebia)	86
<i>Vesper</i> , Butler (Euptychia)	22
<i>Vesta</i> , Butler (Euptychia)	27
<i>vestigata</i> , Butler (Euptychia) ...	20
<i>vicinus</i> , Felder (Cœlites)	111
<i>vidua</i> , Müller (Papilio).....	64
<i>Vindhya</i> , Felder (Debis).....	119
<i>Vindhya</i> , Butler (Lethe).....	119
<i>Virbius</i> , Freyer (Pap. Hipparchia)	61
<i>Virbius</i> , Herrich-Schäffer (Satyrus)	61
<i>Virgilia</i> , Cramer (Papilio).....	9
<i>Virgilia</i> , Hübner (Taygetis)	9
* <i>Visala</i> , Moore (Mycalesis)	133
<i>Visrava</i> , Moore (Debis)	116
<i>Visrava</i> , Butler (Lethe)	116
<i>vulgaris</i> , Butler (Mycalesis)	130

W.

<i>Wagneri</i> , Herrich-Schäffer (Satyrus)	70
<i>Wagneri</i> , Butler (Epinephele) ...	70
E. Doubleday (Hipparchia)	70
<i>Wanga</i> , Bremer (Erebia)	92
<i>Werang</i> , Lang (Aulocera)	188
<i>Westwoodii</i> , Butler (Euptychia)	27
<i>Whitelyi</i> , Butler (Lethe)	117
<i>Williamsianus</i> , Butler (Argyro-phorus).....	159

X.

<i>Xeneas</i> , Hewitson (Mycalesis) ...	131
<i>Xenia</i> , Freyer (Pap. Hipparchia)	156
<i>XENICA</i> , Westwood (subgenus)...	70

	Page
<i>Xicaque</i> , <i>Reakirt</i> (Neonympha)	39, 99
<i>Xicaque</i> , Butler (Euptychia)	39
Butler (Paramecera)	99
<i>Xiphia</i> , Fabricius (Papilio)	121
<i>Boisduval</i> (Satyrus).....	125
<i>Xiphia</i> , Butler (Amecera)	125
Butler (Pararge)	121
Duponchel (Satyrus)	120
E. Doubleday (Lasiommata)	121
Godart (Satyrus)	121
Herrich-Schäffer (Satyrus)...	120
<i>Xois</i> , Hewitson	147
<i>Xyphia</i> , Hübner (Pararge).....	121

Y.


<i>Yama</i> , Moore (Zophoessa)	109
<i>ypthima</i> , Felder (Neonympha)	26
<i>ypthima</i> , Butler (Euptychia) ...	26
<i>YPTHIMA</i> , Westwood.....	148
<i>Ypthima</i> , Hübner (Taygetis).....	9

Z.

* <i>Zacheus</i> , Fabricius (Papilio) ...	145
Godart (Satyrus).....	145
Westwood (Lasiommata) ...	145
<i>Zangis</i> , Fabricius (Papilio)	97
<i>Zangis</i> , Godart (Satyrus)	97
Hewitson (Calisto)	97
Hübner (Hipparchia).....	97
<i>Zapatoza</i> , Westwood (Pronophila)	181
<i>Zapatoza</i> , Butler (Lasiophila) ...	181
<i>Zelinde</i> , Hübner (Oreas & Tisiphone)	70
<i>ZETHERA</i> , Felder	109
<i>Zilia</i> , Borkhausen (Papilio)	87
<i>ZIPAETIS</i> , Hewitson	98
<i>ZOPHOESSA</i> , Westwood	108
<i>Zophyrus</i> , Westwood (Satyrus) ...	132
<i>Zopyrus</i> , Kollar (Satyrus)	132

THE END.






A.G. Butler del. et lith. Feb. & March 1868.

W. West, imp.

- | | |
|-------------------------------|--------|
| 1. <i>Taygetis excavata</i> , | Butler |
| 2. " <i>erubescens</i> | " |
| 3. " <i>armillata</i> | " |
| 4. <i>Euptychia rustica</i> . | " |

- | | |
|--------------------------------|--------|
| 5. <i>Taygetis Euptychidia</i> | Butler |
| 6. " <i>nymphosa</i> | " |
| 7. " <i>nympha</i> | " |

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1




A.G. Butler del et lith, March 1868.

W. West imp.

- | | | | |
|---------------------------------|----------|-----------------------------------|--------|
| 1. <i>Melanitis Gnophodes</i> , | Butler. | 7. <i>Erebia Hispania</i> , | Butler |
| 2. <i>Æneis gigas</i> , | " | 8. " <i>fasciata</i> , | " |
| 3. <i>Lethe arcuata</i> , | " | 9. " <i>episnodea</i> , | " |
| 4. <i>Erites elegans</i> , | " | 10. <i>Æneis assimilis</i> , | " |
| 5. <i>Paramacera xicaque</i> , | Reakirt. | 11. <i>Tetraphlebia plumbeola</i> | " |
| 6. <i>Erebia mantoides</i> , | Butler. | 12. <i>Leptoneura cassina</i> , | " |






A. G. Butler del. et lith. April. 1868.

W. West imp.

- | | | | |
|----------------------------------|---------|------------------------------|--------|
| 1. <i>Mycalesis suga</i> | Butler. | 8. <i>Mycalesis angulosa</i> | Butler |
| 2. " <i>vulgaris</i> | " | 9. " <i>mundosa</i> | " |
| 3. <i>Epinephele pallescens?</i> | " | 10. " <i>Madjicosu</i> | " |
| 4. <i>Mycalesis junonia</i> | " | 11. " <i>æthiops</i> | " |
| 5. " <i>Felderi</i> | " | 12. " <i>justinella</i> | " |
| 6. " <i>auricruda</i> | " | 13. " <i>fraternu</i> | " |
| 7. " <i>Semper</i> | " | 14. " <i>malsaridu</i> | " |






A. G. Butler del et lith. May 1868.

W. West, imp.

- | | |
|--------------------------------------|--------|
| 1. <i>Argyrophorus Williamsianus</i> | Butler |
| 2. <i>Oxeoschistus simplex</i> | " |
| 3. <i>Lymanopoda ferruginosa</i> | " |
| 4. <i>Lasiophila Hewitsonia</i> | " |
| 5. <i>Lymanopoda venosa</i> | " |

- | | |
|--------------------------------|---------|
| 6. <i>Lymanopoda Acraeida</i> | Butler. |
| 7. <i>Pedaliodes angularis</i> | " |
| 8. " <i>granulata</i> | " |
| 9. <i>Hypocysta antirius</i> | " |
| 10. <i>Pronophila obscura</i> | " |





A. G. Butler del. et lith. May 1868.

W. West, imp.

1. *Hipio Crameri* Butler.
 2. *Anchiphlebia Thygetina* "
 3. " *archaea* Hübner.
 4. *Ctenonympha marginalis* Butler.

5. *Euptychia perfuscata* Butler.
 6. *Lasiophila orbifera* "
 7. *Steroma superba* "
 8. *Erites argentina* "





